



Camel tick species distribution in Saudi Arabia and United Arab Emirates using MaxEnt modelling

Research Article

Cite this article: Perveen N, Muzaffar SB, Jaradat A, Sparagano OA, Willingham AL (2024). Camel tick species distribution in Saudi Arabia and United Arab Emirates using MaxEnt modelling. *Parasitology* **151**, 1024–1034. <https://doi.org/10.1017/S0031182024001161>

Received: 4 June 2024
Revised: 19 August 2024
Accepted: 4 September 2024

Keywords:

camel tick; *Hyalomma dromedarii*; MaxEnt; modelling; Saudi Arabia; species distribution; UAE

Corresponding author:

Arve L. Willingham;
Email: awillingham@uaeu.ac.ae

Nighat Perveen^{1,2} , Sabir B. Muzaffar^{1,3} , Areej Jaradat¹ ,
Olivier A. Sparagano^{4,5} and Arve L. Willingham²

¹Department of Biology, College of Science, United Arab Emirates University, Al-Ain, UAE; ²Department of Veterinary Medicine, College of Agriculture and Veterinary Medicine, United Arab Emirates University, Al-Ain, UAE; ³Department of Science, The Natural History Museum, London, UK; ⁴Agricultural Sciences and Practice, Royal Agricultural University, Cirencester, UK and ⁵Department of Infectious Diseases and Public Health, Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong Kong, Kowloon, Hong Kong SAR, China

Abstract

Ticks are important vectors and reservoirs of pathogens causing zoonotic diseases in camels and other livestock, rodents and other small mammals, birds and humans. *Hyalomma dromedarii* is the most abundant tick species in Saudi Arabia and United Arab Emirates (UAE) affecting primarily camels, and to a lesser extent, other livestock. Species presence data, land use/landcover, elevation, slope and 19 bioclimatic variables were used to model current and future distribution of *H. dromedarii* ticks using maximum entropy species distribution modelling (MaxEnt.). The model highlighted areas in the northern, eastern and southwestern parts of the study area as highly suitable for ticks. Several variables including land use/land cover (LULC) (53.1%), precipitation of coldest quarter (Bio19) (21.8%), elevation (20.6%), isothermality (Bio3) (1.9%), mean diurnal range [mean of monthly (max temp – min temp)] (Bio2) (1.8%), slope (0.5%), precipitation, seasonality (Bio15) (0.2%) influenced habitat suitability of ticks, predicting high tick density or abundance. Middle of the road scenario (ssp2-4.5) where CO₂ levels remain similar to current levels, did not indicate a major change in the tick distributions. This tick distribution model could be used for targeting surveillance efforts and increasing the efficiency and accuracy of public health investigations and vector control strategies.

Introduction

Ticks are haematophagous parasites that have great economic and ecological significance due to their capacity to transmit a variety of pathogens including viruses, bacteria and parasites to animals and humans. Expansion of the range of ticks due to rapid climate change carries profound threats for public health and society (Illoldi-Rangel *et al.*, 2012; Rochlin and Toledo, 2020; Nuttall, 2022). Some of the most common tick-borne infections in the Middle East and North Africa (MENA) include Crimean–Congo haemorrhagic fever, anaplasmosis, theileriosis and babesiosis (Perveen *et al.*, 2021c).

The 1-humped camel (*Camelus dromedarius*) is a highly valued species of livestock in Saudi Arabia and UAE (Gharbi *et al.*, 2013). The current population of camels in UAE and Saudi Arabia is approximately 1 million (<https://worldpopulationreview.com/country-rankings/camel-population-by-country>). *Hyalomma dromedarii* ticks feed on the blood of camels and has been reported with high prevalence in the UAE (Perveen *et al.*, 2020, 2021b; Perveen, 2021) and Saudi Arabia (Alanazi *et al.*, 2018, 2020; Zakham *et al.*, 2021). Crimean–Congo haemorrhagic fever (CCHF) is a deadly viral disease and virus transmitted by *Hyalomma* ticks (Perveen and Khan, 2022). In a systematic review of Crimean–Congo haemorrhagic fever in the Arab world (1978–2021), a total of 65 confirmed human cases have been reported from the 2 countries. Lately, *H. dromedarii* ticks were found positive for CCHF virus (CCHFV) in both UAE (Camp *et al.*, 2020) and Saudi Arabia (Mohamed *et al.*, 2017). In the MENA Region, *H. dromedarii* appears to be the vector of *Theileria annulata* where camels are raised together with cattle (Jacquiet *et al.*, 1994), therefore, raising camels in mixed patterns could cause cross-infection in livestock (Tomassone *et al.*, 2012). Furthermore, *H. dromedarii* can transmit various disease-causing pathogens for example, Dhori virus (Hoogstraal *et al.*, 1981; Champour *et al.*, 2016), the tropical theileriosis, *T. annulata* and *T. camelensis* (Hoogstraal *et al.*, 1981; Hamed *et al.*, 2011) Sindbis virus, Chick Ross and Kadam viruses (Al-Khalifa *et al.*, 2007), *Coxiella burnetii* (Abdullah *et al.*, 2018) and spotted fever rickettsia (Hoogstraal *et al.*, 1981; Abdel-Shafy *et al.*, 2012; Demoncheaux *et al.*, 2012; Kernif *et al.*, 2012; Kleinerman *et al.*, 2013; Elzein *et al.*, 2020).

Due to anthropogenic factors and climate change, tick-borne infectious diseases are increasingly becoming a significant public health threat (Gray *et al.*, 2009). Ticks have unique physiological habits and spend their life cycle feeding on the host and in the habitat of the host. A range of environmental factors, such as substrate type, relative humidity and vegetation

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

associated with the host habitat can affect tick abundance and distribution patterns (Ma *et al.*, 2023). *Hyalomma dromedarii* may act as a 3-, 2- or 1-host species (Hoogstraal, 1956; Walker *et al.*, 2003) and engorged female burrows a few centimetres under the ground to lay eggs in suitable microhabitats to avoid desiccation of eggs and new larvae (Alahmed and Kheir, 2003). Environmental factors and host range may help in the assessment of risk factors determining the distribution of tick-borne pathogens. Increased temperatures may positively affect the survival and reproduction of ticks (Ma *et al.*, 2023). For example, rising temperatures in temperate and cold environments contributes to faster nymph maturation and shorter life cycles that increase tick abundance and also extend the period of ticks' host-seeking activity (Gray *et al.*, 2009), thus encourage range expansion through establishment in new geographical ranges. Furthermore, global climate change not only influences tick distribution and abundance, but also affects tick-borne pathogen transmission by impacting land use, vegetation cover and distribution, and the abundance of reservoir hosts (Gray *et al.*, 2009). Consequently, it is crucial to assess the current and future distribution of ticks for better management of tick-borne pathogens.

MaxEnt is a widely used technique in species distribution modelling because of its compatibility with presence-only (PO) data (Merow *et al.*, 2013; Bradie and Leung, 2017; Phillips *et al.*, 2022). Its algorithm is known for its robustness (Phillips *et al.*, 2006) and outperforms many other PO modelling methods (Phillips *et al.*, 2008; Wisz *et al.*, 2008; Elith *et al.*, 2011; Merow *et al.*, 2013). MaxEnt was developed specifically for low sample-size data (PO locations) for multiple species (Phillips *et al.*, 2006). This model used extensively to determine the distribution of numerous arthropods such as hard ticks and soft ticks that transmit important pathogens (Table 1).

In the MENA region, camel husbandry has increased in recent years with rapid economic development (Abahussain *et al.*, 2002).

In addition, climatic conditions of Saudi Arabia and UAE provide favourable conditions for tick species that are adapted to dry environments. Human activities and international trade increase the risk of tick expansion into new geographic zones. The present study was conducted to use MaxEnt modelling and the ArcGIS spatial technology platform to describe the current and predicted future distribution of the camel tick, *H. dromedarii* using occurrences of ticks in Saudi Arabia and UAE for the monitoring and surveillance of tick-borne pathogens associated with this species in the region.

Materials and methods

Collection and preparation of *H. dromedarii* occurrence data

Geo-referenced location points on ticks from Saudi Arabia and UAE from various resources including field collections and information from prior publications (Supplementary Table S1) were compiled. For Saudi Arabia, a literature review was conducted by search engines, Google Scholar, PubMed and Web of Science databases using the keywords 'Saudi Arabia', 'tick', 'ticks' and 'tick-borne pathogens' for articles published in the last 10 years. Only full-length research articles were used in this study. Review articles, letters to editors, short reports/communications, abstracts and conference proceedings were excluded. Literature containing geographical distribution information of camel tick species was filtered and extracted for their geographical location coordinates, Al-Khurma, Al-Kharj, Al-Hasa, Al-Qassim, Riyadh, Hail area, Amman Road, Madinah Road, Duba Road, Industrial area and Taif (Alanazi *et al.*, 2018, 2020, 2021; Alreshidi *et al.*, 2020; Zakham *et al.*, 2021; Al Thabiani *et al.*, 2022). After removing duplicates, only research articles were selected that provided coordinates for tick locations. Locations were chosen on the basis of presence of both hosts and tick vectors in the area. Occurrence data was rarefied using the spatially rarefy occurrence data tool in SDM toolbox in ArcGIS ver. 10.8.1 at a resolution of 15 km to avoid model over fitting and bias. It resulted in 32 occurrences which were later used in the spatial modelling.

Collection and preparation of key variables influencing tick distribution

For the current model, land use/land cover (LULC), elevation, slope and 19 bioclimatic variables (Table 2) were selected initially. The bioclimatic variables and the DEM were obtained from WorldClim database (version 2.1) at ~1 km² resolution (Fick and Hijmans, 2017) covering the period 1970–2000. Slope was calculated from elevation using the slope tool in ArcGIS. LULC was obtained from the European Space Agency climate change initiative for the year 2020 at 300 m resolution (Defourny *et al.*, 2023). The extent and resolution of all selected variables were harmonized to the same study area size and resolution of ~1 km² matching the bioclimatic variables using spatial analyst toolbox in ArcGIS.

Multicollinearity was assessed between the environmental variables using variance inflation factor analyses (VIF) in R (version 4.3.0). Highly correlated variables were eliminated considering a VIF <5 as a critical threshold (Akinwande *et al.*, 2015). Variables that did not demonstrate any significant contribution to the model were subsequently removed (Redon and Luque, 2010) and 7 variables were finally selected for modelling: LULC, elevation, slope, mean diurnal range (Bio2), isothermality (Bio3), precipitation seasonality (Bio15) and precipitation of coldest quarter (Bio19). The contribution of each variable is assessed by utilizing jackknife tests to visualize variable significance and calculating the percentage contributions of the variables.

Table 1. MaxEnt model used to predict the distribution of some hard and soft tick species

Tick species	References
<i>Hyalomma asiaticum</i> , <i>Dermacentor nuttalli</i> , <i>Ixodes persulcatus</i> , <i>Dermacentor silvarum</i>	Hu <i>et al.</i> (2022); Ma <i>et al.</i> (2023)
<i>Ixodes scapularis</i>	Johnson <i>et al.</i> (2016); Burrows <i>et al.</i> (2022); Zhang <i>et al.</i> (2022)
<i>Hyalomma marginatum</i>	Celina <i>et al.</i> (2023); Hekimoglu <i>et al.</i> (2023)
<i>Amblyomma testudinarium</i> , <i>Haemaphysalis flava</i> Neumann, <i>Haemaphysalis kitaokai</i> Hoogstraal, <i>Haemaphysalis longicornis</i> Neumann, <i>Haemaphysalis megapinosa</i> Saito, <i>Ixodes ovatus</i> Neumann	Doi <i>et al.</i> (2021)
<i>Argas persicus</i> , <i>Dermacentor marginatus</i> , <i>Haemaphysalis concinna</i> , <i>Haemaphysalis longicornis</i> , <i>Ixodes granulatus</i> , <i>Rhipicephalus microplus</i> , <i>Rhipicephalus sanguineus sensu</i> , <i>Rhipicephalus turanicus</i>	Yang <i>et al.</i> (2020)
<i>Ornithodoros hermsi</i>	Sage <i>et al.</i> (2017)
<i>Amblyomma americanum</i>	Raghavan <i>et al.</i> (2016)
<i>Ixodes ricinus</i> , <i>Rhipicephalus annulatus</i> , <i>Dermacentor marginatus</i> , <i>Haemaphysalis punctata</i>	Williams <i>et al.</i> (2015)
<i>Otobius megnini</i>	Estrada-Peña <i>et al.</i> (2010)

Table 2. Environmental layers for species distribution models

Code	Bioclimatic variables
Bio1	Annual mean temperature
Bio2	Mean diurnal range [mean of monthly (max temp – min temp)]
Bio3	Isothermality (difference in day-to-night temperature oscillations in relation to annual temperature oscillations).
Bio4	Temperature seasonality [standard deviation × 100]
Bio5	Max temperature of warmest month
Bio6	Min temperature of coldest month
Bio7	Temperature annual range
Bio8	Mean temperature of wettest quarter
Bio9	Mean temperature of driest quarter
Bio10	Mean temperature of warmest quarter
Bio11	Mean temperature of coldest quarter
Bio12	Annual precipitation
Bio13	Precipitation of wettest month
Bio14	Precipitation of driest month
Bio15	Precipitation seasonality [coefficient of variation]
Bio16	Precipitation of wettest quarter
Bio17	Precipitation of driest quarter
Bio18	Precipitation of warmest quarter
Bio19	Precipitation of coldest quarter

To assess the impact of different climate change scenarios on the spatial distribution of *H. dromedarii*, we excluded LULC. The future scenario used was ssp2-4.5, model CanESM5 which is among the most sensitive models in climate equilibrium (Swart *et al.*, 2019) and covered the periods 2021–2040 and 2041–2060. The ssp2-4.5 Scenario is considered to be ‘middle of the road’ in which CO₂ emissions remain approximately close to current levels before diminishing by the 2050s without achieving net-zero level emissions, and mean temperatures increase to 2.7°C by the end of the century. Socioeconomic factors continue to remain similar to historic trends and progress towards sustainability is slow, with development and income growing unevenly.

Bioclimatic variables were obtained from WorldClim database (version 2.1) at 30-arc-second (~1 km) resolution (Fick and Hijmans, 2017) except elevation and slope as it remained unchanged throughout the duration of the study. The extent and resolution of all selected variables were harmonized to the same study area size and resolution of ~1 km² matching the bioclimatic variables using spatial analyst toolbox in ArcGIS. All variables were processed to have the same spatial extent and resolution of ~1 km² using spatial analyst toolbox in ArcGIS.

Maxent modelling procedures and calibration

For the modelling analyses, MaxEnt 3.4.3 (Phillips *et al.*, 2022) was utilized. To address species-specific conditions and research objectives and to avoid relying solely on MaxEnt as a ‘blackbox’ tool (Hernandez *et al.*, 2006; Phillips *et al.*, 2008; Merow *et al.*, 2013), we used the spatial jackknifing tool within the SDM toolbox in ArcGIS (Brown *et al.*, 2017). This approach allowed for consideration of biological factors and provided a more comprehensive analysis. The tool tests the model using varying parameters and independently evaluates feature class parameters and the regularization multiplier (RM) to produce a model with the best performance. The RM enhances the model’s predictive accuracy and

achieve maximum entropy or a more uniform distribution which reduces model overfitting (Hernandez *et al.*, 2006; Phillips *et al.*, 2008). In addition, a bias file was generated using a gaussian kernel density approach for the sampling localities within the SDM toolbox. This bias file considers any sampling bias by providing MaxEnt with a background file that exhibits a similar level of bias observed in the presence localities. The bias file also enables the model to regulate the density and locations of background points, thereby avoiding the inclusion of less informative points that fall outside the known species range (Brown *et al.*, 2017).

The final 7 environmental variables along with 32 presence points were used to run 10 replicates by the cross-validation method. An RM of 5 with linear features and cross-validation method for all analyses. Iterations were raised to 5000 to prevent under- or overprediction of spatial relationships, considering the recommended convergence threshold of 10⁻⁵. The bioclimatic model was then projected onto 2040 and 2060 years under ssp2-4.5 scenario with no extrapolation. Recognizing the limitation in the predictive capabilities of the modelling algorithm during projection (Merow *et al.*, 2013), MaxEnt was prevented from extrapolating. In other words, MaxEnt did not make predictions beyond its training data during projecting. In addition, the provided bias file makes MaxEnt avoid sampling habitat outside the species’ known occurrence (Brown *et al.*, 2017).

Model evaluation/performance assessment

To assess the model’s performance, the receiver operating characteristic (ROC) curve was utilized, and the area under the curve (AUC) was calculated as a threshold-independent measure. AUC values range from 0 to 1, with higher values indicating better model performance (Merckx *et al.*, 2011). For a threshold-dependent measure, the true skill statistics (TSS) method was employed, using the threshold of maximum training sensitivity and specificity (West *et al.*, 2016). TSS accounts for both omission and commission errors and is less influenced by prevalence (Allouche *et al.*, 2006). TSS was interpreted based on ranges: <0.4 poor, 0.4–0.8 useful and >0.8 good-to-excellent performance (Zhang *et al.*, 2015). TSS calculations were conducted using Microsoft Excel.

To determine the relative importance of each variable in the model, contribution percentage and jackknife analysis were conducted using MaxEnt. Response curves were also measured for each predictor variable to illustrate the changes in habitat suitability corresponding to varying levels of the environmental variables.

Model exploration between current and future variables

To examine the differences between current and future variables, multivariate environmental similarity surfaces (MESS) and the most dissimilar variable (MoD) of the MESS map were computed using MaxEnt (Elith *et al.*, 2010). MESS shows similarity of a given point in the future to its reference current set of environmental layers. MoD shows the variable with the smallest similarity at each point (Elith *et al.*, 2010). Limiting factor analyses (LF) were also conducted in MaxEnt to examine the most influencing variable on model prediction at each point for current and future predictions (Elith *et al.*, 2010). All maps were processed and visualized in ArcGIS.

Results

Model evaluation and sensitivity analysis

The current distribution map of *H. dromedarii* is given in Fig. 1 (Supplementary Table S1). The distribution model showed consistent spatial distribution, with AUC-test at 0.772 and




Figure 1. *Hyalomma dromedarii* occurrence points in Saudi Arabia and UAE used for the distribution modelling.

AUC-train at 0.798, meaning the model had 77.2% performance (Table 1). The TSS result also indicated that the model is useful as the averaged value was TSS = 0.563 (Table 3) (Zhang *et al.*, 2015). LULC, Bio19 and elevation were the top contributors to the model, with 53.1, 21.8 and 20.6% (Table 3), respectively. The jack-knife test revealed that when LULC was used in isolation, the environmental variable showed the highest gain (Fig. 2) and this variable when omitted decreased the gain. The second most significant variable was the Bio19 when not used in the model, dropped the average gain followed closely by elevation (Fig. 2).

Predicted areas of *Hyalomma dromedarii* with potential suitability

Highly suitable areas (>0.6) existed mostly in the northern and eastern parts of the study area, with a considerable area near

Table 3. Evaluation test, sensitivity test and each variable contribution percentage in the model of *Hyalomma dromedarii*

Evaluation test	Result
AUCtest	0.772
AUCtrain	0.798
TSS	0.563
Variable	Contribution to the model (%)
LULC	53.1
Bio19	21.8
Elevation	20.6
Bio3	1.9
Bio2	1.8
Slope	0.5
Bio15	0.2

the Red Sea coast in the south (Fig. 3a, b). Very highly suitable areas (>0.8) corresponded mostly with urban cities across the study area (Fig. 3a, b). The UAE ranged from high (western region) to very high (mid- and northern region) in suitability. Highly suitable areas covered a good portion of Saudi Arabia and extended from north to east. Moderately suitable areas were mainly centred in the western and mid-regions of Saudi Arabia.

The potential future distribution is not predicted to change dramatically under the ssp2-4.5 scenario for the years 2040 and 2060 (Fig. 3c, d). A noticeable change is seen in Riyadh and Najran cities where suitability decreased sharply.

Model exploration

With the increasing value of slope, elevation and Bio2 (mean diurnal range) variables, the probability of occurrence declined sharply (Fig. 4a, b, g). High suitability (>0.6) occurred in areas where elevation ≤ 500 m and b2 $\leq 14.5^\circ\text{C}$. In contrast, potential suitability increased with increasing Bio19 (precipitation of coldest quarter) with highly suitable areas existing in areas receiving ≥ 38 mm (Fig. 4d). Suitability also slightly increased with increasing Bio15 (precipitation seasonality) and Bio3 (isothermality) (Fig. 4e, f). For LULC (land use/land cover) variable, suitability was moderate for croplands (10, 30), tree cover (60, 80), shrubland (120), sparse vegetation (150) and high for urban areas (190) (Fig. 4c).

The MESS analysis values ranged from -46 to 71 and from -46 to 73 for the years 2040 and 2060, respectively (Fig. 5a, b). Most areas were similar with varying degrees to current environmental conditions. Some areas had negative values demonstrating a degree of newness in environmental space. The highest novelty was in Riyadh city followed by Najran city where the novelty was mostly driven by Bio3 and Bio15 in both years, respectively (Fig. 5c, d).

Bio19 is the main limiting factor over the predicted existing range, followed by Bio2 in the limiting factor analyses (Fig. 6a, b).




Figure 2. Jackknife test (evaluation of each variable significance). LULC (land use/land cover), Bio2 (mean diurnal range), Bio3 (isothermality), Bio15 (precipitation seasonality), Bio19 (precipitation of coldest quarter).

Southern region in Saudi Arabia had elevation, slope and Bio15 slope as the limiting factors. Similarly, northern areas in the UAE had slope and Bio15 as the limiting factors. LULC was also limiting the distribution on the northern west coast of the UAE (Fig. 6a). For the potential future distribution (Fig. 6c, d), Bio3 limiting effect increased over Bio2 in northern Saudi areas as well in eastern UAE areas. Elevation limiting effect also increased in middle Saudi areas.

Discussion

The MaxEnt model has been widely used for predicting distributions of hundreds of animal species (Elith *et al.*, 2006). Our model helped us to better understand the environmental niche of *H. dromedarii* tick species in Saudi Arabia and UAE. The predicted maps developed from this model on current occurrences of *H. dromedarii* with a high probability based on suitable




Figure 3. Geographic distribution of *Hyalomma dromedarii* for (a) current, (b) current based on bioclimatic variables, (c) year 2040 under ssp2-4.5 scenario and (d) year 2060 under ssp2-4.5 scenario both based on bioclimatic variables.




Figure 4. Environmental variables response curves display probable occurrence of *Hyalomma dromedarii*. (a) Slope (m), (b) elevation (m), (c) LULC (land use/land cover), (d) Bio19 (precipitation of coldest quarter, mm), (e) Bio15 (precipitation seasonality, per cent), (f) Bio3 (isothermality, percent), (g) Bio2 (mean diurnal range, °C).




Figure 5. Model maps: (a, b) MESS (multivariate environmental similarity surfaces) analysis presenting the degree of resemblance between future and current set of environmental layers; (c, d) most dissimilar variable (MoD) analysis. (a, c) year 2040 and (b, d) year 2060, all under the bioclimatic distribution model.

environmental conditions. The modelled distribution of *H. dromedarii* indicated that highly suitable areas existed mostly in the northern, eastern and southern portions of the study range. In the UAE, the eastern to northern regions were classified as highly suitable for ticks. These regions include portions of eastern Abu Dhabi emirate, including Al Ain, bordering Oman, various cities within north-eastern portions of Abu Dhabi, such as Mafrak, Dubai and the northern Emirates, where the coastal zones influence moisture and temperature profiles. This increases the suitability of ticks in these regions. In addition, farms abound with high density of camels and other livestock that help to sustain tick populations (Perveen *et al.*, 2021c). In Saudi Arabia, areas that were highly suitable were within the north-central region, the eastern edge south of the UAE and the southwestern shoreline bordering the Red Sea, whereas western and mid-regions were of moderate suitability for ticks. The north central regions with high suitability were east of the Al Nafud desert region. This area is adjacent to areas that are around 500 m in elevation, and receives more precipitation, making them suitable for ticks. Similarly, the southwestern regions are immediately adjacent to the Asir Mountain range that has elevations of over 2000 m, with significant moisture draining out of them into the Red Sea, increasing moisture content and influencing temperature along the southwestern coastline. Saudi Arabia and UAE camel and livestock farming represent an essential habitat for *H. dromedarii* and

other ticks, which are likely enhanced in these regions due to better environmental conditions compared to the remaining parts of the study area.

Bioclimatic factors combined with LULC have a cumulative influence on determining the suitability of tick habitats. The survival of ticks during its off-host phase is heavily reliant on variables like temperature and humidity (Apanaskevich *et al.*, 2013; Pascoe *et al.*, 2019). The elevation of an area affects its microclimate, the presence of hosts and the vegetation. Additionally, slope serves as an indicator of subsurface water flow velocity, runoff rate and soil moisture content (Pascoe *et al.*, 2019). The presence and characteristics of vegetation also impact the suitability of tick habitats. Vegetation plays a crucial role in the life cycle of tick hosts, such as camels, where the availability of cropland, herbaceous cover and water are decisive factors for host presence in a given area (Apanaskevich *et al.*, 2013; Pascoe *et al.*, 2019). Based on the current climate, several variables influenced the *H. dromedarii* tick distribution including land use/land cover LULC (53.1%), Bio19 (21.8%), elevation (20.6%), Bio3 (1.9%), Bio2 (1.8%), slope (0.5%) and Bio15 (0.2%). The precipitation of coldest quarter is significant to the tick survival in the winter season. For land use/land cover variable, suitability was modest for sparse vegetation (150) and high for urban areas (190), maybe due to anthropogenic factors such as land-use change, agriculture practices, forest fragmentation or urbanization.




Figure 6. Limiting factor (LF) analyses highlighting the most influencing variable on model prediction for (a) current, (b) current, based on bioclimatic variables, (c) year 2040 under ssp2-4.5 scenario and (d) year 2060 under ssp2-4.5 scenario both based on bioclimatic variables.

The jackknife results showed that the LULC when used in isolation was with the highest gain. The high suitability/high tick density sites are the areas categorized by a high proportion of land cover (Burrows *et al.*, 2022). Moreover, the land-use and land cover patterns may provide microclimatic conditions through vegetation covers that serve as ticks habitats (Doi *et al.*, 2021; Khwarahm, 2023). Our findings are almost similar to the one conducted in Iraq on distribution of the *Hyalomma* spp. where distribution influenced by LULC (50.8%), followed by elevation (30.4%) (Khwarahm, 2023). A similar study was conducted on distribution of *Ornithodoros hermsi* where annual temperature range (Bio7) contribution was highest (18.9%) in the model, followed by elevation (18.1%), and precipitation of the warmest quarter (Bio18) in Sage *et al.* (2017). However, in another study minimum temperature of coldest month (Bio6) and precipitation of driest quarter (Bio17) strongly influenced the model (Porretta *et al.*, 2013). In Mongolia, annual precipitation (Bio12) and elevation influenced the *Hyalomma asiaticum* distribution (Ma *et al.*, 2023). The differences in the impact of environmental variables in other studies are most likely due to species-specific niche requirements, an area that requires extensive study in the Middle East region.

Response curves of the environmental variables showed that *H. dromedarii* potential suitability increased with increasing precipitation of coldest quarter (Bio19), in areas receiving ≥ 38 mm while declining sharply with the increasing value of slope and elevation.

The potential future distribution is not predicted to change dramatically under the ssp2-4.5 scenario for 2040 and 2060 years; however, a visible change has been seen in Riyadh and Najran cities where suitability decreased sharply mostly driven by isothermality and precipitation seasonality in both years, respectively. Therefore, climate change can impact the tick distribution in forthcoming years. This species disperses naturally with the help of infested animals, but it can also inhabit new ranges through travel and transportation of animals across the borders. Due to its high prevalence on camels in both countries (Alanazi *et al.*, 2020; Perveen *et al.*, 2020; Perveen, 2021), its distribution is a continued threat in the region. In addition, it serves as a reservoir of many tick-borne pathogens. Therefore, the current study will assist researchers and health care managers to devise the strategies to limit the distribution of ticks to better avoid tick-borne zoonotic diseases in the future. Various zoonotic pathogens have been reported previously in the region including MERS-CoV (Zaki, 2012; de Groot *et al.*, 2013; Perveen *et al.*, 2021a) and CCHFV (Camp *et al.*, 2020; Perveen and Khan, 2022) posing a serious threat to camel farming and human and animal health. *Hyalomma* spp. are primary vectors of CCHFV (Perveen and Khan, 2022). Recently, *H. dromedarii* tested positive for CCHFV in both countries (Mohamed *et al.*, 2017; Camp *et al.*, 2020). Previously, a virus related to the tick-borne encephalitis complex has also been detected in Saudi Arabia (Zaki, 1997).

The true distribution of ticks and other invertebrates are difficult to determine as widespread sampling of ticks, hosts and their environments are not available in the MENA region (Perveen et al., 2021c). In the absence of true distribution, model validation provides the best way of assessing if the predicted distribution falls within reasonable bounds of statistical uncertainty (Convertino et al., 2014; Liu et al., 2016; Chen et al., 2019). We used AUC and TSS values that indicated that the model was able to predict the distribution based on the quality of the data available. The Kappa statistic is typically used for validating the accuracy of the distribution models, including MaxEnt models (Allouche et al., 2006). However, the Kappa statistic has a linear relationship with prevalence (the proportion of sites in which the species is present), making it a potential statistical artefact (Allouche et al., 2006; Liu et al., 2016). In comparison, the TSS is independent of prevalence making it better in accurately predicting distribution (Allouche et al., 2006; Liu et al., 2016). Values of TSS that we obtained were in the useful category. This may suggest that further data from more sampling sites could improve the model.

In summary, the model showed that LULC, precipitation of coldest quarter, and elevation were most influential for predicting the areas as highly suitable for ticks. Other environmental variables which contributed in the model were isothermality, mean diurnal range [mean of monthly (max temp – min temp)], slope, precipitation seasonality. The model presented here provides valuable information on camel tick species distribution in Saudi Arabia and UAE. The predicted distribution of *H. dromedarii* may allow researchers and health officials to conduct risk assessments targeting specific pathogens and potentially reduce the chance of outbreaks through surveillance and mitigation efforts.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182024001161>.

Data availability statement. Locations used in this study are available in supplementary material.

Acknowledgements. The authors thank Daniil Iliashevich, Department of Veterinary Medicine, College of Agriculture and Veterinary Medicine, United Arab Emirates University, Al-Ain, P.O. Box 15551, UAE, for his assistance in collecting camel tick locations in the UAE.

Author contributions. N. P., S. B. M. and A. L. W. conceived and designed the study. N. P. conducted data gathering. A. J., N. P., S. B. M. and O. A. S. performed analysis. N. P., S. B. M. and A. J. wrote original draft. N. P., S. B. M., O. A. S., and A. L. W. reviewed and edited the manuscript. A. L. W. supervised, administered the project and acquired funding. Nighat Perveen and Sabir B. Muzaffar. These authors contributed equally to this article.

Financial support. This study was funded by the UAE University through UPAR Grant # G00003709 to Arve Lee Willingham.

Competing interests. The authors declare there are no conflicts of interest.

Ethical standards. Tick collection in the UAE was performed in accordance with the experimental protocol approved by the Animal Research Ethics Committee of the United Arab Emirates University (ethical approval #ERA_2022-1647).

References

- Abahussain AA, Abdu AS, Al-Zubari WK, El-Deen NA and Abdul-Raheem M (2002) Desertification in the Arab Region: analysis of current status and trends. *Journal of Arid Environments* **51**, 521–545.
- Abdel-Shafy S, Allam NAT, Mediannikov O, Parola P and Raoult D (2012) Molecular detection of spotted fever group Rickettsiae associated with ixodid ticks in Egypt. *Vector-Borne and Zoonotic Diseases* **12**, 346–359.
- Abdullah HHAM, El-shanawany EE, Abdel-shafy S, Abou-zeina HAA and Abdel-rahman EH (2018) Molecular and immunological characterization of *Hyalomma dromedarii* and *Hyalomma excavatum* (Acari: Ixodidae) vectors of Q fever in camels. *Veterinary World* **11**, 1109–1119.
- Akinwande MO, Dikko HG and Samson A (2015) Variance inflation factor: as a condition for the inclusion of suppressor variable(s) in regression analysis. *Open Journal of Statistics* **5**, 754–767.
- Alahmed AM and Kheir SM (2003) Life cycle and survival of *Hyalomma dromedarii* (Acari: Ixodidae) under laboratory conditions. *Agricultural and Marine Sciences* **8**, 11–14.
- Alanazi AD, Abdullah S, Wall R and Alharbi SA (2018) Tick-borne pathogens in ticks and blood samples collected from camels in Riyadh province, Saudi Arabia. *International Journal of Zoological Research* **14**, 30–36.
- Alanazi AD, Nguyen VL, Alyousif MS, Manoj RRS, Alouffi AS, Donato R, Sazmand A, Mendoza-Roldan JA, Dantas-Torres F and Otranto D (2020) Ticks and associated pathogens in camels (*Camelus dromedarius*) from Riyadh province, Saudi Arabia. *Parasites and Vectors* **13**, 1–9.
- Alanazi AD, Alouffi AS, Alshahrani MY, Alyousif MS, Abdullah HHAM, Allam AM, Elsayy BSM, Abdel-Shafy S, Alsulami MN, Khan A and Iqbal F (2021) A report on tick burden and molecular detection of tick-borne pathogens in cattle blood samples collected from four regions in Saudi Arabia. *Ticks and Tick-Borne Diseases* **12**, 101652.
- Al-Khalifa MS, Diab FM and Khalil GM (2007) Man-threatening viruses isolated from ticks in Saudi Arabia. *Saudi Medical Journal* **28**, 1864–1867.
- Allouche O, Tsoar A and Kadmon R (2006) Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology* **43**, 1223–1232.
- Alreshidi MM, Veettil VN, Noumi E, Del Campo R and Snoussi M (2020) Description of microbial diversity associated with ticks *Hyalomma dromedarii* (Acari: Ixodidae) isolated from camels in Hail region (Saudi Arabia) using massive sequencing of 16S rDNA. *Bioinformatics* **16**, 602–610.
- Al Thabiani A, Panneerselvam C, Alshehri MA, Asiry KA, Alsaif M and Alhowity Y (2022) Efficacy of synthetic pyrethroids on camel ticks *Hyalomma dromedarii* 'Acari: Ixodidae' in Saudi Arabia. *Entomology and Applied Science Letters* **8**, 27–32.
- Apanaskevich DA, Oliver JH, Sonenshine DE and Roe RM (2013) Life cycles and natural history of ticks. In Sonenshine DE and Roe RM (eds), *Biology of Ticks*, Vol. 1. New York: Oxford University Press, pp. 59–73.
- Bradie J and Leung B (2017) A quantitative synthesis of the importance of variables used in MaxEnt species distribution models. *Journal of Biogeography* **44**, 1344–1361.
- Brown JL, Bennett JR and French CM (2017) SDMtoolbox 2.0: the next generation Python-based GIS toolkit for landscape genetic, biogeographic and species distribution model analyses. *PeerJ* **2017**, 1–12.
- Burrows H, Slatculescu AM, Feng CX, Clow KM, Guillot C, Jardine CM, Leighton PA, Krause PJ and Kulkarni MA (2022) The utility of a maximum entropy species distribution model for *Ixodes scapularis* in predicting the public health risk of Lyme disease in Ontario, Canada. *Ticks and Tick-Borne Diseases* **13**, 1–9.
- Camp JV, Kannan DO, Osman BM, Shah MS, Howarth B, Khafaga T, Weidinger P, Karuvantevida N, Kolodziejek J, Mazrooei H, Wolf N, Loney T and Nowotny N (2020) Crimean-Congo hemorrhagic fever virus endemicity in United Arab Emirates, 2019. *Emerging Infectious Diseases* **26**, 1019–1021.
- Celina SS, Černý J and Samy AM (2023) Mapping the potential distribution of the principal vector of Crimean-Congo haemorrhagic fever virus *Hyalomma marginatum* in the Old World. *PLoS Neglected Tropical Diseases* **17**, e0010855.
- Champour M, Chinikar S and Mohammadi G (2016) Molecular epidemiology of Crimean-Congo hemorrhagic fever virus detected from ticks of one humped camels (*Camelus dromedarius*) population in northeastern Iran. *Journal of Parasitic Diseases* **40**, 110–115.
- Chen X, Dimitrov NB and Meyers LA (2019) Uncertainty analysis of species distribution models. *PLoS ONE* **14**, 1–11.
- Convertino M, Muñoz-Carpena R, Chu-Agor ML, Kiker GA and Linkov I (2014) Untangling drivers of species distributions: global sensitivity and uncertainty analyses of MaxEnt. *Environmental Modelling and Software* **51**, 296–309.
- Defourny P, Lamarche C, Brockmann C, Boettcher M, Bontemps S, De Maet T, Duveiller GL, Harper K, Hartley A, Kirches G, Moreau I, Peylin P, Ottlé C, Radoux J, Van Bogaert E, Ramoino F, Albergel C and Arino O (2023) *Climate Change Initiative Data: Observed Annual Global Land-Use Change from 1992 to 2020 Three Times More Dynamic than Reported by Inventory-Based Statistics*. Oxfordshire, UK: European Space Agency Climate Change Initiative (ESA CCI).

- de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, Fouchier RAM, Galiano M, Gorbalenya AE, Memish ZA, Perlman S, Poon LLM, Snijder EJ, Stephens GM, Woo PCY, Zaki AM, Zambon M and Ziebuhr J (2013) Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the coronavirus study group. *Journal of Virology* **87**, 7790–7792.
- Demoncheaux JP, Socolovschi C, Davoust B, Haddad S, Raoult D and Parola P (2012) First detection of *Rickettsia aeschlimannii* in *Hyalomma dromedarii* ticks from Tunisia. *Ticks and Tick-Borne Diseases* **3**, 398–402.
- Doi K, Kato T, Tabata I and Hayama S (2021) Mapping the potential distribution of ticks in the western Kanto region, Japan: predictions based on land-use, climate, and wildlife. *Insects* **12**, 1095.
- Elith J, Graham CH, Anderson RP, Dudík M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JMcCM, Townsend Peterson A, ... Zimmermann NE (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography* **29**, 129–151.
- Elith J, Kearney M and Phillips SJ (2010) The art of modelling range-shifting species. *Methods in Ecology and Evolution* **1**, 330–342.
- Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE and Yates CJ (2011) A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* **17**, 43–57.
- Elzein FE, Aloteibi M, Alanazi W, Alsaed M and Almaghaslah M (2020) A rickettsia infection from Saudi Arabia. *International Journal of Infectious Diseases* **90**, 167–169.
- Estrada-Peña A, Nava S, Horak IG and Guglielmo AA (2010) Using ground-derived data to assess the environmental niche of the spinose ear tick, *Otobius megnini*. *Entomologia Experimentalis et Applicata* **137**, 132–142.
- Fick SE and Hijmans RJ (2017) Worldclim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* **37**, 4302–4315.
- Gharbi M, Moussi N, Jedidi M, Mhadhbi M and Sassi L (2013) Population dynamics of ticks infesting the one-humped camel (*Camelus dromedarius*) in central Tunisia. *Ticks and Tick-Borne Diseases* **4**, 488–491.
- Gray JS, Dautel H, Estrada-Pena A, Kahl O and Lindgren E (2009) Effects of climate change on ticks and tick-borne diseases in Europe. *Interdisciplinary Perspectives on Infectious Diseases* **2009**, e593232.
- Hamed MI, Zaitoun AMA, El-Allawy TAA and Mourad MI (2011) Investigation of *Theileria camelensis* in camels infested by *Hyalomma dromedarii* ticks in Upper Egypt. *Journal of Advanced Veterinary Research* **1**, 4–7.
- Hekimoglu O, Elverici C and Kuyucu AC (2023) Predicting climate-driven distribution shifts in *Hyalomma marginatum* (Ixodidae). *Parasitology* **150**, 883–893.
- Hernandez PA, Graham CH, Master LL and Albert DL (2006) The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography* **29**, 773–785.
- Hoogstraal H (1956) African Ixodidae: ticks of Sudan (with special reference to Equatoria province and with preliminary reviews of the genera *Boophilus*, *Margaropus*, and *Hyalomma*). In *Research Report NM 005 050.29.07*. Cairo, Egypt: Department of Medical Zoology U. S. Naval Medical Research Unit No. 3, pp. 598.
- Hoogstraal H, Wassef HY and Buttiker W (1981) Ticks of Saudi Arabia. *Fauna of Arabia* **3**, 25–110.
- Hu E, Hu Z, Mi X, Li C, He W, Gan L, Li Y, Zhang W, Meng Y and Gailike B (2022) Distribution prediction of *Hyalomma asiaticum* (Acari: Ixodidae) in a localized region in northwestern China. *Journal of Parasitology* **108**, 330–336.
- Illoldi-Rangel P, Rivaldi C-L, Sissel B, Trout Fryxell R, Gordillo-Pérez G, Rodríguez-Moreno A, Williamson P, Montiel-Parra G, Sánchez-Cordero V and Sarkar S (2012) Species distribution models and ecological suitability analysis for potential tick vectors of lyme disease in Mexico. *Journal of Tropical Medicine* **2012**, 1–10.
- Jacquet P, Colas F, Cheikh D, Thiam E and Ly BA (1994) Epidémiologie descriptive de la theilériose bovine à *Theileria annulata* en Mauritanie, l'Afrique de l'Ouest-saharienne. *Revue D'élevage Et de Médecine Veterinaire Des Pays Tropicaux* **47**, 147–155.
- Johnson TL, Bjork JKH, Neitzel DF, Dorr FM, Schiffman EK and Eisen RJ (2016) Habitat suitability model for the distribution of *Ixodes scapularis* (acari: Ixodidae) in Minnesota. *Journal of Medical Entomology* **53**, 598–606.
- Kernif T, Djerbouh A, Mediannikov O, Ayach B, Rolain JM, Raoult D, Parola P and Bitam I (2012) *Rickettsia africae* in *Hyalomma dromedarii* ticks from sub-Saharan Algeria. *Ticks and Tick-Borne Diseases* **3**, 377–379.
- Khwarahm NR (2023) Predicting the spatial distribution of *Hyalomma* spp., vector ticks of Crimean–Congo haemorrhagic fever in Iraq. *Sustainability* **15**, 13669.
- Kleinerman G, Baneth G, Mumcuoglu KY, van Straten M, Berlin D, Apanaskevich DA, Abdeen Z, Nasereddin A and Harrus S (2013) Molecular detection of *Rickettsia africae*, *Rickettsia aeschlimannii*, and *Rickettsia sibirica mongolitimonae* in camels and *Hyalomma* spp. ticks from Israel Gabriela. *Vector-Borne and Zoonotic Diseases* **13**, 851–856.
- Liu C, Newell G and White M (2016) On the selection of thresholds for predicting species occurrence with presence-only data. *Ecology and Evolution* **6**, 337–348.
- Ma R, Li C, Tian H, Zhang Y, Feng X, Li J and Hu W (2023) The current distribution of tick species in Inner Mongolia and inferring potential suitability areas for dominant tick species based on the MaxEnt model. *Parasites and Vectors* **16**, 1–15.
- Merckx B, Steyaert M, Vanreusel A, Vincx M and Vanaverbeke J (2011) Null models reveal preferential sampling, spatial autocorrelation and overfitting in habitat suitability modelling. *Ecological Modelling* **222**, 588–597.
- Merow C, Smith MJ and Silander JA (2013) A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography* **36**, 1058–1069.
- Mohamed RAEH, Mohamed N, Aleanizy FS, Alqahtani FY, Al Khalaf A and Al-Keridis LA (2017) Investigation of hemorrhagic fever viruses inside wild populations of ticks: one of the pioneer studies in Saudi Arabia. *Asian Pacific Journal of Tropical Disease* **7**, 299–303.
- Nuttall PA (2022) Climate change impacts on ticks and tick-borne infections. *Biologia* **77**, 1503–1512.
- Pascoe EL, Marcantonio M, Caminade C and Foley JE (2019) Modeling potential habitat for *Amblyomma* tick species in California. *Insects* **10**, 1–19.
- Perveen N (2021) *Livestock Ticks in the UAE: Prevalence, Distribution, Population Dynamics, and Associated Microorganisms*. PhD thesis, Department of Biology, College of Science, United Arab Emirates University, Al Ain, UAE, p. 206.
- Perveen N and Khan G (2022) Crimean – Congo hemorrhagic fever in the Arab world: a systematic review. *Frontiers in Veterinary Science* **9**, 1–16.
- Perveen N, Muzaffar SB and Al-Deeb MA (2020) Population dynamics of *Hyalomma dromedarii* on camels in the United Arab Emirates. *Insects* **11**, 1–9.
- Perveen N, Muzaffar SB and Al-Deeb MA (2021a) Exploring human-animal host interactions and emergence of COVID-19: evolutionary and ecological dynamics. *Saudi Journal of Biological Sciences* **28**, 1417–1425.
- Perveen N, Muzaffar SB and Al-Deeb MA (2021b) Four tick-borne microorganisms and their prevalence in *Hyalomma* ticks collected from livestock in United Arab Emirates. *Pathogens* **10**, 1005.
- Perveen N, Muzaffar SB and Al-Deeb MA (2021c) Ticks and tick-borne diseases of livestock in the Middle East and north Africa: a review. *Insects* **12**, 1–34.
- Phillips SB, Aneja VP, Kang D and Arya SP (2006) Modelling and analysis of the atmospheric nitrogen deposition in North Carolina. *International Journal of Global Environmental Issues* **6**, 231–252.
- Phillips SJ, Dudík M and Phillips SJ (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* **31**, 161–175.
- Phillips SJ, Dudík M and Schapire RE (2022) Maxent software for modeling species niches and distributions Version 3.4.3.
- Porretta D, Mastrantonio V, Amendolia S, Gaiarsa S, Epis S, Genchi C, Bandi C, Otranto D and Urbanelli S (2013) Effects of global changes on the climatic niche of the tick *Ixodes ricinus* inferred by species distribution modelling. *Parasites & Vectors* **6**, 271.
- Raghavan RK, Goodin DG, Hanzlicek GA, Zolnerowich G, Dryden MW, Anderson GA and Ganta RR (2016) Maximum entropy-based ecological niche model and bio-climatic determinants of lone star tick (*Amblyomma americanum*) niche. *Vector-Borne and Zoonotic Diseases* **16**, 205–211.
- Redon M and Luque S (2010) Presence-only modelling for indicator species distribution: biodiversity monitoring in the French Alps. In *6th Spatial Analysis and Geomatics International Conference (SAGEO 2010)*, Vol. 1. Université de Toulouse, pp. 42–55. <https://hal.science/hal-00558859>
- Rochlin I and Toledo A (2020) Emerging tick-borne pathogens of public health importance: a mini-review. *Journal of Medical Microbiology* **69**, 781–791.

- Sage KM, Johnson TL, Teglas MB, Nieto NC and Schwan TG (2017) Ecological niche modeling and distribution of *Ornithodoros hermsi* associated with tick-borne relapsing fever in western North America. *PLoS Neglected Tropical Diseases* **11**, 1–18.
- Swart N, Cole JNS, Kharin VV, Lazare M, Scinocca JF, Gillett NP, Anstey J, Arora V, Christian JR, Hanna S, Jiao Y, Lee WG, Majaess F, Saenko OA, Seiler C, Seinen C, Shao A, Sigmond M, Solheim L, von Salzen K, Yang D and Winter B (2019) The Canadian earth system model version 5 (CanESM5.0.3). *Geoscientific Model Development* **12**, 4823–4873.
- Tomassone L, Grego E, Callà G, Rodighiero P, Pressi G, Gebre S, Zeleke B and De Meneghi D (2012) Ticks and tick-borne pathogens in livestock from nomadic herds in the Somali region, Ethiopia. *Experimental and Applied Acarology* **56**, 391–401.
- Walker AR, Bouattour A, Camicas JL, Estrada-Peña A, Horak IG, Latif AA, Pegram RG and Preston PM (2003) *Ticks of Domestic Animals in Africa: A Guide to Identification of Species*. Edinburgh, UK: Bioscience Reports. Available at <http://www.alanwalker.com/assets/PDF/tickguide-africa.pdf>.
- West AM, Kumar S, Brown CS, Stohlgren TJ and Bromberg J (2016) Field validation of an invasive species Maxent model. *Ecological Informatics* **36**, 126–134.
- Williams HW, Cross DE, Crump HL, Drost CJ and Thomas CJ (2015) Climate suitability for European ticks: assessing species distribution models against null models and projection under AR5 climate. *Parasites and Vectors* **8**, 1–15.
- Wisn MS, Hijmans RJ, Li J, Peterson AT, Graham CH, Guisan A, and NCEAS Predicting Species Distributions Working Group (2008) Effects of sample size on the performance of species distribution models. *Diversity and Distributions* **14**, 763–773.
- Yang X, Gao Z, Zhou T, Zhang J, Wang L, Xiao L, Wu H and Li S (2020) Mapping the potential distribution of major tick species in China. *International Journal of Environmental Research and Public Health* **17**, 1–15.
- Zakham F, Albalawi AE, Alanazi AD, Nguyen PT, Alouffi AS, Alaoui A, Sironen T, Smura T and Vapalahti O (2021) Viral RNA metagenomics of *Hyalomma* ticks collected from dromedary camels in Makkah province, Saudi Arabia. *Viruses* **13**, 1–13.
- Zaki AM (1997) Isolation of a flavivirus related to the tick-borne encephalitis complex from human cases in Saudi Arabia. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **91**, 179–181.
- Zaki AM (2012) Novel coronavirus – Saudi Arabia: human isolate. *International Society for Infectious Diseases*. <http://www.promedmail.org/direct.php?id=20120920.1302733>
- Zhang L, Liu S, Sun P, Wang T, Wang G, Zhang X and Wang L (2015) Consensus forecasting of species distributions: the effects of niche model performance and niche properties. *PLoS ONE* **10**, 1–18.
- Zhang L, Ma D, Li C, Zhou R, Wang J and Liu Q (2022) Projecting the potential distribution areas of *Ixodes scapularis* (Acari: Ixodidae) driven by climate change. *Biology* **11**, 1–13.