

A metered intake of milk following exercise and thermal dehydration restores whole-body net fluid balance better than a carbohydrate–electrolyte solution or water in healthy young men

Suzanne Seery¹ and Philip Jakeman^{1,2*}

¹Department of Physical Education and Sport Sciences, University of Limerick, Limerick V94 YDE9, Republic of Ireland

²Human Science Research Unit, Centre for Interventions in Inflammation, Infection and Immunity, University of Limerick, Limerick V94 YDE9, Republic of Ireland

(Submitted 18 November 2015 – Final revision received 23 June 2016 – Accepted 29 June 2016 – First published online 1 August 2016)

Abstract

Appropriate rehydration and nutrient intake in recovery is a key component of exercise performance. This study investigated whether the recovery of body net fluid balance (NFB) following exercise and thermal dehydration to –2% of body mass (BM) was enhanced by a metered rate of ingestion of milk (M) compared with a carbohydrate–electrolyte solution (CE) or water (W). In randomised order, seven active men (aged 26.2 (SD 6.1) years) undertook exercise and thermal dehydration to –2% of BM on three occasions. A metered replacement volume of M, CE or W equivalent to 150% of the BM loss was then consumed within 2–3 h. NFB was subsequently measured for 5 h from commencement of rehydration. A higher overall NFB in M than CE ($P=0.001$) and W ($P=0.006$) was observed, with no difference between CE and W ($P=0.69$). After 5 h, NFB in M remained positive (+117 (SD 122) ml) compared with basal, and it was greater than W (–539 (SD 390) ml, $P=0.011$) but not CE (–381 (SD 460) ml, $P=0.077$, $d=1.6$). Plasma osmolality (P_{osm}) and K remained elevated above basal in M compared with CE and W. The change in P_{osm} was associated with circulating pre-provasopressin ($r_s 0.348$, $P<0.001$), a biomarker of arginine vasopressin, but could not account fully for the augmented NFB in M compared with CE and W. These data suggest that a metered approach to fluid ingestion acts in synergy with the nutrient composition of M in the restoration of NFB following exercise and thermal dehydration.

Key words: Rehydration: Metered fluid intake: Milk: Carbohydrate electrolyte: Net water balance

Normal physiological and behavioural perturbation of net body water (fluid) balance changes daily total body water (TBW) by <1%⁽¹⁾. The demands of physical activity perturb whole-body fluid balance and results in a net water deficit (dehydration) that may affect exercise tolerance and/or performance^(2–4). Physical activity accelerates the rate of substrate oxidation and the depletion of body energy stores. Particularly during intense exercise, the preferential use of carbohydrate as a substrate leads to accelerated use of muscle glycogen and an increase in branched chain amino acid oxidation by muscle that is inversely related to muscle glycogen content⁽⁵⁾. After exercise, amino acids, particularly the essential amino acids (EAA), are required in support of tissue repair and remodelling. A recovery beverage should, therefore, fulfil both fluid and nutrient requirements. Supported by evidence that milk can act as an effective rehydration fluid^(6,7), it is reasoned that the nutrient composition of milk offers a superior recovery fluid than water or carbohydrate–electrolyte solutions (CE).

A common, laboratory-based, experimental paradigm to investigate the optimal strategy for fluid replacement is to

undertake prolonged (approximately 90 min) exercise and thermal dehydration to a fixed decrease in body mass (–2% BM), a surrogate of change in TBW^(8,9). The outcome of these studies recommends a fluid replacement regimen that requires the ingestion of a fluid volume equivalent to 150% of the BM loss during exercise to be consumed within 1 h after exercise⁽¹⁰⁾. The volume of fluid equivalent to 150% of an approximately 2% loss in BM for an 80 kg male approximates to 2.5 litres, which, when ingested in bolus mode over a short time (i.e. 1 h), results in enhanced diuresis and failure to restore net fluid balance (NFB) within 4–6 h^(8,9,11). The rate and magnitude of fluid retention following exercise and thermal dehydration is enhanced by the addition of electrolytes^(11–13), carbohydrate^(14–17) and protein^(18–21) affecting singly, or in combination, the inter-related processes of gastric emptying, intestinal absorption and retention. Addition of electrolytes, carbohydrate and protein may also fulfil the requirement to replenish body reserves and recovery of muscle glycogen.

Recent evidence suggests that a ‘metered’ approach to fluid replacement post exercise is a more effective strategy⁽²²⁾.

Abbreviations: AVP, arginine vasopressin; BM, body mass; CE, carbohydrate–electrolyte solution; CTproAVP, pre-provasopressin; M, milk; NFB, net fluid balance; P_{osm} , plasma osmolality; W, water.

* **Corresponding author:** Professor P. Jakeman, fax +353 6120 2841, email phil.jakeman@ul.ie

A metered fluid intake has greater ecological validity for the athlete, but whether the presence of electrolytes or macronutrients in the rehydration fluid would act in synergy with a metered fluid intake to improve body NFB following exercise is not known. In this study, exercise and thermal dehydration to -2% BM was used to examine net body fluid balance over a 5-h period following metered replacement of three rehydration fluids of constant volume, equivalent to 150% of the exercise-induced BM loss. Plain water (W) was used as the control fluid against which a proprietary CE and low-fat milk (M) was compared. CE is purported to be optimised for post-exercise restoration of fluid and carbohydrate and is a popular form of replenishment fluid following exercise⁽²³⁾. M was chosen for the enhanced electrolyte (K) and macro-nutrient (carbohydrate, protein and fat) composition and healthier nutrient profile compared with proprietary CE products. The primary aim was to investigate whether NFB following exercise and thermal dehydration to -2% BM was enhanced by a metered rate of ingestion of M or CE compared with W. A secondary aim was to investigate whether the composition of the rehydration fluid was sufficient to affect a change in circulating pre-provasopressin (CTproAVP) – a novel biomarker of the potent antidiuretic arginine vasopressin (AVP) and signature of an augmented osmoregulatory response.

Methods

Subjects

All procedures were carried out in accordance with the ethical standards outlined in the most recent version of the Declaration of Helsinki, and procedures involving human subjects were approved by the Faculty of Education & Health Sciences Research Ethics Committee (EHSREC10/45), University of Limerick. Subjects were provided with full details of the study, and queries related to the study were addressed before a written statement of informed consent was obtained. Seven healthy men (age 26.2 (SD 6.1) years, height 1.79 (SD 0.08) m, BM 86.4 (SD 11.5) kg and body fat 21.4 (SD 4.5)%) volunteered to participate in the investigation. At the time of the study, all were engaged in regular physical activity (>3 h/week) but were not accustomed to exercise in a warm environment. Because of the nature of the investigation, those with known lactose intolerance were excluded.

Experimental protocol

All subjects attended two habituation and three experimental trials. In the 24 h preceding each trial, subjects were instructed to abstain from alcohol and strenuous physical activity. Subjects also recorded a weighed food and fluid intake and physical activity diary and were asked to replicate these patterns of dietary intake and physical activity before each subsequent trial. In the 24 h preceding the first trial, subjects provided a record of weighed food and fluid intake and a diary of physical activity. Subjects were instructed to repeat these behaviours before subsequent trials. All trials commenced in the morning, normally 07.00 hours, and followed a standardised protocol.

On the evening preceding a trial day, subjects were instructed to fast from midnight and to consume 500 ml of water 1 h before arrival at the laboratory. On arrival, subjects completed a subjective feelings questionnaire and provided a urine and venous blood sample. Height, to the nearest 0.1 cm (Seca 213 Stadiometer; Seca), BM, to the nearest 0.1 kg, TBW (bioelectric impedance analysis; MC-180MA; Tanita UK Ltd) and body composition (dual energy X-ray absorptiometry; Lunar iDXATM; GE Healthcare) were measured during the first habituation trial. Measurement of BM and body water content was repeated on all subsequent trial days. At the second habituation trial, subjects completed a familiarisation session on a cycle ergometer (Monark Ergonomic 874E; Monark Exercise AB) to confirm the steady-state work rate to be undertaken during the dehydration phase of the three experimental trials. The experimental trials were randomised and separated by 7 d. The experimental trial proceeded as follows: after provision of a urine sample, an 18G cannula was inserted into a superficial vein located at the antecubital fossa and a basal (resting) blood sample was drawn. Subjects then consumed a standardised breakfast comprising 40 g of cornflakes, full-fat milk (200 ml), a breakfast bar (37 g; Kellogg's[®] Nutri-Grain[®]; Kellogg Company) and plain water (100 ml) 1 h before the dehydration phase. Further blood samples were drawn post exercise and at 1, 2, 3, 4 and 5 h of the rehydration phase.

Dehydration phase

Before entering an environmental chamber maintained at 30 (SD 1.1)°C and 58 (SD 8)% relative humidity, subjects voided their bladder and BM was measured. Subjects then undertook a continuous bout of cycle exercise for 50 min, towelled dry and reweighed. Subjects returned to the environmental chamber to continue exercise for 10-min intervals until a BM loss of approximately 1.8% was achieved. Having showered, dried thoroughly and voided their bladder subjects were reweighed to determine total BM loss during the dehydration phase. Expired air was collected and analysed for the rate of VO₂ and carbon dioxide production (VCO₂) by indirect calorimetry (AMIS 2000; Innovision A/S) from which estimates of the rate of energy expenditure, carbohydrate and fat metabolism were calculated⁽²⁴⁾.

Rehydration phase

Rehydration commenced 25 min after the end of the dehydration phase and lasted for 5 h. For this phase, subjects remained seated in a laboratory (20.5 (SD 1.2)°C, 35 (SD 8)% relative humidity) and, other than the rehydration fluid, no food or drink was consumed. The rehydration fluid was randomly assigned and administered to subjects at room temperature (18–20°C). The composition of the rehydration fluids, that is, milk (0.1% skimmed milk; Glanbia Consumer Foods Ireland Plc; trial M), CE (Powerade; Coca Cola Ltd; trial CE) and water (Centra; Musgrave Retail Partners Ireland; trial W), is provided in Table 1.

Subjects ingested a volume (litres) equivalent to 150% of the BM loss (kg). Using a metered protocol, 1000 ml was provided

Table 1. Energy density, osmolality, protein, carbohydrate and fat content, sodium and potassium concentration of the rehydration fluids (Mean values and standard deviations)

	M		CE		W	
	Mean	SD	Mean	SD	Mean	SD
Energy density (kJ/l)	1450	0	700	0	0	0
Osmolality (mosmol/kg)	280	2	299	2	11	1
Protein (g/l)	33	0	0	0	0	0
Carbohydrate (g/l)	50	0	39	0	0	0
Fat (g/l)	1	0	0	0	0	0
Na (mmol/l)	17.9	0	21.7	0	0.7	0

M, milk; CE, carbohydrate–electrolyte solution; W, water.

to be ingested within the first 30 min and 500 ml every 30 min thereafter until the required volume of fluid was consumed. Subjects were allowed 15 min in which to consume each aliquot. All subjects completed the ingestion of the required volume of all fluids within the specified time. Subjects were instructed to empty their bladder as completely as possible at the end of each hour. The volume of urine provided was measured, and a 5 ml aliquot was retained for analysis. Any additional urine produced during the recovery period was collected and added to the bolus produced at the end of each hour. Subjective feelings relating to hydration (thirst, hunger, bloatedness, mouth taste, alertness, headache, tiredness and refreshment) were assessed by 100 mm visual analogue scales. Subjective responses for hydration were provided for the basal state, following the completion of the dehydration phase and at the end of each hour during the rehydration phase. Subjective responses relating to drink taste (visual appeal, palatability, aftertaste, sweetness and saltiness) were obtained for 2 h of the rehydration period.

Biochemical analysis

Whole-blood samples were analysed immediately after collection for Na and K concentration by ion-selective potentiometry (I-STAT[®]; Abbott Laboratories). Intra-assay CV using control solutions was <1% for both Na and K. Inter-assay CV was <1.5% for Na and <11% for K, using baseline samples. Plasma and urine osmolality was measured by freezing-point depression (Advanced Micro-Osmometer Model 3220; Advanced Instruments Inc.). The intra-assay CV for P_{osm} was 1.2%. CTproAVP was measured in a single batch with a commercial sandwich immunoluminometric assay (B.R.A.H.M.S LUMItest CTproAVP; B.R.A.H.M.S AG). The lower detection limit of this assay was 0.4 pM, and the functional assay sensitivity (<20% inter-assay CV) was <1 pM. The intra-assay CV was <10%.

Statistical analysis

Unless stated otherwise, data are presented as means and standard deviations. Statistical analysis was performed within the Statistical Package for the Social Sciences (SPSS, 21.0; SPSS Inc.). Data were checked for normality of distribution by Shapiro–Wilk test. Data containing two independent factors were analysed by two-factor (time × trial) repeated-measures

ANOVA. Variables containing one factor were analysed using a one-factor repeated-measures ANOVA. Bonferroni-adjusted paired sample Student's *t* test was used to compare the means and locate significant differences. The 95% level of confidence was the criterion of statistical difference ($P < 0.05$). Where statistical significance marginally exceeded 0.05 the effect size, Cohen's *d* was calculated to indicate the standardised difference between means. Friedman's test and *post hoc* Wilcoxon's signed-rank tests were used to compare differences between trials for non-parametric data relating to subjective responses rated by visual analogue scales.

Results

Basal hydration status

No difference was observed in basal BM, hydration status or blood electrolyte concentration for each trial (Table 2), indicating that the subjects complied fully with the prescribed pre-experimental protocol.

Dehydration phase

The dehydration protocol achieved a BM loss of −1.84 (SD 0.28) kg (W), −1.86 (SD 0.37) kg (CE) and −1.82 (SD 0.28) kg (M), which did not differ between trials ($P = 0.84$). This corresponded to −2.2 (SD 0.2)% (W), −2.2 (SD 0.2)% (CE) and −2.1 (SD 0.2)% (M) loss of the subjects' basal BM. The duration of exercise required to achieve this was not different between trials ($P = 0.16$): 94 (SD 17) min (W), 93 (SD 11) min (CE) and 101 (SD 16) min (M). Estimates of energy expenditure, carbohydrate and fat oxidation are provided in Table 3.

Rehydration phase

Aligned with the outcome of the dehydration phase, the volume of fluid ingested during the rehydration phase was not significantly different between trials ($P = 0.84$); 2764 (SD 416) ml (W), 2796 (SD 556) ml (CE) and 2732 (SD 426) ml (M). Significantly lower mean hourly urine volume was observed in the rehydration phase in M compared with CE and W (time × trial, $P = 0.01$), resulting in a lower cumulative urine volume in M compared with CE ($P = 0.013$) and W ($P = 0.007$) but no difference between CE and W ($P = 0.68$) (Fig. 1(a)). At 5 h, total

Table 2. Body mass, total body water, blood and urine osmolality and electrolyte concentration at basal (Mean values and standard deviations)

	M		CE		W		<i>P</i> *
	Mean	SD	Mean	SD	Mean	SD	
Body mass (kg)	85.6	11.5	86.2	11.6	85.7	11.0	0.56
Total body water (litres)	50.0	4.46	49.8	4.23	50.0	4.17	0.91
Plasma osmolality (mosmol/kg)	286	3.4	287	3.2	288	4.4	0.33
Plasma Na (mmol/l)	140.7	1.7	140.6	2.9	139.9	2.0	0.55
Plasma K (mmol/l)	3.9	0.1	4.0	0.3	3.9	0.3	0.95

M, milk; CE, carbohydrate–electrolyte solution; W, water.

* *P* values for the difference between groups analysed by univariate ANOVA.

Table 3. Estimate of energy expenditure, carbohydrate and fat oxidation during the exercise and dehydration phase (Mean values and standard deviations)

	M		CE		W		<i>P</i> *
	Mean	SD	Mean	SD	Mean	SD	
Total energy (kJ)	3730	663	3440	559	3877	950	0.44
Carbohydrate oxidation (g)	167	31	149	28	171	40	0.36
Fat oxidation (g)	28	6	29	5	31	9	0.63

M, milk; CE, carbohydrate–electrolyte solution; W, water.

* *P* values for the difference between groups analysed by univariate ANOVA.

urine volume in M (794 (SD 99) ml) was lower than W (1429 (SD 131) ml) ($P=0.018$) but not CE (1314 (SD 164) ml) ($P=0.057$; $d=1.7$). Percentage fluid retention reflected these differences between trials: 71 (SD 4)% (M), 52 (SD 16)% (CE, $P=0.048$) and 47 (SD 15)% (W, $P=0.014$).

Whole-body NFB was calculated from the change in BM (reflecting the loss of fluid by sweating), fluid ingested and urine produced. No significant difference in NFB was observed between trials following dehydration ($P=0.84$), but a significant difference in the time course of change in NFB was observed between trials (time \times trial, $P=0.003$) in the rehydration phase (Fig. 1(b)). A higher overall NFB was maintained throughout the rehydration phase in M than CE ($P=0.001$) and W ($P=0.006$), with no difference between CE and W ($P=0.69$). At 2 h into rehydration, NFB was positive compared with basal in M and significantly greater than CE ($P=0.031$) but not W ($P=0.09$, $d=1.38$). At 3 h, NFB remained positive in M and greater than W ($P=0.019$) but not CE ($P=0.069$, $d=1.21$), and this trend continued to 5 h, at which point NFB in M remained positive compared with basal and greater than W ($P=0.011$) but not CE ($P=0.077$, $d=1.6$).

Plasma osmolality and pre-provasopressin

The relationship between plasma osmolality (P_{osm}) and CTproAVP is reported in Fig. 2. Spearman's rank correlation revealed a moderate, but statistically significant ($P<0.0001$), association between P_{osm} and serum CTproAVP concentration (Fig. 2(a)) and between the change in CTproAVP induced by a change in P_{osm} (Fig. 2(b)). A threshold of effect on serum CTproAVP is observed when P_{osm} increased above

approximately 290 mosmol/kg, which approximates to approximately 3 mosmol/kg increase in P_{osm} from basal values.

P_{osm} was similar for all trials at basal (288 (SD 4.5) mosmol/kg, W; 287 (SD 3.2) mosmol/kg, CE; 288 (SD 3.4) mosmol/kg, M, $P=0.33$) and increased following dehydration, with no significant difference between trials ($P=0.22$) (online Supplementary Fig. S1(a)). A significant difference in the time course of change in P_{osm} during rehydration was observed between trials (time \times trial, $P=0.003$) (Fig. 3(a)), remaining elevated above basal levels in trial M compared with trial CE and W ($P=0.001$). At 1 h, P_{osm} remained elevated above basal in M compared with CE ($P=0.016$) but not W ($P=0.051$, $d=1.74$); at 2 h, M was greater than CE ($P=0.002$) and W ($P=0.002$); at 3 h, M was greater than W ($P=0.04$) but not CE ($P=0.052$, $d=1.13$); at 4 h, M was greater than CE ($P=0.046$) and W ($P=0.001$) and CE was greater than W ($P=0.037$). At the end of the 5-h rehydration period, P_{osm} returned to within basal levels for M 288 (SD 3.9) mosmol/kg, CE 287 (SD 2.6) mosmol/kg and W 286 (SD 1.5) mosmol/kg, with no difference ($P=0.13$) between trials.

No significant difference in the time course of change in serum CTproAVP was observed between trials (time \times treatment, $P=0.96$) (online Supplementary Fig. S1(b)). Serum CTproAVP was similar for all trials at basal (4.6 (SD 1.2) pmol/l, W; 4.3 (SD 1.6) pmol/l, CE; 4.1 (SD 1.8) pmol/l, M, $P=0.73$) and increased approximately 5-fold from basal in response to dehydration to 29.1 (SD 26.9) pmol/l (W), 23.8 (SD 13.5) pmol/l (CE) and 27.8 (SD 13.4) pmol/l (M) ($P=0.72$) (Fig. 3(b)). CTproAVP decreased during rehydration to values that were not significantly different from basal at 1 h ($P>0.05$), and remained unchanged thereafter.




Fig 1. Urinary volume (a) and net fluid balance (b) over the course of the experimental trials. Values are means and standard deviations represented by vertical bars. * Value of milk (M) significantly different from carbohydrate-electrolyte solution (CE) and water (W) ($P < 0.05$). † Value of M is significantly different from W ($P < 0.05$). ○—, Trial M; ●—, trial CE; △—, trial W.

Plasma electrolytes

Plasma Na was similar for all trials at basal (140 (SD 2.0) mmol/l, W; 141 (SD 2.9) mmol/l, CE; 141 (SD 1.7) mmol/l, M, $P = 0.55$), and no significant difference in plasma Na in response over the duration of the experiment was observed between trials (time \times trial, $P = 0.22$) (online Supplementary Fig. S2(a)) (Fig. 4(a)). Basal plasma K was similar for all trials (3.93 (SD 0.30) mmol/l, W; 3.96 (SD 0.32) mmol/l, CE; 3.91 (SD 0.15) mmol/l, M, $P = 0.95$), increasing by 0.25 (SD 0.22) mmol/l following dehydration, with no significant difference in response between trials ($P = 0.71$) (online Supplementary Fig. S2(a)). A significant difference in plasma K during rehydration was observed between trials (time \times treatment, $P < 0.001$), remaining elevated above basal for the 5-h rehydration period in M and greater than in trial CE ($P = 0.006$) and W ($P = 0.008$), with no difference in response between CE and W ($P = 0.11$). At 1 h into rehydration, K remained significantly elevated above basal in M compared with CE ($P = 0.001$) but not W ($P = 0.063$, $d = 1.74$); at 2 h, M was greater than CE ($P = 0.005$) and W ($P = 0.021$); at 3 h, M was greater than CE ($P = 0.024$) and W ($P = 0.013$); at 4 h, M was greater than CE ($P = 0.004$) and




Fig 2. Relationship between (a) plasma osmolality and pre-provasopressin (CTproAVP) concentration and (b) change in pre-provasopressin (Δ CTproAVP) measured over the course of the experimental trials shown as scatter plots. r_s , Spearman's rank correlation coefficient ($n = 126$); ○, trial milk; ●, trial carbohydrate-electrolyte solution; △, trial water.

W ($P = 0.022$) and greater in M compared with CE ($P = 0.003$) but not W ($P = 0.88$) at 5 h (Fig. 4(b)).

Subjective feelings of hydration and drink acceptance

There was a marked increase in perceived thirst score immediately post exercise for all trials, although no overall difference in change in thirst perception scores was observed between test drinks during the rehydration period ($P = 0.53$). Ingestion of a rehydration fluid reduced the feeling of thirst from a pooled mean of 81 (SD 6) to 45 (SD 4) mm and gradually returned to basal feelings of thirst by the end of the 5-h rehydration period for all three trials: M 51 (SD 26) mm ($P = 0.67$); CE 50 (SD 24) mm ($P = 0.79$); and W trials 51 (SD 27) mm ($P = 0.39$). Participants scored mouth taste lowest in W trial (14 (SD 4) mm) compared with CE (35 (SD 13) mm) ($P = 0.034$) and M (44 (SD 8) mm) ($P = 0.013$). Perceived feeling of hunger differed between test drinks during rehydration (time \times trial) ($P = 0.037$), with a lower overall mean score in the M trial 51 (SD 9) mm compared with W 79 (SD 9) mm, and CE 66 (SD 14) mm. Similar perceived levels of bloatedness ($P = 0.15$), alertness ($P = 0.28$), tiredness ($P = 0.063$), headache ($P = 0.16$) and level of refreshment ($P = 0.06$) were reported for all rehydration fluids. Subjective feelings associated with drink taste were analysed for the 2nd hour of fluid ingestion. There was no difference in perceived visual appeal ($P = 0.54$), palatability ($P = 0.57$) or aftertaste ($P = 0.06$)




Fig 3. Change in plasma osmolality (ΔP_{osm}) (a) and serum pre-provasopressin ($\Delta CTproAVP$) (b) from basal values over the course of the experimental trials. Values are the means and standard deviations represented by vertical bars. * Value of milk (M) significantly different from carbohydrate–electrolyte solution (CE) and water (W) ($P < 0.05$). † Value of M significantly different from W ($P < 0.05$). ‡ Value of M significantly different from CE ($P < 0.05$). § Value of CE significantly different from W ($P < 0.05$). —○—, Trial M; —●—, trial CE; —△—, trial W.

between trials. There was a higher perceived sweetness for CE compared with M ($P = 0.048$) and W ($P = 0.018$). Perceived saltiness also scored highest in the CE trial – 29 (SD 28) mm compared with M 18 (SD 30) mm ($P = 0.046$) and W 3 (SD 3) mm ($P = 0.028$) trials, although overall scores were low across trials.

Energy and substrate intake in comparison with expenditure

A comparison of the total energy, carbohydrate and fat intake provided in the rehydration phase for M and CE is provided in Table 4. The mean total energy provision was approximately 2-fold greater for M than CE, the difference being less because of carbohydrate (approximately 40% greater in M) than for the additional protein and fat provided in M. Relative to the mean estimates of energy expenditure, M provided 106% of total energy, 82% of total carbohydrate and 10% of total fat oxidised during exercise. In comparison, CE provided 57% of total energy and 73% of total carbohydrate used during exercise.




Fig 4. Change in plasma sodium (a) and plasma potassium (b) from basal values over the course of the experimental trials. Values are the means and standard deviations represented by vertical bars. * Value of milk (M) significantly different from carbohydrate–electrolyte solution (CE) and water (W) ($P < 0.05$). † Value of M significantly different from CE ($P < 0.05$). ‡ Value of CE significantly different from W ($P < 0.05$). —○—, Trial M; —●—, trial CE; —△—, trial W.

Table 4. Estimate of energy, carbohydrate and fat provided during the rehydration phase (Mean values and standard deviations)

	M		CE		<i>P</i> *
	Mean	SD	Mean	SD	
Total energy (kJ)	3962	234	1958	147	<0.001
Carbohydrate (g)	137	8	109	8	<0.001
Fat (g)	2.7	0.2	0	0	<0.001
Protein (g)	90	5.3	0	0	<0.001

M, milk; CE, carbohydrate–electrolyte solution.
* *P* values for the difference between groups analysed by paired sample *t* test.

Discussion

The purpose of a metered rate of fluid ingestion was to optimise the inter-related processes of gastric emptying, intestinal absorption and retention of the rehydration fluid. The rate of gastric emptying is related to rate of fluid ingestion that is

modulated by the energy density and osmolality of the fluid, with energy density exerting a greater influence than osmolality^(25,26). M contains approximately double the energy density compared with the CE, and rehydration studies using fluids containing milk protein confirm a slowing in the rate of gastric emptying and a delay in the rate of intestinal water absorption into the circulation⁽²⁶⁾. Once fluid is absorbed and released into the circulation, the magnitude and time course of change in P_{osm} is a key determinant of water retention. The loss of plasma volume and associated increase in P_{osm} during dehydration was associated with a 5-fold increase in CTproAVP, a marker of the antidiuretic hormone AVP. CTproAVP is produced during synthesis of mature AVP in a 1:1 ratio as a stable peptide, sensitive to measurement by specific immunoassay and representative of the antidiuretic effect of AVP⁽²⁷⁾. Analysed over all samples and all time points, a highly significant association between P_{osm} and CTproAVP (r_s 0.348; $P < 0.001$) was observed. The mean increase in CTproAVP across all trials in response to exercise and thermal-induced dehydration was 5-fold the basal level but, as shown in Fig. 2, plasma CTproAVP concentration at a given osmolality varied widely between individuals, particularly at osmolalities above 287 mosmol/kg, and highlights the polygenetic variability of this response^(28,29). However, despite the continued elevation in P_{osm} in the early phase of rehydration in M, CTproAVP returned to basal levels within 1 h of rehydration in all trials.

An increase in AVP is predicted to promote a compensatory change in renal water conservation and decrease in water excretion subsequently potentiated by ingestion of the rehydration fluid. Ingestion of W caused the largest decrease in P_{osm} (approximately 12 mosmol/l) from 295 (SD 2.2 mosmol/l) following dehydration to 283 (SD 2.9) mosmol/l at 2 h rehydration. The response to ingestion of CE did not differ significantly from W, but the outcome in M was to maintain an elevated P_{osm} of +5 mosmol/l during the early phase of rehydration (Fig. 3). The maintenance of an elevated P_{osm} in M might be predicted by the protein content in M, the higher energy density resulting in a decrease in the rate of gastric emptying⁽²⁶⁾. In addition, the protein in M could augment the osmotic/oncotic pressure via an increase in plasma albumin and/or post ingestion increase in plasma amino acid concentration^(19,21). Plasma albumin is the major contributor to oncotic pressure that trends towards an increase over time when milk-based proteins are added to a carbohydrate beverage⁽³⁰⁾. On the basis of pilot data (unpublished results), total free amino acid concentration following ingestion of M or CE increases approximately 500 $\mu\text{mol/l}$ in M and decreases approximately 400 $\mu\text{mol/l}$ in CE, 1 h post ingestion. Acting independently, the change in plasma albumin and amino acid concentration of this magnitude may be too low to alter serum osmolality, but could act in synergy to augment NFB.

Analysed across all trials, no effect of the experimental protocol on plasma Na was observed (online Supplementary Fig. S2(a)). Considering the strong influence that Na and chloride, its principal anion, exerts on P_{osm} , these data are perplexing, possibly reflecting the relative imprecision offered by the point-of-care platform of measurement. In contrast, a significant increase in plasma K was observed over the

rehydration phase in M (online Supplementary Fig. S2(b)). The maintenance of plasma K in M relative to CE and W could have contributed to water retention via its effect on the renin-angiotensin-aldosterone axis. Controlled infusion of potassium chloride results in an increase in circulating aldosterone; the response is most sensitive to an increase in plasma K within a range of 0.1–0.5 mmol/l⁽³¹⁾ – that is, of comparable magnitude to that observed in the present study.

The main considerations for an athlete in recovery from exercise are the replacement of fluid and electrolytes lost in sweat, repletion of body energy reserves, principally muscle and liver glycogen, and the provision of nutrient support to the regeneration or adaptation of skeletal muscle⁽³²⁾. In recovery, M provided 106% of total energy and 82% of total carbohydrate oxidised during exercise compared with 57% of total energy and 73% of total carbohydrate provided by CE. The recommended intake of carbohydrate in recovery is 1.0–1.2 g/kg per h to be consumed at frequent intervals during the first 4–6 h post exercise⁽³²⁾. The mean intake for M (1.6 g/kg per 5 h) was higher than CE (1.3 g/kg per 5 h), but both rehydration solutions fell short of the recommended recovery rate for carbohydrate intake. Although protein intake in recovery may enhance glycogen resynthesis, particularly in the early (0–2 h) phase or when the carbohydrate intake is sub-optimal⁽³³⁾, the primary effect of adding protein to a recovery fluid, particularly when consumed in close proximity to exercise, would be to stimulate skeletal muscle protein synthesis (MPS)⁽³⁴⁾. The mean protein intake in M was 90 g, equivalent to 1.1 g/kg BM, and greater than the recommended intake of protein to stimulate MPS post exercise⁽³²⁾. Furthermore, the EAA composition of milk proteins approximates to 50%. EAA, particularly leucine, are the main nutrient regulators of MPS, which lends support to the use of milk protein as a natural and effective source of protein for recovery⁽³⁵⁾. However, in practice, an athlete would probably use a recovery beverage as a supplement to habitual dietary intake, and the present investigation did not address the effect of habitual dietary intake in combination with a recovery beverage, and it is probable that solid food and beverage intake would influence the temporal pattern of restoration of body NFB⁽³⁶⁾. However, when time between exercise bouts is short (<6 h), the potential for gastrointestinal disturbance in the following training session is an important consideration in the recovery plan and a liquid recovery strategy may be more acceptable.

Finally, gastrointestinal disturbance is a relevant consideration during recovery post exercise, particularly when large volumes of nutrient-dense fluids, such as M, are consumed. Although published evidence is lacking⁽³⁷⁾, a metered rate of fluid ingestion is considered to reflect, more closely, *ad libitum* drinking patterns, avoiding over-distension of the stomach and potential for negative gastrointestinal effects, such as bloatedness. Subjects were screened for lactose intolerance, as those with lactose intolerance generally can only tolerate relatively small dose of lactose approximately 12 g (approximately 200 ml of milk)⁽³⁸⁾. However, large lactose loads can result in gastrointestinal disturbances such as bloating and diarrhoea that may be further exacerbated in those with undiagnosed lactase deficiency. Possibly reflecting the lower

rate of lactose load, the subjective feeling of bloatedness post ingestion of milk in the first 2 h of rehydration in this study scored relatively low (≤ 34 mm) compared with previous studies using a non-metered approach⁽⁵⁾.

In conclusion, it was possible to demonstrate a significant advantage in the restoration of body NFB over a 5-h period following exercise and thermal dehydration to -2% BM by a metered replacement of M compared with CE or W. Consumed as the sole recovery diet, neither M nor CE provided optimal nutrition for recovery. However, the protein component of M, essentially absent in CE and W, may further enhance glycogen resynthesis and promote MPS. In addition, the minerals (Ca, P, Zn, I) and vitamins (A, E, B group vitamins) in M that are not present in CE contribute to micro-nutrient requirements of the athlete.

Acknowledgements

The authors would like to thank Gavin Connolly, Alexandra Cremona, Katie Hughes, Robert Davies and Beate Gilson for their assistance in the data collection and the subjects for providing their time and commitment to the project.

This work was supported by the National Dairy Council, Ireland. The National Dairy Council, Ireland had no role in the design, analysis or writing of this article.

S. S. and P. J. conceived the study design. S. S. conducted the study and managed the data collection. S. S. and P. J. undertook the data analysis and co-authored the manuscript.

Supported by National Dairy Council Research Scholarship to S. S.

Supplementary material

For supplementary material/s referred to in this article, please visit <http://dx.doi.org/10.1017/S0007114516002907>

References

- Cheuvront SN, Ely BR, Kenefick RW, *et al.* (2010) Biological variation and diagnostic accuracy of dehydration assessment markers. *Am J Clin Nutr* **92**, 565–573.
- Goulet DB (2013) Effect of exercise-induced dehydration on endurance performance: evaluating the impact of exercise protocols on outcomes using a meta-analytic procedure. *Br J Sports Med* **47**, 679–686.
- Cheuvront SN & Kenefick RW (2014) Dehydration: physiology, assessment, and performance effects. *Compr Physiol* **4**, 257–285.
- Wall BA, Watson G, Peiffer JJ, *et al.* (2015) Current hydration guidelines are erroneous: dehydration does not impair exercise performance in the heat. *Br J Sports Med* **49**, 1077–1083.
- Kasperek GJ & Snider RD (1987) Effect of exercise intensity and starvation on activation of branched-chain keto acid dehydrogenase by exercise. *Am J Physiol* **252**, E33–E37.
- Shirreffs SM, Watson P & Maughan RJ (2007) Milk as an effective post-exercise rehydration drink. *Br J Nutr* **98**, 173–180.
- Watson P, Love TD, Maughan RJ, *et al.* (2008) A comparison of the effects of milk and a carbohydrate-electrolyte drink on the restoration of fluid balance and exercise capacity in a hot, humid environment. *Eur J Appl Physiol* **104**, 633–642.
- Shirreffs SM, Taylor AJ, Leiper JB, *et al.* (1996) Post-exercise rehydration in man: effects of volume consumed and drink sodium content. *Med Sci Sports Exerc* **28**, 1260–1271.
- Shirreffs SM & Maughan RJ (1998) Volume repletion after exercise-induced volume depletion in humans: replacement of water and sodium losses. *Am J Physiol* **274**, F868–F875.
- Sawka MN, Burke LM, Eichner ER, *et al.* (2007) American college of sports medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc* **39**, 377–390.
- Nose H, Mack GW, Shi X, *et al.* (1988) Role of osmolality and plasma volume during rehydration in humans. *J Appl Physiol* (1985) **65**, 325–331.
- Maughan RJ, Shirreffs SM & Leiper JB (1994) Post-exercise rehydration in man: effects of electrolyte addition to ingested fluids. *Eur J Appl Physiol Occup Physiol* **69**, 209–215.
- Merson SJ, Maughan RJ & Shirreffs SM (2008) Rehydration with drinks differing in sodium concentration and recovery from moderate exercise-induced hypohydration in man. *Eur J Appl Physiol* **103**, 585–594.
- Nielsen B, Sjogaard G, Ugelvig J, *et al.* (1986) Fluid balance in exercise dehydration and rehydration with different glucose-electrolyte drinks. *Eur J Appl Physiol Occup Physiol* **55**, 318–325.
- Evans GH, Shirreffs SM & Maughan RJ (2009) Postexercise rehydration in man: the effects of osmolality and carbohydrate content of ingested drinks. *Nutrition* **25**, 905–913.
- Osterberg KL, Pallardy SE, Johnson RJ, *et al.* (2010) Carbohydrate exerts a mild influence on fluid retention following exercise-induced dehydration. *J Appl Physiol* (1985) **108**, 245–250.
- Clayton DJ, Evans GH & James LJ (2014) Effect of drink carbohydrate content on postexercise gastric emptying, rehydration, and the calculation of net fluid balance. *Int J Sport Nutr Exerc Metab* **24**, 79–89.
- Seifert J, Harmon J & DeClercq P (2006) Protein added to a sports drink improves fluid retention. *Int J Sport Nutr Exerc Metab* **16**, 420–429.
- James LJ, Clayton D & Evans G H (2011) Effect of milk protein addition to a carbohydrate-electrolyte rehydration solution ingested after exercise in the heat. *Br J Nutr* **105**, 393–399.
- James LJ, Evans GH, Madin J, *et al.* (2013) Effect of varying the concentrations of carbohydrate and milk protein in rehydration solutions ingested after exercise in the heat. *Br J Nutr* **110**, 1285–1291.
- James LJ, Mattin L, Aldiss P, *et al.* (2014) Effect of whey protein isolate on rehydration after exercise. *Amino Acids* **46**, 1217–1224.
- Jones EJ, Bishop PA, Green JM, *et al.* (2010) Effects of metered versus bolus water consumption on urine production and rehydration. *Int J Sport Nutr Exerc Metab* **20**, 39–44.
- Euromonitor (2014) Sports and energy drinks in Ireland. <http://www.euromonitor.com/sports-and-energy-drinks-in-ireland/report> (accessed September 2014).
- Frayn KN (1983) Calculation of substrate oxidation rates *in vivo* from gas exchange. *J App Physiol Respir Environ Exerc Physiol* **55**, 628–634.
- Vist GE & Maughan RJ (1995) The effect of osmolality and carbohydrate content on the rate of gastric emptying of liquids in man. *J Physiol* **486**, 523–531.
- Calbet JA & MacLean DA (1997) Role of caloric content on gastric emptying in humans. *J Physiol* **498**, 553–559.
- Balancescu S, Kopp P, Gaskill MB, *et al.* (2011) Correlation of plasma copeptin and vasopressin concentrations in hypo-, iso-, and hyperosmolar states. *J Clin Endocrinol Metab* **96**, 1046–1052.

28. Cheuvront SN, Kenefick RW, Charkoudian N, *et al.* (2013) Physiologic basis for understanding quantitative dehydration assessment. *Am J Clin Nutr* **97**, 455–462.
29. Zerbe RL, Miller JZ & Robertson GL (1991) The reproducibility and heritability of individual differences in osmoregulatory function in normal human subjects. *J Lab Clin Med* **117**, 51–59.
30. Hobson R & James L (2015) The addition of whey protein to a carbohydrate–electrolyte drink does not influence post-exercise rehydration. *J Sports Sci* **33**, 77–84.
31. Himathongkam T, Dluhy RG & Williams GH (1974) Potassium-aldosterone-renin interrelationships. *J Clin Endocrinol Metab* **41**, 153–159.
32. Thomas DT, Erdman KA & Burke LM (2016) Position of the Academy of Nutrition and Dietetics, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and Athletic Performance. *J Acad Nutr Diet* **116**, 501–528.
33. Ivy JL, Goforth HW Jr, Damon BM, *et al.* (2002) Early post-exercise muscle glycogen recovery is enhanced with a carbohydrate-protein supplement. *J Appl Physiol (1985)* **93**, 1337–1344.
34. Tipton KD, Rasmussen BB, Miller SL, *et al.* (2001) Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab* **281**, E197–E206.
35. Tipton KD, Elliott TA, Cree MG, *et al.* (2007) Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *Am J Physiol Endocrinol Metab* **292**, E71–E76.
36. Maughan RJ, Leiper JB & Shirreffs SM (1977) Factors influencing the restoration of fluid and electrolyte balance after exercise in the heat. *Br J Sports Med* **31**, 175–182.
37. Wong SH, Williams C, Simpson M, *et al.* (1998) Influence of fluid intake pattern on short-term recovery from prolonged, submaximal running and subsequent exercise capacity. *J Sports Sci* **16**, 143–152.
38. National Centre for Biotechnology Information (2010) Lactose intolerance and health, Evidence Reports/Technology Assessments No. 192. Bethesda, MD. <http://www.ncbi.nlm.nih.gov/books/NBK44616/> (accessed September 2014).