




RESEARCH ARTICLE

Equivariant geometry of singular cubic threefolds

Ivan Cheltsov ¹, Yuri Tschinkel ^{2,3} and Zhijia Zhang ⁴

¹Department of Mathematics, University of Edinburgh, UK; E-mail: I.Cheltsov@ed.ac.uk (corresponding author).

²Courant Institute, 251 Mercer Street, New York, NY, 10012, USA; E-mail: tschinkel@cims.nyu.edu.

³Simons Foundation, 160 Fifth Avenue, New York, NY, 10010, USA.

⁴Courant Institute, 251 Mercer Street, New York, NY, 10012, USA; E-mail: zz1753@nyu.edu.

Received: 23 March 2024; **Revised:** 23 August 2024; **Accepted:** 17 November 2024

2020 Mathematical Subject Classification: *Primary* – 14E07; *Secondary* – 14E30

Abstract

We study linearizability of actions of finite groups on singular cubic threefolds, using cohomological tools, intermediate Jacobians, Burnside invariants, and the equivariant Minimal Model Program.

1. Introduction

In this paper, we continue our investigations of actions of finite groups on rational threefolds over an algebraically closed field k of characteristic zero, up to equivariant birationality. The main problem is to decide *linearizability* (i.e., birationality of the given action to a *linear* action on projective space); see, for example, [22] for background and references. The linearizability problem is essentially settled in dimension 2 [36, 50] but remains largely open in dimension 3. Here, we focus on the following:

Problem 1. Let $X \subset \mathbb{P}^4$ be a singular rational cubic threefold and let G be a finite subgroup of its automorphisms. When is the G -action on X linearizable?

Note that linearizability of a G -action for a cubic threefold X is equivalent to projective linearizability since the action lifts to GL_5 (see Section 2 for a proof, and [42] for a general discussion of these notions).

Smooth cubic threefolds are not rational, and their automorphisms have been classified in [57, Theorem 1.1]: there are six maximal groups

$$C_3^4 \rtimes \mathfrak{S}_5, ((C_3^2 \rtimes C_3) \rtimes C_4) \times \mathfrak{S}_3, C_{24}, C_{16}, \mathrm{PSL}_2(\mathbb{F}_{11}), C_3 \times \mathfrak{S}_5.$$

However, all singular ones, except cones over smooth cubic curves, are rational. Cubic threefolds with isolated singularities have been classified in [55], but it is not immediately clear how to identify possible symmetries from that analysis.

Here, we restrict our attention to nodal cubics (i.e., those with ordinary double points), as this is the most interesting and difficult class of singular cubics from the perspective of equivariant geometry. In all these cases, the automorphism group $\mathrm{Aut}(X)$ is finite, by, for example, [2, Theorem 1.1].

Note that the existence of a G -fixed node yields a straightforward linearization construction: projection from this node gives an equivariant birational map to \mathbb{P}^3 with linear action. Thus, we will be primarily interested in actions not fixing a singular point of X . Another such construction comes from a G -stable plane and a disjoint G -stable line:

Lemma 1.1. *Let X be a nodal cubic threefold. Let $G \subseteq \text{Aut}(X)$ be such that it preserves a plane $\Pi \subset X$ and a line $l \subset X$, disjoint from Π . Then the G -action on X is linearizable.*

Proof. Let $\phi : X \dashrightarrow X_{2,2}$ be the unprojection from Π ; $X_{2,2} \subset \mathbb{P}^5$ is a (nodal) complete intersection of two quadrics. Then ϕ is G -equivariant, and $\phi(l)$ is a G -invariant line. Taking a projection from this line, we obtain a G -equivariant birational map $X \dashrightarrow \mathbb{P}^3$; see [29, Proposition 2.2] for an application of this construction over nonclosed fields. \square

Let $s = s(X)$ be the cardinality of the set $\text{Sing}(X)$ of nodes of X . It is well known that $s \leq 10$. Moreover, there is a unique cubic threefold X with $s = 10$, the Segre cubic, treated in [5, 24]. In [5], it is shown that the subgroup $\mathfrak{A}_5 \subset \mathfrak{S}_6 = \text{Aut}(X)$ that leaves invariant a hyperplane section is not linearizable. In [24], we have completed this analysis by proving that the action of $G \subseteq \mathfrak{S}_6$ is linearizable if and only if

- G fixes a singular point of X , or
- G is contained in the subgroup $\mathfrak{S}_5 \subset \mathfrak{S}_6$ that does not leave invariant a hyperplane section of X , or
- $G \simeq C_2^2$, X contains three G -invariant planes, and $\text{Sing}(X)$ is a union of five G -orbits of length 2.

Using this description, one can list all subgroups of \mathfrak{S}_6 giving rise to linearizable actions – there are 37 such subgroups up to conjugation (among 56 conjugacy classes of subgroups of \mathfrak{S}_6).

In this paper, we study the cases where

$$2 \leq s(X) \leq 9.$$

To address the linearizability problem for these outstanding cases, we use explicit geometric constructions, as well as the following techniques:

- cohomological tools [14, 45],
- intermediate Jacobians, in the equivariant context [10],
- Burnside invariants and their specialization [46],
- G -birational rigidity and G -solidity; see, for example, [20].

To describe our results, we distinguish cases based on linear position properties of nodes, following [37]. According to [37], there are 15 configurations, labeled (J1), ..., (J15), with (J1), ..., (J5) corresponding to 1, ..., 5 nodes in general linear position, and (J15) corresponding to the Segre cubic threefold. The relevant invariants are

- s , the number of nodes of X ,
- $d = \text{rk Cl}(X) - 1$, the defect of X , which equals the number of dependent linear conditions imposed on $H^0(X, \mathcal{O}_X(1))$ by the nodes, and
- p – the number of planes $\Pi \subset X$.

We list all possibilities for the triples (s, d, p) and describe our results in each of the cases:

- $s = 2, d = 0, p = 0$: We prove that the G -action is linearizable if and only if G fixes each of the two nodes, and classify actions of cyclic groups not fixing any node; see Section 3.
- $s = 3, d = 0, p = 0$: We conjecture that the G -action is linearizable if and only if it fixes a node, and classify all automorphism groups not fixing any node. We provide examples of nonlinearizable actions of $G = C_3^2$; see Section 4.
- $s = 4$:
 - $d = 0, p = 0$: There is an equivariant birational map to a smooth divisor of degree $(1, 1, 1, 1)$ in $(\mathbb{P}^1)^4$. Following considerations over nonclosed fields in [48, Conjecture 1.3], we conjecture that the G -action is not linearizable if it is transitive on the nodes. We provide examples of nonlinearizable actions of $G = C_2^2$ confirming this conjecture. We classify all automorphism groups not fixing any node.
 - $d = 1, p = 1$: An action is nonlinearizable if and only if it does not fix a node and X does not contain G -stable lines disjoint from the unique plane in X ; see Section 5.

- $s = 5$:
 - $d = 0, p = 0$: All actions are linearizable, except for actions of \mathfrak{A}_5 and \mathfrak{S}_5 , which are birational to standard actions on a smooth quadric threefold. The \mathfrak{S}_5 -action is not linearizable [21]; we conjecture that the \mathfrak{A}_5 -action is also not linearizable; see Section 6.
 - $d = 1, p = 1$: All actions are linearizable, as there is a unique node outside the plane and fixed by the action [37].
- $s = 6$:
 - $d = 1, p = 0$: We classify all automorphism groups not fixing any node, establish a sufficient condition for nonlinearizability, and apply it to provide examples of nonlinearizable actions of $G = C_2$.
 - $d = 1, p = 1$: All actions are linearizable by Lemma 1.1.
 - $d = 2, p = 3$: We classify all actions and solve the linearizability problem for most of them; see Section 7.
- $s = 7$:
 - $d = 2, p = 2$: All actions are linearizable, since each of the two planes contains four nodes, and exactly one of the nodes is on both planes, thus preserved by the action [37].
 - $d = 2, p = 3$: All actions are linearizable, since there is a unique node not contained in any plane in X , thus fixed by the action [37].
- $s = 8, d = 3, p = 5$: We classify automorphism groups not fixing any node and solve the linearizability problem in Section 8.
- $s = 9, d = 4, p = 9$: Linearizability problem is solved, except for specific actions of \mathfrak{S}_3 and \mathfrak{D}_6 , which are birational to actions on a smooth quadric; see Section 9.

We conclude the introduction by summarizing the cases for which the linearizability problem remains open.

- $s = 3, d = 0, p = 0$: Actions in Proposition 4.1 not containing the C_3^2 in Example 4.2 and not fixing any node.
- $s = 4, d = 0, p = 0$: Actions in Theorem 5.1 not containing the C_2^2 in Example 5.2 and not fixing any node.
- $s = 5, d = 0, p = 0$: A unique \mathfrak{A}_5 -action described in Section 6, which is equivariantly birational to the \mathfrak{A}_5 -action on a smooth quadric (14).
- $s = 6, d = 1, p = 0$: Actions in Proposition 7.3 not fixing any node and not containing an involution not fixing any nodes.
- $s = 6, d = 2, p = 3$: Actions in Proposition 7.11 not containing the C_2^2 in Lemma 7.14, not containing the C_2^2 or \mathfrak{S}_3 in Remark 7.15, not contained in the C_2^2 in Lemma 7.13, and not fixing any node.
- $s = 9, d = 4, p = 9$: The actions of \mathfrak{D}_6 , \mathfrak{S}_3 and \mathfrak{S}_3' specified in (26); these are also equivariantly birational to actions on a smooth quadric (27).

In many of these cases, equivariant specialization of [46], applied here in the geometric context for the first time, shows nonlinearizability of the actions for a very general member of the family; see Propositions 4.3, 5.3, 5.4, Lemmas 7.17, 7.18 and Remark 7.19.

2. Obstructions to linearizability

Among available obstruction theories to linearizability are

- Existence of fixed points upon restriction to abelian subgroups,
- Group cohomology,
- Intermediate Jacobians, and their equivariant versions,
- Burnside invariants,
- Specialization of birational types,
- Birational rigidity.

We briefly review relevant results and constructions.

Fixed points by abelian subgroups

Recall that existence of fixed points for actions of abelian groups is an equivariant birational invariant; see [51]. Precisely, let G be a finite abelian group acting generically freely on a smooth projective variety V . Assume that there exists a G -equivariant birational map $W \dashrightarrow V$ from a smooth G -variety W . Then

$$W^G \neq \emptyset \iff V^G \neq \emptyset.$$

Linear actions of abelian groups on projective spaces always have fixed points, and thus, we have the following.

Lemma 2.1. *Let V be a smooth projective variety with generically free and linearizable action by a finite group G . Then $V^H \neq \emptyset$ for all abelian subgroups $H \subseteq G$.*

In particular, let X be a nodal cubic threefold and $G \subseteq \text{Aut}(X)$. The G -action on X is linearizable if and only if it is projectively linearizable. To see this, one can apply the argument in [38]. Alternatively, we provide a direct proof below.

First, we show that the G -action is induced from an action of the ambient \mathbb{P}^4 (i.e., $G \subset \text{PGL}_5$). Indeed, by Lefschetz hyperplane theorem, the Picard group $\text{Pic}(X) = \mathbb{Z}$ is generated by a general hyperplane section of X . The induced G -action on $\text{Pic}(X)$ is trivial, sending a hyperplane to another hyperplane in \mathbb{P}^4 . This implies that the G -action on X lifts to \mathbb{P}^4 .

It follows that the G -action naturally lifts to $\mathcal{O}_X(-5)$, the restriction of the canonical bundle of \mathbb{P}^4 to X . Similarly, the G -action lifts to $\mathcal{O}_X(-2)$, the canonical bundle of X . Since $\mathcal{O}_X(-5)$ and $\mathcal{O}_X(-2)$ generate $\mathcal{O}_X(1)$, we know that the G -action lifts to $\mathcal{O}_X(1)$, and thus to $H^0(X, \mathcal{O}_X(1))$ and $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(1))$. Therefore, the G -action lifts to GL_5 .

This also shows that the Amitsur group $\text{Am}(X, G)$ (see [13, Section 6]) is trivial. If the G -action is projectively linearizable (i.e., equivariantly birational to a G -action on \mathbb{P}^3), then $\text{Am}(\mathbb{P}^3, G) = 0$ since the Amitsur group is an equivariant birational invariant. This implies that the G -action on \mathbb{P}^3 is linear; namely, it lifts to GL_4 . So the notions of linearizable and projectively linearizable actions on X are equivalent.

Thus, if an abelian subgroup $H \subseteq G$ does not fix a point in the standard desingularization \tilde{X} of X (the blowup of the nodes), then there exists no G -equivariant birational map $X \dashrightarrow \mathbb{P}^3$. We found two applications of this obstruction; see Example 4.2 and Section 9.

Cohomology

Let X be a nodal cubic threefold, $\tilde{X} \rightarrow X$ its standard desingularization, and $G \subseteq \text{Aut}(X)$. Here, we consider the induced G -actions on the Picard group $\text{Pic}(\tilde{X})$ and the class group $\text{Cl}(X)$; we often identify divisors and their classes.

A well-studied obstruction to (stable) linearizability is the failure of $\text{Pic}(\tilde{X})$ to be a *stably permutation module*; we call this the **(SP)**-obstruction. In turn, this is implied by the nonvanishing of

$$H^1(G', \text{Pic}(\tilde{X})), \quad \text{or} \quad H^1(G', \text{Pic}(\tilde{X})^\vee), \quad \text{for some } G' \subseteq G.$$

We call this the **(H1)**-obstruction; see [24, Section 2].

Proposition 2.2. *When $s(X) \leq 7$ and $s(X) \neq 6$, or when $s = 6$ and the nodes are not in general linear position, $\text{Pic}(\tilde{X})$ is a permutation module.*

Proof. We use labels for configurations of nodes and planes from [37].

- o $s = 1, \dots, 5, p = 0$; (J1–J5):

$\text{Cl}(X) = \mathbb{Z}$ is freely generated by the hyperplane section, with trivial G -action, and $\text{Pic}(\tilde{X})$ is freely generated by the exceptional divisors of the blowup $\tilde{X} \rightarrow X$ and the pullback to \tilde{X} of the basis of $\text{Cl}(X)$. The G -action permutes the exceptional divisors. So $\text{Pic}(\tilde{X})$ is a permutation module.

- $s = 4, 5, 6, p = 1$; (J6–J8):
 $\text{Cl}(X) = \mathbb{Z}^2$ is freely generated by the hyperplane section and the unique, necessarily G -stable, plane in X , with trivial G -action on their classes. Thus, $\text{Pic}(\tilde{X})$ is a permutation module.
- $s = 7, p = 2$; (J10):
 $\text{Cl}(X) = \mathbb{Z}^3$ is freely generated by the hyperplane section and the classes of the two planes in X , with G possibly permuting these two classes. Thus, $\text{Pic}(\tilde{X})$ is a permutation module.
- $s = 6, 7, p = 3$; (J11–J12):
 $\text{Cl}(X) = \mathbb{Z}^3$ is freely generated by the classes of the three planes in X which form a tetrahedron (with one face missing), with G possibly permuting these classes. So $\text{Pic}(\tilde{X})$ is a permutation module. □

The remaining cases are more involved; we handle these in subsequent sections.

Intermediate Jacobians

Applications of intermediate Jacobians to rationality problems (over $k = \mathbb{C}$) go back to the seminal work of Clemens–Griffiths [27]: if a smooth threefold X is rational, then its intermediate Jacobian $\text{IJ}(X)$ is a product of Jacobians of curves. Refinements of this, taking into account group actions, have appeared in, for example, [8]; an arithmetic analog of these arguments has been developed in [10]. In particular, intermediate Jacobians exist over arbitrary fields; see [1, 11].

The key point is that, geometrically, $\text{IJ}(X)$ could be a product of Jacobians of curves, but this does not necessarily hold equivariantly, respectively, over a nonclosed base field. This idea is implemented in, for example, [10, Theorem 1.1]. Pursuing the analogy, we have the following:

Theorem 2.3. *Let X be a smooth projective rationally connected threefold over an algebraically closed field such that its intermediate Jacobian $\text{IJ}(X)$ is isomorphic to the Jacobian of a smooth nonhyperelliptic curve C of genus $g \geq 3$, as a principally polarized abelian variety. Suppose that $\text{Aut}(X)$ contains an involution τ acting on $\text{IJ}(X)$ by multiplication by (-1) . Then X is not $\langle \tau \rangle$ -equivariantly birational to any smooth projective variety with trivial intermediate Jacobian.*

Proof. Suppose first that there exists a $\langle \tau \rangle$ -equivariant blowup $\pi: X \rightarrow Y$ of a nonhyperelliptic curve $C \subset Y$, where Y is a smooth threefold with trivial intermediate Jacobian. Since C is not hyperelliptic, it follows from Theorem 3 in [49, Appendix] that

$$\text{Aut}(\text{IJ}(X)) \simeq \text{Aut}(C) \times C_2, \tag{1}$$

where the second factor corresponds to the action of multiplication by (-1) . If C is pointwise fixed by τ , then τ acts trivially on $\text{IJ}(X)$. If τ acts faithfully on C , then its action on $\text{IJ}(X) = \text{J}(C)$ is induced by the action on C ; thus, τ cannot project nontrivially to the second factor in (1).

The general case is treated similarly, using equivariant weak factorization. □

Example 2.4. Consider the conic bundle

$$x_1x_2 = f(y_1, y_2, y_3) \subset \mathbb{A}^2 \times \mathbb{P}^2,$$

where f is a form of degree ≥ 4 defining a smooth curve in \mathbb{P}^2 , and C_2 -action via permutation on x_1, x_2 . Then this action is not linearizable by Theorem 2.3; see the proof of Theorem 3.3.

Remark 2.5. In the assumptions and notation of Theorem 2.3, suppose that there exists a G -equivariant birational map $X \dashrightarrow \mathbb{P}^3$, for some subgroup $G \subseteq \text{Aut}(X)$. From the isomorphism (1), we deduce that the G -action on $\text{IJ}(X)$ gives rise to a homomorphism

$$\nu: G \rightarrow \text{Aut}(\text{IJ}(X)) = \text{Aut}(C) \times C_2.$$

The projection of $\nu(G)$ to the C_2 -factor must be trivial; cf. the proof of [10, Proposition 3.2].

Burnside obstructions

It is well known that the classification of involutions $\tau \in \text{Cr}_2$, the plane Cremona group, is based on the geometry of τ -fixed loci $F(\tau)$; see, for example, [7]. The different cases are characterized by geometric properties of a (necessarily unique) curve C of genus ≥ 1 in $F(\tau)$, primarily by whether or not this curve is hyperelliptic. A more refined birational invariant of actions of general cyclic groups on rational surfaces, the *normalized fixed curve with action*, appeared in [34] and [12]; the invariant takes into account the stabilizer of the fixed curve, as well as the residual action on it.

These invariants are special cases of the Burnside formalism of [46], which applies to actions of arbitrary finite groups and takes into account *all* strata with nontrivial generic stabilizers. We will use a simplified version, explained in [24, Section 4]. It is based on the notion of *incompressible divisorial symbols*, which should be viewed as analogs of higher-genus curves in the classification of involutions on rational surfaces. A sample result in our context is the following:

Proposition 2.6. *Let X be a nodal cubic threefold, with a regular action of G , and assume that there is an element $\tau \in G$ such that*

- (1) *the τ -fixed locus contains a cubic surface $S \subset X$,*
- (2) *the subgroup $Y \subseteq G$ preserving S acts generically nontrivially on S and contains an element fixing a curve of genus ≥ 1 .*

Then the G -action on X is not linearizable.

Proof. Let $H = \langle \tau \rangle$; the action produces the symbol

$$(H, Y/H \curvearrowright k(S), (b)).$$

By [14], $H^1(Y/H, \text{Pic}(S)) \neq 0$, which implies that the symbol is *incompressible*; see [24, Section 4]. Such symbols cannot appear for linear actions; see [53, Corollary 6.1]. □

Example 2.7. Let $X \subset \mathbb{P}^4$ be a 2-nodal cubic given by

$$x_1x_2x_3 + x_1(x_4^2 + x_5^2) + x_2(x_4^2 - x_5^2) + x_3^3 = 0,$$

with $G \simeq C_4$ -action generated by

$$\iota : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_1, x_3, x_4, \zeta_4x_5).$$

The model satisfies the conditions in Proposition 2.6. In particular, the subgroup $\langle \iota^2 \rangle$ fixes the cubic surface $S = X \cap \{x_5 = 0\}$. The residual C_2 -action fixes a genus 1 curve $S \cap \{x_1 = x_2\}$. By Proposition 2.6, the G -action on X is not linearizable.

Specialization of birational types

We will use the specialization homomorphism for Burnside groups

$$\rho_{\pi}^G : \text{Burn}_{n,K}(G) \rightarrow \text{Burn}_{n,k}(G)$$

from [46, Definition 6.4], and in particular, [46, Corollary 6.8]. Here, K is the fraction field of a DVR, and k its residue field. In applications, one considers the local geometry of fibrations, seeking to specialize the birational type of the generic fiber X to a special fiber X_0 . In practice, the special fiber X_0 is an irreducible variety, with *mild* singularities; the relevant notion of BG -rational singularities on the special fiber X_0 is in [46, Definition 6.9].

Example 2.8. Let $\mathcal{X} \rightarrow \mathcal{B}$ be a G -equivariant flat and projective morphism onto a smooth curve \mathcal{B} , with smooth generic fiber X and a special fiber X_0 with ordinary double points. Then the singularities of X_0 are BG -rational singularities in the following situations:

- G -orbits of isolated ordinary double points, with trivial stabilizers [46, Example 6.10];
- $G = C_2$, fixing a singular point; one verifies directly that the required condition for BG -rational singularities holds – namely,

$$\rho_\pi^G([X \curvearrowright G]) = [X_0 \curvearrowright G].$$

A Hilbert scheme argument, used in [56], [28] and [40, Theorem 9] in the context of specialization of rationality properties, implies the following:

Proposition 2.9. *Let k be an uncountable algebraically closed field of characteristic zero and G a finite group. Let*

$$\pi : \mathcal{X} \rightarrow B$$

be a G -equivariant flat and projective morphism onto a smooth curve over k with smooth generic fiber, such that

- G acts trivially on B and generically freely on the fibers of π ,
- for some $b_0 \in B$, the special fiber \mathcal{X}_{b_0} is irreducible and has BG -rational singularities, and the G -action on \mathcal{X}_{b_0} is not linearizable.

Then, for very general $b \in B$, the G -action on the special fiber \mathcal{X}_b is not linearizable.

Specialization allows to exhibit nonlinearizable actions which are ‘invisible’ to classical obstructions (i.e., cannot be distinguished from linearizable actions with other available tools). However, the *very general* condition makes it difficult to determine linearizability for any specific variety in the family. A central problem is to give *criteria* for linearization.

In our applications, we work with models with nodes in the generic fiber. We reduce to the situation of Proposition 2.9 by equivariantly resolving the nodes in the generic fiber.

Example 2.10. Let $\mathcal{X} \rightarrow \mathbb{A}_k^1$ be a family of cubic threefolds with fibers \mathcal{X}_a over $a \in k$ given in \mathbb{P}^4 by

$$\begin{aligned} &a(x_1x_2^2 - 4x_3^2x_4 + x_3x_4^2 - 3x_3^2x_5 - x_4^2x_5) \\ &\quad + (a + 1)(x_1x_2x_3 + x_1x_2x_4 + x_1x_2x_5 - x_1x_4x_5 - x_2x_4x_5) \\ &\quad + x_1x_3x_4 - 3x_2x_3x_4 - 3x_1x_3x_5 + x_2x_3x_5 + (5a + 3)x_3x_4x_5 = 0. \end{aligned}$$

One can check that the family carries a $G = \langle \iota \rangle \simeq C_2$ -action, with ι acting on \mathbb{P}^4 via

$$(x_1, \dots, x_5) \mapsto (-x_3 + x_5, -x_3 + x_4, -x_3, -x_3 + x_2, -x_3 + x_1).$$

For a very general $a \in k$, \mathcal{X}_a is a 2-nodal cubic threefold with nodes at

$$p_1 = [1 : 0 : 0 : 0 : 0] \quad \text{and} \quad p_2 = [0 : 0 : 0 : 0 : 1].$$

But the special fiber over $a = 0$ is 6-nodal; the nodes are in general linear position, and ι does not fix any of the nodes. By Proposition 7.5, the G -action on the special fiber \mathcal{X}_0 is not stably linearizable. The four additional nodes form two G -orbits with trivial stabilizer, so they are BG -rational singularities, by Example 2.8. Blowing up the singularities in the generic fiber, we are in the situation of Proposition 2.9. This allows us to conclude that the G -action on a very general member of the family \mathcal{X} is not stably linearizable.

Birational rigidity

Let $X \subset \mathbb{P}^4$ be a nodal cubic threefold and let $G \subseteq \text{Aut}(X)$. If $\text{rk Cl}^G(X) = 1$, then X is a G -Mori fiber space [23, Definition 1.1.5], and every G -birational map from X to another G -Mori fiber space can be

decomposed into a sequence of elementary links, known as G -Sarkisov links [31, 39]. If there are no G -Sarkisov links that start at X , we say that X is G -birationally super-rigid. Similarly, if every G -Sarkisov link that starts at X also ends at X , we say that X is G -birationally rigid. Finally, if X is not G -birational to any G -Mori fiber space with a positive dimensional base (a conic bundle or a Del Pezzo fibration), we say that X is G -solid. We have the following implications:

$$G\text{-birationally super-rigid} \Rightarrow G\text{-birationally rigid} \Rightarrow G\text{-solid.}$$

Note that all of these conditions assume that $\text{rk Cl}^G(X) = 1$.

Recall that the G -action on X lifts to \mathbb{P}^4 . Using the G -action on \mathbb{P}^4 , we can state an obstruction for a cubic threefold X to be G -solid:

Lemma 2.11 [5, Lemma 2.6]. *If G leaves invariant a line or a plane in \mathbb{P}^4 , then X is not G -solid.*

Proof. Note that G leaves invariant a plane in \mathbb{P}^4 if and only if it leaves invariant a line. Thus, we may assume that there exists a G -stable plane in \mathbb{P}^4 . Linear projection $\mathbb{P}^4 \dashrightarrow \mathbb{P}^1$ from this plane induces a rational dominant map $X \dashrightarrow \mathbb{P}^1$ whose general fiber is a (possibly singular) cubic surface. Taking a G -equivariant resolution of indeterminacies of this map, a G -invariant resolution of singularities (if necessarily), and applying the G -equivariant Minimal Model program over \mathbb{P}^1 , we obtain a G -birational map from X to a G -Mori fiber space with a positive-dimensional base. \square

This yields the following result:

Theorem 2.12 (Avilov). *Let $X \subset \mathbb{P}^4$ be a nodal cubic threefold and $G \subseteq \text{Aut}(X)$ a finite subgroup such that $\text{rk Cl}^G(X) = 1$. If X is G -solid, then one of the following holds:*

- (1) $|\text{Sing}(X)| = 10$, X is the Segre cubic, $\text{Aut}(X) \simeq \mathfrak{S}_6$, and G contains a subgroup isomorphic to \mathfrak{A}_5 that leaves invariant a hyperplane section of X ,
- (2) $|\text{Sing}(X)| = 9$, X is given in \mathbb{P}^5 by

$$x_1x_2x_3 - x_4x_5x_6 = \sum_{i=1}^6 x_i = 0,$$

$\text{Aut}(X) \simeq \mathfrak{S}_3 \rtimes C_2$, G acts transitively on $\text{Sing}(X)$ and is isomorphic to $\mathfrak{S}_3 \rtimes C_2$, \mathfrak{S}_3^2 or $C_3^2 \rtimes C_4$,


- (3) $|\text{Sing}(X)| = 5$, $X \subset \mathbb{P}^4$ is given by

$$x_1x_2x_3 + x_1x_2x_4 + x_1x_2x_5 + x_1x_3x_4 + x_1x_3x_5 + x_1x_4x_5 + x_2x_3x_4 + x_2x_3x_5 + x_2x_4x_5 + x_3x_4x_5 = 0,$$

$\text{Aut}(X) \simeq \mathfrak{S}_5$, and either $G \simeq \mathfrak{S}_5$ or $G \simeq \mathfrak{A}_5$.

Proof. Suppose that X is G -solid. If there exists a G -equivariant birational map $X \dashrightarrow \mathbb{P}^3$, then \mathbb{P}^3 is G -solid, which contradicts [20]. Thus, the G -action on X is not linearizable. It follows from [6, 4, 5] and the proofs of the main results in these papers that either X and G are as in (1), (2), (3), or X is the cubic threefold in (3) and $G \simeq C_4 \rtimes C_5$. Let us show that X is not G -solid in the latter case, contradicting the assumption.

Namely, suppose X is the threefold from (3), and $G \simeq C_4 \rtimes C_5$. By [5, 21], there exists the following \mathfrak{S}_5 -Sarkisov link:



where Q is the smooth quadric threefold

$$\{x_1x_2 + x_2x_3 + \cdots + x_4x_5 = 0\} \subset \mathbb{P}^4,$$

\mathfrak{S}_5 acts on Q by permuting the coordinates, χ is the birational map induced by the standard Cremona involution of \mathbb{P}^4 , α is the standard resolution of singularities, β is a composition of 10 Atiyah flops, and γ is a blowup of an \mathfrak{S}_5 -orbit of length 5. Let $\eta: Q \rightarrow \mathbb{P}^3$ be the double cover induced by the projection from the \mathfrak{S}_5 -fixed point in \mathbb{P}^4 . Then η is \mathfrak{S}_5 -equivariant, and \mathbb{P}^3 contains two skew lines L_1 and L_2 such that the curve $L_1 + L_2$ is G -invariant. Let C_1 and C_2 be the preimages of these lines on Q . Then C_1 and C_2 are disjoint conics, and the curve $C_1 + C_2$ is G -invariant. Blowing up these two conics, we obtain a G -equivariant birational map from Q to a conic bundle over $\mathbb{P}^1 \times \mathbb{P}^1$; in particular, X is not G -solid, which contradicts our assumption. In fact, the G -action on Q is linearizable; see Section 6. \square

Moreover, in Case (1) in Theorem 2.12, X is G -birationally super-rigid [6]. Similarly, it follows from [21] that X is G -solid in Case (3) when $G \simeq \mathfrak{S}_5$. In Section 9, we show that X is G -birationally super-rigid in Case (2) when $G \simeq \mathfrak{S}_3^2 \rtimes C_2$. We believe that X is G -solid for the remaining groups G in Cases (2) and (3).

3. Two nodes

Standard form

We may assume that $\text{Sing}(X)$ consists of the points

$$[1 : 0 : 0 : 0 : 0], \quad [0 : 1 : 0 : 0 : 0],$$

and that $G = \text{Aut}(X)$ swaps these points. Then X can be given by

$$x_1x_2x_3 + x_1q_1 + x_2q_2 + f_3 = 0, \tag{2}$$

for forms $q_1, q_2 \in k[x_4, x_5]$, and $f_3 \in k[x_3, x_4, x_5]$, of degree 2, 2, 3, respectively.

Conic bundle

Introducing new coordinates $y_1 = x_1x_3$ and $y_2 = x_2x_3$ (of weight two), and multiplying (2) by x_3 , we rewrite (2) as

$$y_1y_2 + y_1q_1 + y_2q_2 + x_3f_3 = 0,$$

which defines a quartic hypersurface $V_4 \subset \mathbb{P}(1, 1, 1, 2, 2)$; the coordinate change defines a G -equivariant birational map

$$\chi: X \dashrightarrow V_4.$$


We can G -equivariantly simplify the equation of V_4 further as

$$z_1z_2 = q_1q_2 - x_3f_3,$$

where $z_1 = y_1 + q_2$ and $z_2 = y_2 + q_1$. Observe that V_4 has two singular points of type $\frac{1}{2}(1, 1, 1)$ – these are the points

$$[0 : 0 : 0 : 1 : 0], \quad \text{and} \quad [0 : 0 : 0 : 0 : 1],$$

in coordinates $(x_3, x_4, x_5, z_1, z_2)$. This yields the following $\text{Aut}(X)$ -equivariant commutative diagram:



where α is an extremal divisorial contraction of a surface to the line $\{x_3 = x_4 = x_5 = 0\} \subset X$, and β is an extremal divisorial contraction of the strict transform of the non-normal cubic surface $\{x_3 = 0\} \cap X$. The description of the morphism α can be found in the proof of [18, Proposition 6.1]; see also [54]. Note that \widehat{X} has two singular points of type $\frac{1}{2}(1, 1, 1)$, which are mapped to the nodes of X .

Let D be the quartic curve $\{q_1q_2 - x_3f_3 = 0\} \subset \mathbb{P}^2_{x_3, x_4, x_5}$. Then D is smooth, and we have the following G -equivariant commutative diagram



where γ is the blowup of the singular points of V_4 , π is a conic bundle with discriminant curve D , and the dashed arrow is the projection

$$(x_3, x_4, x_5, z_1, z_2) \mapsto (x_3, x_4, x_5).$$

This gives a natural homomorphism

$$\gamma : \text{Aut}(X) \rightarrow \text{Aut}(D).$$

Automorphisms

The full classification of automorphisms of 2-nodal cubics can be addressed via the conic bundle presentation, combined with the (classically known) classification of automorphisms of smooth quartic plane curves (see, for example, [36, Lemma 6.16 and Table 6], [47]); and using Torelli for nodal cubics, as in [15, Section 7]. Starting with equation (2) and passing to the conic bundle, we see that the G -action on X gives rise to

- a linear representation on \mathbb{P}^2 , preserving a line, corresponding to $x_3 = 0$, and thus a fixed point in \mathbb{P}^2 ,
- an automorphism of the discriminant curve D .

Combining these two conditions with the list of automorphisms of plane quartic curves, we find that the possible images of the G -actions on the base $\mathbb{P}^2_{x_3, x_4, x_5}$ of the conic bundle are

$$\begin{aligned}
 &C_2, C_3, C_4, C_2^2, \mathfrak{S}_3, C_6, C_7, C_2 \times C_4, C_8, Q_8, C_9, C_4^2, \\
 &C_{12}, D_4 \rtimes C_2, \text{SL}_2(\mathbb{F}_3), OD16, \mathfrak{D}_4, C_4 \text{ wr } C_2, \text{SL}_2(\mathbb{F}_3) \rtimes C_2.
 \end{aligned}$$

Here are examples with interesting groups $\text{Aut}(X)$:

Example 3.1. We keep the notation of (2), with $X \subset \mathbb{P}^4$ and the discriminant curve $D \subset \mathbb{P}^2_{x_3, x_4, x_5}$, with $G = \text{Aut}(X)$ and $G' = \text{Aut}(D)$.

(1) Let $D = x_3^4 + x_4^4 + x_5^4 + (4\rho + 2)x_3^2x_4^2$, and X be given by

$$f_3 = x_3^3, \quad q_1 = x_4^2 + (2\rho + 1 + 2i)x_5^2, \quad q_2 = x_4^2 + (2\rho + 1 - 2i)x_5^2.$$

Then $G = \mathfrak{D}_4 \rtimes C_2$ and $G' = \mathrm{SL}_2(\mathbb{F}_3) \rtimes C_2$.

(2) Let $D = x_3^4 + x_4^4 + x_5^4$, and X be given by

$$f_3 = -x_3^3, \quad q_1 = x_4^2 + ix_5^2, \quad q_2 = x_4^2 - ix_5^2.$$

Then $G = C_4 \wr C_2$ and $G' = C_4^2 \rtimes \mathfrak{S}_3$.

Proposition 3.2. Let $X \subset \mathbb{P}^4$ be a 2-nodal cubic threefold with an action of a cyclic group $G = \langle \iota \rangle \subseteq \mathrm{Aut}(X)$ not fixing any node. Then, up to a change of coordinates, X is given by

$$x_1x_2x_3 + x_1q_1 + x_2q_2 + f_3 = 0,$$

for $q_1, q_2 \in k[x_4, x_5]$ and $f_3 \in k[x_3, x_4, x_5]$ that can be described together with ι as follows.

(C₂) $\iota(x_1, x_2, x_3, x_4, x_5) = (x_2, x_1, x_3, x_4, x_5),$

$$q_1 = q_2 = x_4x_5, \\ f_3 \in k[x_3, x_4, x_5];$$

(C'₂) $\iota(x_1, x_2, x_3, x_4, x_5) = (-x_2, -x_1, x_3, x_4, -x_5),$

$$q_1 = a_1x_4^2 + x_4x_5 + a_3x_5^2, \\ q_2 = -a_1x_4^2 + x_4x_5 - a_3x_5^2, \\ f_3 = c_1x_3^3 + d_1x_3^2x_4 + x_3(e_1x_4^2 + e_3x_5^2) + r_1x_4^3 + r_3x_4x_5^2,$$

for some $a_1, a_3, c_1, d_1, e_1, e_3, r_1, r_3 \in k$;

(C''₂) $\iota(x_1, x_2, x_3, x_4, x_5) = (x_2, x_1, x_3, x_4, -x_5),$

$$q_1 = q_2 = x_4^2 + x_5^2 \\ f_3 = c_1x_3^3 + d_1x_3^2x_4 + x_3(e_1x_4^2 + e_3x_5^2) + r_1x_4^3 + r_3x_4x_5^2,$$

for some $c_1, d_1, e_1, e_3, r_1, r_3 \in k$;

(C₄) $\iota(x_1, x_2, x_3, x_4, x_5) = (x_2, x_1, x_3, \zeta_4x_4, -\zeta_4x_5), \quad \zeta_4 = e^{\frac{2\pi i}{4}},$

$$q_1 = a_1x_4^2 + x_4x_5 + a_3x_5^2, \\ q_2 = -a_1x_4^2 + x_4x_5 - a_3x_5^2, \\ f_3 = x_3^3 + e_2x_3x_4x_5,$$

for some $a_1, a_2, e_2 \in k$;

(C'₄) $\iota(x_1, x_2, x_3, x_4, x_5) = (x_2, x_1, x_3, x_4, \zeta_4x_5),$

$$q_1 = x_4^2 - x_5^2, \\ q_2 = x_4^2 + x_5^2, \\ f_3 \in k[x_3, x_4];$$

$$(C_4'') \iota(x_1, x_2, x_3, x_4, x_5) = (x_2, x_1, x_3, -x_4, \zeta_4 x_5),$$

$$\begin{aligned} q_1 &= x_4^2 - x_5^2, \\ q_2 &= x_4^2 + x_5^2, \\ f_3 &= x_3^3 + e_1 x_3 x_4^2 + r_3 x_4 x_5^2, \end{aligned}$$

for some $e_1, r_3 \in k$;

$$(C_6) \iota(x_1, x_2, x_3, x_4, x_5) = (x_2, x_1, x_3, \zeta_3 x_4, \zeta_3^2 x_5) \text{ for } \zeta_3 = e^{\frac{2\pi i}{3}},$$

$$\begin{aligned} q_1 &= q_2 = x_4 x_5, \\ f_3 &= x_3^3 + e_2 x_3 x_4 x_5 + r_1 (x_4^3 + x_5^3), \end{aligned}$$

for some $e_2, r_1 \in k$;

$$(C_6') \iota(x_1, x_2, x_3, x_4, x_5) = (\zeta_6 x_2, \zeta_6 x_1, x_3, \zeta_6^5 x_4, \zeta_6^2 x_5) \text{ for } \zeta_6 = e^{\frac{2\pi i}{6}},$$

$$\begin{aligned} q_1 &= a_1 x_4^2 + x_4 x_5 + a_3 x_5^2, \\ q_2 &= -a_1 x_4^2 + x_4 x_5 - a_3 x_5^2, \\ f_3 &= x_3^2 x_5, \end{aligned}$$

for some $a_1, a_3 \in k$;

$$(C_{12}) \iota(x_1, x_2, x_3, x_4, x_5) = (\zeta_{12}^8 x_2, \zeta_{12}^8 x_1, x_3, \zeta_{12}^4 x_4, \zeta_{12} x_5) \text{ for } \zeta_{12} = e^{\frac{2\pi i}{12}},$$

$$\begin{aligned} q_1 &= (x_4^2 - x_5^2), \\ q_2 &= (x_4^2 + x_5^2), \\ f_3 &= x_3^2 x_4. \end{aligned}$$

Proof. We can choose the coordinates so that

$$\iota : (x_1, x_2, x_3, x_4, x_5) \mapsto (sx_2, tx_1, x_3, ux_4, vx_5),$$

for some $s, t, u, v \in k^\times$, and X is given by

$$x_1 x_2 x_3 + x_1 q_1 + x_2 q_2 + f_3 = 0,$$

for

$$\begin{aligned} q_1 &= a_1 x_4^2 + a_2 x_4 x_5 + a_3 x_5^2 + a_4 x_3^2 + a_5 x_3 x_4 + a_6 x_3 x_5, \\ q_2 &= b_1 x_4^2 + b_2 x_4 x_5 + b_3 x_5^2 + b_4 x_3^2 + b_5 x_3 x_4 + b_6 x_3 x_5, \\ f_3 &= c_1 x_3^3 + x_3^2 d(x_4, x_5) + x_3 e(x_4, x_5) + r(x_4, x_5), \end{aligned}$$

where

$$\begin{aligned} d &= d_1 x_4 + d_2 x_5, \\ e &= e_1 x_4^2 + e_2 x_4 x_5 + e_3 x_5^2, \\ r &= r_1 x_4^3 + r_2 x_4^2 x_5 + r_3 x_4 x_5^2 + r_4 x_5^3. \end{aligned}$$

Since X is $\langle \iota \rangle$ -invariant, one has $\iota^*(f) = stf$, and thus, the zero loci of q_1q_2 and f_3 are preserved; and these polynomials cannot identically vanish, under our assumptions on singularities of X . Concretely,

$$\iota^*(f) = stx_1x_2x_3 + sx_2\iota^*(q_2) + tx_1\iota^*(q_2) + \iota^*(f_3) = stf,$$

which implies that

$$\iota^*(q_2) = tq_1, \quad \iota^*(q_1) = sq_2, \tag{4}$$

and

$$\iota^*(f_3) = stf_3. \tag{5}$$

Expanding and substituting into (4), we obtain 12 equations:

$$\begin{aligned} u^2a_1 - tb_1 &= uva_2 - tb_2 = v^2a_3 - tb_3 = 0, \\ sa : 1 - u^2b_1 &= sa : 2 - uvb_2 = sa : 3 - v^2b_3 = 0, \\ a_4 - tb_4 &= ua_5 - tb_5 = va_6 - tb_6 = 0, \\ sa : 4 - b_4 &= sa : 5 - ub_5 = sa : 6 - vb_6 = 0, \end{aligned}$$

and, writing down the (5) constraints on f_3 , additional equations

$$\begin{aligned} c_1(1 - st) &= 0, \\ d_1(u - st) &= d_2(v - st) = 0, \\ e_1(u^2 - st) &= e_2(uv - st) = e_3(v^2 - st) = 0, \\ r_1(u^3 - st) &= r_2(uv - st) = r_3(uv^2 - st) = r_4(v^3 - st) = 0, \end{aligned}$$

in the variables

$$a_1, \dots, a_6, b_1, \dots, b_6, c_1, d_1, d_2, e_1, e_2, r_1, \dots, r_4 \in k.$$

Since X has nodes at $[1 : 0 : 0 : 0 : 0]$ and $[0 : 1 : 0 : 0 : 0]$, we have

$$a_2^2 - 4a_1a_3 \neq 0, \quad b_2^2 - 4b_1b_3 \neq 0.$$

Thus, up to scaling x_4, x_5 and swapping them, we may further assume that one of the following holds:

- $a_2 = b_2 = 1$, or
- $a_2 = b_1 = b_3 = 1, b_2 = 0$, or
- $a_1 = b_1 = b_3 = 1, a_2 = b_2 = 0$.

The second option is impossible since $b_2 = 0$ forces $a_2 = 0$. Solving the system of equations for the remaining two options using Magma, we obtain a complete set of solutions:

	$(a_1, a_2, a_3, a_4, a_5, a_6)$ $(b_1, b_2, b_3, b_4, b_5, b_6)$	(d_1, d_2) (e_1, e_2, e_3)	c_1 (r_1, r_2, r_3, r_4)	s t	u v
(1)	$(a_1, 1, a_3, a_4, a_5, a_6)$ $(a_1, 1, a_3, a_4, a_5, a_6)$			1 1	1 1
(2)	$(a_1, 1, a_3, a_4, a_5, a_6)$ $(-a_1, 1, -a_3, -a_4, -a_5, a_6)$	$(d_1, 0)$ $(e_1, 0, e_3)$	$(r_1, 0, r_3, 0)$	-1 -1	1 -1
(3)	$(0, 1, 0, a_4, 0, 0)$ $(0, 1, 0, a_4, 0, 0)$	$(0, 0)$ $(0, e_2, 0)$	$(r_1, 0, 0, r_4)$	1 1	ζ_3 ζ_3^2
(4)	$(a_1, 1, a_3, a_4, 0, 0)$ $(-a_1, 1, -a_3, a_4, 0, 0)$	$(0, 0)$ $(0, e_2, 0)$	$(0, 0, 0, 0)$	1 1	ζ_4 $-\zeta_4$
(5)	$(a_1, 1, a_3, 0, 0, 0)$ $(-a_1, 1, -a_3, 0, 0, 0)$	$(0, d_2)$ $(0, 0, 0)$	0 $(0, 0, 0, 0)$	ζ_6 ζ_6	ζ_6^5 ζ_6^2
(6)	$(1, 0, 1, a_4, a_5, a_6)$ $(1, 0, 1, a_4, a_5, a_6)$			1 1	1 1
(7)	$(1, 0, 1, a_4, a_5, a_6)$ $(1, 0, 1, a_4, a_5, -a_6)$	$(d_1, 0)$ $(e_1, 0, e_3)$	$(r_1, 0, r_3, 0)$	1 1	1 -1
(8)	$(1, 0, -1, a_4, a_5, 0)$ $(1, 0, 1, a_4, a_5, 0)$	$(d_1, 0)$ $(e_1, 0, 0)$	$(r_1, 0, 0, 0)$	1 1	1 ζ_4
(9)	$(1, 0, -1, a_4, a_5, 0)$ $(1, 0, 1, a_4, -a_5, 0)$	$(0, 0)$ $(e_1, 0, 0)$	$(0, 0, r_3, 0)$	1 1	-1 ζ_4
(10)	$(1, 0, -1, 0, 0, 0)$ $(1, 0, 1, 0, 0, 0)$	$(d_1, 0)$ $(0, 0, 0)$	0 $(0, 0, 0, 0)$	ζ_{12}^8 ζ_{12}^8	ζ_{12}^4 ζ_{12}^4

Here, we omitted solutions obtained by swapping coordinates x_4 and x_5 and scaling coordinates x_1 and x_2 . After an additional equivariant change of coordinates, we obtain the required assertion. \square

Intermediate Jacobian

Using arguments as in Section 2, we settle the linearizability problem for 2-nodal cubic threefolds.

Theorem 3.3. *Let $X \subset \mathbb{P}^4$ be a 2-nodal cubic, and $G \subseteq \text{Aut}(X)$ a subgroup not fixing any node of X . Then the G -action on X is not linearizable.*

Proof. By the assumptions, G contains an element ι switching the nodes of X . It suffices to prove the required assertion for $G = \langle \iota \rangle$. With the notation as above, we may assume that X is given by (2); that is,

$$f = x_1x_2x_3 + x_1q_2 + x_2q_2 + f_3 = 0,$$

and ι acts on the coordinates via

$$\iota : (x_1, x_2, x_3, x_4, x_5) \mapsto (sx_2, tx_1, x_3, ux_4, vx_5),$$

for roots of unity s, t, u, v . Introducing new coordinates

$$w_1 = \sqrt{t}x_1 + \sqrt{s}x_2, \quad w_2 = \sqrt{t}x_1 - \sqrt{s}x_2,$$

we diagonalize ι , so that it acts via

$$\iota : (w_1, w_2, x_3, x_4, x_5) \mapsto (\lambda w_1, -\lambda w_2, x_3, ux_4, vx_5), \tag{6}$$

where $\lambda = \sqrt{st}$. The equation of X in the new coordinates becomes

$$f' = (w_1^2 - w_2^2)x_3 + (w_1 + w_2)2\sqrt{s}q_1 + (w_1 - w_2)2\sqrt{t}q_2 + 4\sqrt{st}f_3 = 0.$$

Note that $\iota^*(f') = \lambda^2 f'$.

Recall that X is G -birational to the conic bundle (3). The conic bundle is not standard. In particular, the intermediate Jacobian of \tilde{X} and the Jacobian of the curve D are isomorphic, as principally polarized abelian varieties:

$$\text{IJ}(\tilde{X}) \simeq \text{IJ}(Y) \simeq \text{J}(D),$$

where \tilde{X} is the standard desingularization of X , Y is the conic bundle in (3) and D is its discriminant curve in $\mathbb{P}^2_{x_3, x_4, x_5}$ given by the quartic form $h = q_1q_2 - x_3f_3$. Note that $\iota^*(h) = \lambda^2h$.

Arguing as in the proof of [9, Lemma 1], we see that ι acts faithfully on $\text{IJ}(\tilde{X})$ and preserves the principal polarization. However, the ι -action on coordinates x_3, x_4, x_5 induces an action on D and its Jacobian $\text{J}(D)$. We claim that the ι -actions on $\text{IJ}(\tilde{X})$ and $\text{J}(D)$ differ by multiplication by -1 . Since D is not hyperelliptic, this would imply that the G -action on X is not linearizable, by Remark 2.5.

To compute the action of ι on $\text{IJ}(\tilde{X})$, recall that its tangent space at zero $\mathcal{T}_0 \text{IJ}(\tilde{X})$ is isomorphic to $\text{H}^2(\tilde{X}, \Omega^1_{\tilde{X}})$. We show that $\text{H}^2(\tilde{X}, \Omega^1_{\tilde{X}})^\vee$ is canonically isomorphic to the linear subspace in

$$\text{H}^0(\mathbb{P}^4, \Omega^4_{\mathbb{P}^4} \otimes \mathcal{O}_{\mathbb{P}^4}(2X)) \cong \text{H}^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(K_{\mathbb{P}^4} + 2X)) \cong \text{H}^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(1))$$

consisting of all sections that vanish at the nodes of X . The proof is essentially contained in [33]. We follow the proof of [9, Lemma 1]. Let $\pi : \tilde{\mathbb{P}}^4 \rightarrow \mathbb{P}^4$ be the blowup of \mathbb{P}^4 centered at two nodes p_1 and p_2 of X , and identify \tilde{X} with the strict transform of X in $\tilde{\mathbb{P}}^4$. The exact sequence

$$0 \rightarrow (\mathcal{N}_{\tilde{X}/\tilde{\mathbb{P}}^4})^\vee \rightarrow \Omega^1_{\tilde{\mathbb{P}}^4|\tilde{X}} \rightarrow \Omega^1_{\tilde{X}} \rightarrow 0$$

gives rise to a $\langle \iota \rangle$ -equivariant exact sequence

$$0 \rightarrow \text{H}^2(\tilde{X}, \Omega^1_{\tilde{X}}) \rightarrow \text{H}^3(\tilde{X}, (\mathcal{N}_{\tilde{X}/\tilde{\mathbb{P}}^4})^\vee) \rightarrow \text{H}^3(\tilde{X}, \Omega^1_{\tilde{\mathbb{P}}^4|\tilde{X}}) \rightarrow 0.$$

By [33], the dimension of $\text{H}^3(\tilde{X}, \Omega^1_{\tilde{\mathbb{P}}^4|\tilde{X}})$ equals the defect of X , which is 0 in our case. It follows that

$$\text{H}^2(\tilde{X}, \Omega^1_{\tilde{X}}) \cong \text{H}^3(\tilde{X}, (\mathcal{N}_{\tilde{X}/\tilde{\mathbb{P}}^4})^\vee).$$

Similarly, the $\langle \iota \rangle$ -equivariant exact sequence

$$0 \rightarrow \mathcal{O}_{\tilde{\mathbb{P}}^4}(-2\tilde{X}) \rightarrow \mathcal{O}_{\tilde{\mathbb{P}}^4}(-\tilde{X}) \rightarrow (\mathcal{N}_{\tilde{X}/\tilde{\mathbb{P}}^4})^\vee \rightarrow 0$$

and the vanishing of $\text{H}^i(\tilde{\mathbb{P}}^4, \mathcal{O}_{\tilde{\mathbb{P}}^4}(-\tilde{X}))$ ([33, Corollary 2]) provide an C_2 -isomorphism

$$\text{H}^3(\tilde{X}, (\mathcal{N}_{\tilde{X}/\tilde{\mathbb{P}}^4})^\vee) \cong \text{H}^4(\tilde{\mathbb{P}}^4, \mathcal{O}_{\tilde{\mathbb{P}}^4}(-2\tilde{X})).$$

By Serre duality, we have canonical isomorphisms between

$$\text{H}^4(\tilde{\mathbb{P}}^4, \mathcal{O}_{\tilde{\mathbb{P}}^4}(-2\tilde{X}))^\vee \cong \text{H}^0(\tilde{\mathbb{P}}^4, K_{\tilde{\mathbb{P}}^4} \otimes \mathcal{O}_{\tilde{\mathbb{P}}^4}(2\tilde{X})) \cong \text{H}^0(\tilde{\mathbb{P}}^4, \Omega^4_{\tilde{\mathbb{P}}^4} \otimes \mathcal{O}_{\tilde{\mathbb{P}}^4}(2\tilde{X})).$$

So we have a $\langle \iota \rangle$ -equivariant isomorphism

$$\text{H}^2(\tilde{X}, \Omega^1_{\tilde{X}})^\vee \cong \text{H}^0(\tilde{\mathbb{P}}^4, K_{\tilde{\mathbb{P}}^4} \otimes \mathcal{O}_{\tilde{\mathbb{P}}^4}(2\tilde{X})).$$

Let E_1 and E_2 be the exceptional divisors of π over p_1 and p_2 , respectively. By adjunction,

$$K_{\tilde{\mathbb{P}}^4} = \pi^*(\mathcal{O}_{\mathbb{P}^4}(-5)) \otimes \mathcal{O}_{\tilde{\mathbb{P}}^4}(3E_1 + 3E_2),$$

and

$$\mathcal{O}_{\mathbb{P}^4}(-2\tilde{X}) = \pi^*(\mathcal{O}_{\mathbb{P}^4}(6)) \otimes \mathcal{O}_{\mathbb{P}^4}(-4E_1 - 4E_2).$$

Then we know

$$K_{\mathbb{P}^4} \otimes \mathcal{O}_{\mathbb{P}^4}(-2\tilde{X}) = \pi^*(\mathcal{O}_{\mathbb{P}^4}(1)) \otimes \mathcal{O}_{\mathbb{P}^4}(-E_1 - E_2).$$

It follows that we can canonically identify $H^2(\tilde{X}, \Omega_{\tilde{X}}^1)^\vee$ with linear subspace in $H^0(\mathbb{P}^4, \Omega_{\mathbb{P}^4}^4 \otimes \mathcal{O}_{\mathbb{P}^4}(2X))$, (or equivalently, in $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(1))$), which consists of all sections that vanish at p_1 and p_2 . Now we can compute the induced G -action on $\mathcal{T}_0 \text{IJ}(\tilde{X})^\vee$ explicitly. Set

$$z_2 = \frac{w_2}{w_1}, \quad z_3 = \frac{x_3}{w_1}, \quad z_4 = \frac{x_4}{w_1}, \quad z_5 = \frac{x_5}{w_1},$$

and consider the rational 4-forms

$$z_3\omega, \quad z_4\omega, \quad z_5\omega, \quad \text{where } \omega = \frac{dz_2 \wedge dz_3 \wedge dz_4 \wedge dz_5}{(f'(1, z_2, z_3, z_4, z_5))^2}.$$

These give sections of $H^0(\mathbb{P}^4, \Omega_{\mathbb{P}^4}^4 \otimes \mathcal{O}_{\mathbb{P}^4}(2X))$, forming a basis of the subspace consisting of sections that vanish at the nodes of X . One computes

$$\iota^*(z_2) = -z_2, \quad \iota^*(z_3) = \frac{1}{\lambda}z_3, \quad \iota^*(z_4) = \frac{u}{\lambda}z_4, \quad \iota^*(z_5) = \frac{v}{\lambda}z_5$$

and

$$\iota^*(f'(1, z_2, \dots, z_5)^2) = \iota^*\left(\frac{f'(w_1, w_2, x_3, x_4, x_5)^2}{w_1^6}\right) = \frac{f'(1, z_2, \dots, z_5)^2}{\lambda^2}.$$

Using these, we see that ι acts on $\mathcal{T}_0 \text{IJ}(\tilde{X})^\vee$ with eigenvalues

$$-\frac{uv}{\lambda^2}, \quad -\frac{u^2v}{\lambda^2}, \quad -\frac{uv^2}{\lambda^2}.$$

Similarly, to compute the action of ι on $\text{J}(D)$, we note that $\mathcal{T}_0 \text{J}(D)^\vee$ is canonically isomorphic to

$$H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}^2 \otimes \mathcal{O}_{\mathbb{P}^2}(D)).$$

Set $y_4 = \frac{x_4}{x_3}$ and $y_5 = \frac{x_5}{x_3}$. The rational 2-forms

$$\frac{dy_4 \wedge dy_5}{h(1, y_4, y_5)}, \quad y_4 \frac{dy_4 \wedge dy_5}{h(1, y_4, y_5)}, \quad y_5 \frac{dy_4 \wedge dy_5}{h(1, y_4, y_5)}$$

define sections of $H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}^2 \otimes \mathcal{O}_{\mathbb{P}^2}(D))$, forming its basis. One has

$$\iota^*(y_4) = uy_4, \quad \iota^*(y_5) = vy_5, \quad \iota^*(h(1, y_4, y_5)) = \lambda^2 h(1, y_4, y_5),$$

and ι acts on $\mathcal{T}_0 \text{J}(D)^\vee$ with eigenvalues

$$\frac{uv}{\lambda^2}, \quad \frac{u^2v}{\lambda^2}, \quad \frac{uv^2}{\lambda^2}.$$

This shows that the ι -action on $\text{IJ}(\tilde{X}) \simeq J(D)$ differs from the action on $J(D)$ induced by the action on D by multiplication by -1 , as claimed. Therefore, the G -action on X is not linearizable. \square

4. Three nodes

Standard form

The three nodes are necessarily in general linear position; they span a distinguished G -stable plane, which is *not* contained in X . This case is labeled (J3), in [37]. Assume the nodes are

$$p_1 = [1 : 0 : 0 : 0 : 0], \quad p_2 = [0 : 1 : 0 : 0 : 0], \quad p_3 = [0 : 0 : 1 : 0 : 0].$$

The standard form is given by

$$x_1x_2x_3 + x_1q_1 + x_2q_2 + x_3q_3 + f_3 = 0, \tag{7}$$

with quadratic $q_1, q_2, q_3 \in k[x_4, x_5]$, and cubic $f_3 \in k[x_4, x_5]$. Note that q_1, q_2, q_3 must have rank 2, and q_1, q_2, q_3, f_3 do not share common factors.

Automorphisms

We proceed to classify automorphism groups of 3-nodal cubics acting transitively on nodes.

Proposition 4.1. *Let $X \subset \mathbb{P}^4$ be a 3-nodal cubic threefold. Assume that $\text{Aut}(X)$ contains an element acting transitively on the nodes. Then, up to a change of coordinates, X is given by*

$$x_1x_2x_3 + x_1q_1 + x_2q_2 + x_3q_3 + f_3 = 0,$$

for $q_1, q_2, q_3, f_3 \in k[x_4, x_5]$ that can be described together with $\text{Aut}(X)$ as follows.

1. $\text{Aut}(X) = C_3$, generated by

$$\sigma_1 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, \zeta_6^2 x_5), \quad \zeta_6 = e^{\frac{2\pi i}{6}},$$

◦ with $f_3 = ax_4^3 + bx_5^3$, for $b \neq 0$, $(a, b) \neq (0, 1)$ and

$$\begin{aligned} q_1 &= x_4(x_4 + x_5), \\ q_2 &= x_4(x_4 + \zeta_6^2 x_5), \\ q_3 &= x_4(x_4 + \zeta_6^4 x_5), \quad \text{or} \end{aligned}$$

◦ with $f_3 = dx_4^3 + ex_5^3$,

$$\begin{aligned} q_1 &= x_4^2 + bx_4x_5 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 bx_4x_5 + \zeta_6^4 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 bx_4x_5 + \zeta_6^2 x_5^2, \quad d \neq \pm e, \text{ and } b \neq \pm 2. \end{aligned}$$

2. $\text{Aut}(X) = C_6$ generated by

$$\sigma_2 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, \zeta_6 x_5),$$

$f_3 = dx_4^3$ for some $d \neq 0$, and

$$\begin{aligned} q_1 &= x_4^2 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 x_5^2. \end{aligned}$$

3. $\text{Aut}(X) \simeq \mathfrak{S}_3$ generated by σ_1 and

$$\sigma_3 : (x_1, x_2, x_3, x_4, x_5) \mapsto (\zeta_6^2 x_2, \zeta_6^4 x_1, x_3, \zeta_6^4 x_5, \zeta_6^2 x_4),$$

$$\begin{aligned} f_3 &= d(x_4^3 + x_5^3), \\ q_1 &= x_4^2 + bx_4x_5 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 bx_4x_5 + \zeta_6^4 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 bx_4x_5 + \zeta_6^2 x_5^2, \quad d \neq 0, b \neq \pm 2. \end{aligned}$$

4. $\text{Aut}(X) \simeq C_2 \times \mathfrak{S}_3$ generated by σ_1, σ_3 and

$$\iota : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, -x_4, -x_5),$$

with $f_3 = 0$ for $b \neq \pm 2$ and $b^2 \neq -2$, and

$$\begin{aligned} q_1 &= x_4^2 + bx_4x_5 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 bx_4x_5 + \zeta_6^4 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 bx_4x_5 + \zeta_6^2 x_5^2. \end{aligned}$$

5. $\text{Aut}(X) \simeq C_2 \times \mathfrak{S}_3$ generated by

$$\sigma_4 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, x_5),$$

$$\sigma_5 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_1, x_3, x_4, x_5),$$

$$\sigma_6 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, x_5, x_4),$$

with

$$f_3 = \begin{cases} d(x_4 + x_5)^3 & d \neq 0, \text{ or} \\ d(x_4 + x_5)(x_4 - x_5)^2 & d \neq 0, \text{ or} \\ (x_4 + x_5)(ax_4 + bx_5)(bx_4 + ax_5) & \text{for } a, b \neq 0, \end{cases}$$

$$q_1 = q_2 = q_3 = x_4x_5.$$

6. $\text{Aut}(X) \simeq C_3 \rtimes \mathfrak{D}_4$ generated by σ_2 and

$$\sigma_7 : (x_1, x_2, x_3, x_4, x_5) \mapsto (\zeta_6^4 x_2, \zeta_6^2 x_1, x_3, \zeta_6^2 x_5, \zeta_6 x_4),$$

with $f_3 = 0$ and

$$\begin{aligned} q_1 &= x_4^2 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 x_5^2. \end{aligned}$$

7. $\text{Aut}(X) \simeq \mathfrak{S}_3 \times \mathfrak{S}_3$ generated by $\sigma_4, \sigma_5, \sigma_6$ and

$$\sigma_8 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, \zeta_6^2 x_4, \zeta_6^4 x_5),$$

with $f_3 = d(x_4^3 + x_5^3)$ for some $d \neq 0$ and

$$q_1 = q_2 = q_3 = x_4 x_5.$$

8. $\text{Aut}(X) \simeq \text{GL}_2(\mathbb{F}_3)$ generated by $\iota, \sigma_1, \sigma_3$ and

$$\sigma_9 : (x_1, x_2, x_3, x_4, x_5) \mapsto \left(x_2, \zeta_6^5 x_1, \zeta_6 x_3, \frac{\zeta_6 b x_4 + x_5}{1 - \zeta_6^2}, \frac{\zeta_6 x_4 + b x_5}{1 - \zeta_6^2} \right),$$

with $f_3 = 0, b^2 = -2$, and

$$\begin{aligned} q_1 &= x_4^2 + b x_4 x_5 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 b x_4 x_5 + \zeta_6^4 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 b x_4 x_5 + \zeta_6^2 x_5^2. \end{aligned}$$

9. $\text{Aut}(X) = \mathfrak{S}_3 = \langle \sigma_4, \sigma_5 \rangle, q_1 = q_2 = q_3 = x_4 x_5$ and f_3 such that X is not isomorphic to any cubic in cases (5) and (7).

Proof. Let $X \subset \mathbb{P}^4$ be a 3-nodal cubic threefold given by (7), with $\text{Aut}(X)$ not fixing any node. There exists an exact sequence

$$0 \rightarrow H \rightarrow \text{Aut}(X) \xrightarrow{\rho} \mathfrak{S}_3 \rightarrow 0 \tag{8}$$

and a $\sigma_{123} \in \text{Aut}(X)$ acting transitively on the nodes, so that $\rho(\sigma_{123}) = (1, 2, 3)$. The zeroes of q_1, q_2, q_3 define at most six points on \mathbb{P}_{x_4, x_5}^1 ; thus,

$$\sigma_{123} = \begin{pmatrix} 0 & 0 & s_3 & 0 & 0 \\ s_1 & 0 & 0 & 0 & 0 \\ 0 & s_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & \zeta_6^r \end{pmatrix} \tag{9}$$

for some $s_1, s_2, s_3 \in k^\times$, where $\zeta_6 = e^{\frac{2\pi i}{6}}$. We have the following cases:

(a) $\text{gcd}(q_1, q_2, q_3) = 1$. We may assume that

$$q_1 = x_4^2 + b x_4 x_5 + x_5^2, \quad b \in k, \quad b \neq \pm 2.$$

The cyclic action on x_1, x_2 and x_3 implies that q_2 and q_3 are multiples of $\sigma_{123}^*(q_1)$ and $\sigma_{123}^*(q_2)$, respectively, and $\sigma_{123}^*(f_3) = s_1s_2s_3f_3$. The torus action on the coordinates x_1, x_2 and x_3 allows us to assume that

$$\begin{aligned} q_2 &= \sigma_{123}^*(q_1), \\ q_3 &= \sigma_{123}^*(q_2). \end{aligned}$$

Since q_1, q_2, q_3 are coprime, we have $r \neq 0, 3$. Thus, $r = 1$ or $r = 2$.

- o If $r = 1$, then $b = 0$; the entries in (9) are $s_1 = s_2 = s_3 = \pm 1$, and

$$\begin{aligned} q_1 &= x_4^2 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 x_5^2, \end{aligned}$$

There are subcases:

- $f_3(x_4, x_5) \neq 0$. Then σ_{123} fixes the points defined by f_3 in \mathbb{P}^1 . And up to isomorphism, $f_3 = dx_4^3$ or $dx_4x_5^2$, for some $d \neq 0$. Since $\sigma_{123}^*(f_3) = s_1s_2s_3f_3$, the latter is impossible. So $f_3 = dx_4^3$ and $s_1 = 1$. This gives $\rho(\text{Aut}(X)) = C_3$. However, any $\gamma \in H$ takes the form

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (t_1x_1, t_2x_2, t_3x_3, t_4x_4 + t_5x_5, t_6x_4 + t_7x_5),$$

for some $t_j \in k^\times$. Since γ leaves (7) invariant, one finds $H = C_2 = \langle \sigma_2^3 \rangle$ and $\text{Aut}(X) \simeq C_6 = \langle \sigma_2 \rangle$, where

$$\sigma_2: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, \zeta_6 x_5).$$

- $f_3(x_4, x_5) \equiv 0$. Then

$$H \simeq C_2^2 = \langle \iota, \sigma_2^3 \rangle, \quad \text{Aut}(X) \simeq C_3 \rtimes \mathcal{D}_4 = \langle \sigma_2, \sigma_7 \rangle,$$

where

$$\iota: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, -x_4, -x_5), \tag{10}$$

$$\sigma_7: (x_1, x_2, x_3, x_4, x_5) \mapsto (\zeta_6^4 x_2, \zeta_6^2 x_1, x_3, \zeta_6^2 x_5, \zeta_6 x_4). \tag{11}$$

- o If $r = 2$, then

$$\begin{aligned} q_1 &= x_4^2 + bx_4x_5 + x_5^2, \\ q_2 &= x_4^2 + \zeta_6^2 bx_4x_5 + \zeta_6^4 x_5^2, \\ q_3 &= x_4^2 + \zeta_6^4 bx_4x_5 + \zeta_6^2 x_5^2, \end{aligned}$$

$b \neq 0$ and $s_1 = s_2 = s_3 = \pm 1$. When $b \neq 1$, q_1, q_2, q_3 define six points in \mathbb{P}^1 , but when $b = 1$, they define three points. There are subcases:

- $f_3(x_4, x_5) = el_1l_2l_3$, $e \in k^\times$. Then σ_{123} permutes the points defined by f_3 in \mathbb{P}^1 ; that is,

$$l_1 = x_4 + dx_5, \quad l_2 = x_4 + \zeta_6^2 dx_5, \quad l_3 = x_4 + \zeta_6^4 dx_5, \quad d \in k^\times.$$

In this case, σ_{123} takes the form

$$\sigma_1: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, \zeta_6^2 x_5).$$

One finds that $H = 0$ and

$$\text{Aut}(X) \simeq \begin{cases} C_3 = \langle \sigma_1 \rangle & d^6 \neq 1, \\ \mathfrak{S}_3 = \langle \sigma_1, \sigma_3 \rangle, & d^3 = 1, \\ \mathfrak{S}_3 = \langle \sigma_1, \sigma'_3 \rangle, & d^3 = -1. \end{cases}$$

$$\sigma_3 : (x_1, x_2, x_3, x_4, x_5) \mapsto (\zeta_6^2 x_2, \zeta_6^4 x_1, x_3, \zeta_6^4 x_5, \zeta_6^2 x_4),$$

$$\sigma'_3 : (x_1, x_2, x_3, x_4, x_5) \mapsto (\zeta_6^2 x_2, \zeta_6^4 x_1, x_3, \zeta_6 x_5, \zeta_6^5 x_4).$$

- $f_3 = l_1 l_2^2$. Then σ_{123} fixes two points defined by f_3 in \mathbb{P}^1 , and $f_3 = dx_4^2 x_5$ or $dx_4 x_5^2$, for some $d \in k^\times$. But then (7) cannot be σ_{123} -invariant. So this case does not exist.
- $f_3 = l^3$. Then $f_3 = dx_4^3$ or dx_5^3 , for some $d \neq 0$. One finds

$$H \simeq 0, \quad \text{Aut}(X) \simeq C_3 = \langle \sigma_1 \rangle.$$

- $f_3 \equiv 0$. Then $\text{Aut}(X)$ contains the involution ι from (10). Up to a twist by ι , we may assume $\sigma_{123} = \sigma_1$. Note that $\text{Aut}(X)$ also contains σ_3 . Using the same argument to find H as above, one gets that when $b^2 \neq -2$,

$$H \simeq C_2 = \langle \iota \rangle, \quad \text{Aut}(X) \simeq C_2 \times \mathfrak{S}_3 = \langle \iota, \sigma_1, \sigma_3 \rangle;$$

when $b^2 = -2$,

$$H \simeq Q_8, \quad \text{Aut}(X) \cong \text{GL}_2(\mathbb{F}_3) = \langle \iota, \sigma_1, \sigma_3, \sigma_9 \rangle,$$

$$\sigma_9 : (x_1, x_2, x_3, x_4, x_5) \mapsto \left(x_2, \zeta_6^5 x_1, \zeta_6 x_3, \frac{\zeta_6 b x_4 + x_5}{1 - \zeta_6^2}, \frac{\zeta_6 x_4 + b x_5}{1 - \zeta_6^2} \right).$$

- (b) When $q_1 = l_1 l$, $q_2 = l_2 l$ and $q_3 = l_3 l$ and l_1, l_2, l_3 are coprime. Then $\text{Aut}(X)$ fixes the point in $\mathbb{P}^1_{x_4, x_5}$ defined by l and acts as C_3 on the three points defined by l_1, l_2 and l_3 . This implies that $r = 2$ in (9), and that

$$l = x_4, \quad l_1 = x_4 + x_5, \quad l_2 = x_4 + \zeta_3 x_5, \quad l_3 = x_4 + \zeta_3^2 x_5.$$

Then either $f_3 = l^3$ defines one point and σ_{123} fixes the point, or $f_3 = l'_1 l'_2 l'_3$, defining three distinct points, with σ_{123} permuting them (i.e., $f_3 = ax_4^3 + bx_5^3$ for some $a, b \in k$). Since X is 3-nodal, one has $b \neq 0$ and $(a, b) \neq (0, 1)$. From the form of f_3 , one sees $\sigma_{123} = \sigma_1$. Here, $H = 0$ since any element in H fixes four points on \mathbb{P}^1 , defined by l, l_1, l_2, l_3 , and acts trivially on x_4, x_5 . Moreover, one can show that no action on \mathbb{P}^1 fixes two points defined by l and l_1 and swaps those defined by l_2 and l_3 at the same time. Therefore, $\rho(\text{Aut}(X)) = C_3$ and

$$H \simeq 0, \quad \text{Aut}(X) \simeq C_3 = \langle \sigma_1 \rangle.$$

- (c) $q_1 = q_2 = q_3 = q$: We may assume that $q = x_4 x_5$. In this case, the exact sequence (8) splits and

$$\text{Aut}(X) \simeq H \times \mathfrak{S}_3,$$

with the factor \mathfrak{S}_3 acting via permutations of x_1, x_2 and x_3 and trivially on x_4, x_5 . Moreover, it is easy to see that H must act faithfully on $\mathbb{P}^1_{x_4, x_5}$. Since H preserves the pair of points defined by q in $\mathbb{P}^1_{x_4, x_5}$, it is either cyclic or dihedral. Assume that $H \neq 1$. Then the structure of $\text{Aut}(X)$ depends on f_3 as follows:

- $f_3 = dl^3$, for some $d \neq 0$ and linear form l in x_4 and x_5 . Then H has a fixed point (i.e., H is a cyclic group). Moreover, H swaps two points, and thus, $H \simeq C_2$ with $l = x_4 + x_5$ and H is generated by swapping coordinates x_4 and x_5 .
- $f_3 = dl_1^2l_2$, for some $d \neq 0$ and linear forms l_1 and l_2 defining two distinct points in \mathbb{P}^1 . Then H fixes two points defined by l_1 and l_2 , and swaps two points defined by q . Similarly, we have $H \simeq C_2$ with

$$l_1 = x_4 - x_5, \quad l_2 = x_4 + x_5,$$

where H is generated by swapping x_4 and x_5 .

- $f_3 = dl_1l_2l_3$, defining three distinct points. There are subcases:
 - H permutes three points defined by f_3 and swaps two points defined by q . Then $H \simeq \mathfrak{S}_3$, generated by

$$\begin{aligned} \sigma_8 &: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, \zeta_3x_4, \zeta_3^2x_5) \\ \sigma_6 &: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, x_5, x_4), \end{aligned}$$

and

$$l_1 = x_4 + x_5, \quad l_2 = \zeta_3x_4 + \zeta_3^2x_5, \quad l_3 = \zeta_3^2x_4 + \zeta_3x_5. \tag{12}$$

- H permutes three points defined by f_3 and fixes two points defined by q ; thus, $H \simeq C_3$, and l_1, l_2, l_3 are as in (12). However, we know that X admits an additional symmetry swapping two points in q as in the case above.
- H fixes the point defined by l_1 . Then H swaps two points defined by q and two points defined by l_2 and l_3 because otherwise, H is trivial. In this case $H \simeq C_2$, with

$$l_1 = x_4 + x_5, \quad l_2 = ax_4 + bx_5, \quad l_3 = bx_5 + ax_4,$$

for some $a, b \neq 0$, and $(\frac{a}{b})^3 \neq 1$; H is generated by the involution swapping x_4 and x_5 .

□

Del Pezzo fibration

We have an $\text{Aut}(X)$ -equivariant commutative diagram:

$$\begin{array}{ccc} \widetilde{X} & \xrightarrow{\varrho} & Y \\ \pi \downarrow & & \downarrow \phi \\ X & \xrightarrow{\rho} & \mathbb{P}^1 \end{array}$$

where π is a blowup of the nodes of X , ϱ is a composition of flops in the strict transforms of the lines

$$\{x_1 = x_4 = x_5 = 0\}, \quad \{x_2 = x_4 = x_5 = 0\}, \quad \{x_3 = x_4 = x_5 = 0\},$$

ϕ is a fibration into Del Pezzo surfaces of degree 6, and ρ is the projection given by

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_4, x_5).$$

The anticanonical model of \widetilde{X} is a singular Fano threefold of degree 18 that has three nodes, which can be smoothed to a smooth Fano threefold of the same degree with Picard rank 1.

Fixed point obstruction

Among actions in Proposition 4.1, we find one example where the linearizability is obstructed by the absence of fixed points upon restriction to abelian subgroups.

Example 4.2. Consider the 3-nodal X in Case (7), Proposition 4.1, and the $G = C_3^2 = \langle \sigma_4, \sigma_8 \rangle$ action on it. The G -action does not have a fixed point on X and $\tilde{X}^G = \emptyset$. By Lemma 2.1, the G -action on X is not linearizable.

Specialization

Here, we exhibit specialization to the 9-nodal cubic with C_3 -action giving an (H1)-obstruction to stable linearizability.

Proposition 4.3. Let $\mathcal{X} \rightarrow \mathbb{A}_k^1$ be a family of cubic threefolds $X_b := \mathcal{X}_b$ given by

$$f_b := x_1x_2x_3 + (x_1 + x_2 + x_3)x_4x_5 + (x_4 + x_5)(x_4 + bx_5)(bx_4 + x_5) = 0$$

for $b \in k$. Consider the $G = C_3$ action on X_b generated by

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, x_5).$$

Then a very general element in \mathcal{X} is not G -stably linearizable.

Proof. Arguing as in Example 2.10, let $\mathcal{X} \rightarrow \mathbb{A}_k^1$ be the family given by f_b . For a very general b , \mathcal{X}_b is a 3-nodal cubic described as Case (5) in Proposition 4.1. The special fiber \mathcal{X}_0 is a 9-nodal cubic, and the G -action fixes a smooth genus 1 curve on \mathcal{X}_0 . From computations in Section 9, there exists an (H1)-obstruction to stable linearizability of the G -action on \mathcal{X}_0 . The six additional nodes form two G -orbits with trivial stabilizer. By Proposition 2.9, a very general element in the family \mathcal{X} is not G -stably linearizable. □

5. Four nodes

Factorial cubics

We first consider the case when the four nodes are in general linear position, forming a ‘tetrahedron’. This is case (J4) in [37]. We may assume that the nodes of X are contained in the hyperplane $x_5 = 0$ and are the points

$$[1 : 0 : 0 : 0 : 0], [0 : 1 : 0 : 0 : 0], [0 : 0 : 1 : 0 : 0], [0 : 0 : 0 : 1 : 0].$$

The intersection $X \cap \{x_5 = 0\}$ is the unique cubic surface with four nodes, the Cayley cubic surface. Using this, we see that X can be given by

$$x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4 + ax_5^3 + x_5^2(b_1x_1 + b_2x_2 + b_3x_3 + b_4x_4) + x_5(a_1(x_1x_2 + x_3x_4) + a_2(x_1x_3 + x_2x_4) + a_3(x_1x_4 + x_2x_3)) = 0 \tag{13}$$

for some $a, b_1, b_2, b_3, b_4, a_1, a_2, a_3 \in k$.

Theorem 5.1. Suppose that $X \subset \mathbb{P}^4$ is a 4-nodal cubic threefold and $\text{Aut}(X)$ does not fix any node of X . Then, up to a change of coordinates, one of the following holds:

(C2) $b_1 = b_2$ and $b_3 = b_4$ in (13), and $\text{Aut}(X) \simeq C_2$, generated by

$$\sigma_1 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_1, x_4, x_3, x_5).$$

(C₄) $a = 0, a_1 = a_2 = a_3 = 0, b_1 = -b_2, b_3 = -b_4$ in (13), and $\text{Aut}(X) \simeq C_4$, generated by

$$\sigma'_1: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_1, x_4, x_3, ix_5), \quad i = e^{\frac{2\pi i}{4}}.$$

(C₂²) $b_1 = b_2 = b_3 = b_4$ in (13), and $\text{Aut}(X) \simeq C_2^2$, generated by σ_1 and

$$\sigma_2: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_3, x_4, x_1, x_2, x_5).$$

(C₈) $a = 0, a_1 = a_2 = a_3 = 0, b_1 = 1, b_2 = -\zeta_8^2, b_3 = -1, b_4 = \zeta_8^2$ in (13), and $\text{Aut}(X) \simeq C_8$, generated by

$$\sigma'_3: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_4, x_1, x_2, x_3, \zeta_8 x_5), \quad \zeta_8 = e^{\frac{2\pi i}{8}}.$$

(D₄) $a_1 = a_3 = 0, a_2 = 1, b_1 = b_2 = b_3 = b_4$ in (13), and $\text{Aut}(X) \simeq D_4$, generated by σ_1, σ_2 and

$$\sigma_3: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_4, x_1, x_2, x_3, x_5).$$

(S₄) $a \neq 0, a_1 = a_2 = a_3 = 0, b_1 = b_2 = b_3 = b_4 = 1$ in (13), and $\text{Aut}(X) \simeq S_4$, generated by $\sigma_1, \sigma_2, \sigma_3$ and

$$\sigma_4: (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, x_4, x_5).$$

Proof. Let $\phi: \text{Aut}(X) \rightarrow S_4$ be the homomorphism given by the action on the nodes of X . Since $\text{Aut}(X)$ does not fix nodes, we may assume that there is a $\iota \in \text{Aut}(X)$ such that $\phi(\iota) = (12)(34)$ or $\phi(\iota) = (1234)$.

Suppose that $\phi(\iota) = (12)(34)$. Then ι is given by

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_2 + sx_5, x_1 + sx_5, x_3 + sx_5, x_4 + sx_5, tx_5),$$

for some $s, t \in k$ such that $t \neq 0$. Considering how ι acts on (13), we see that $s = 0$ or $a_1 = a_2 = a_3$. In the former case, we have $b_1 = b_2$ and $b_3 = b_4$, which implies $t = 1$, because otherwise, $t = -1$ and $a = a_1 = a_2 = a_3 = 0$, which implies that X is not 4-nodal. Thus, if $(s, t) = (0, 1)$ and $\text{im}(\phi) \simeq C_2$, then we are in the case (C₂).

If $a_1 = a_2 = a_3$, then after a coordinate change, we may assume that $a_1 = a_2 = a_3 = 0$. In this case, we get

$$0 = s = a(1 - t^3) = b_3 - b_4t^2 = b_4 - b_3t^2 = b_2 - b_1t^2 = b_1 - b_2t^2.$$

Since X is 4-nodal, this gives $a = 0, b_1 = -b_2, b_3 = -b_4$ and $t = \pm i$. Hence, if $\text{im}(\phi) \simeq C_2$, then we are in the case (C₄).

Now, we suppose that $\phi(\iota) = (1234)$. Then ι is given by

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_4 + sx_5, x_1 + sx_5, x_2 + sx_5, x_3 + sx_5, tx_5)$$

for some $s, t \in k$ such that $t \neq 0$. Then

- $a = 2a_2b_4 - a_2^3, a_1 = 2a_2 - a_3, b_1 = b_2 = b_3 = b_4$, or
- $a_1 = a_3, b_1 = b_2 = b_3 = b_4$, or
- $a_1 = a_2 = a_3$.

In the former case, X is not 4-nodal. If $a_1 = a_2 = a_3$, then after a coordinate change, we may assume that $a_1 = a_2 = a_3 = 0$, which gives

$$0 = s = a(1 - t^3) = b_2 - b_3t^2 = b_3 - b_4t^2 = b_4 - b_1t^2 = b_1 - b_2t^2,$$

so, after an appropriate scaling of x_5 , we see that

- $a \neq 1, b_1 = b_2 = b_3 = b_4 = 1$, or
- $a = 0, b_1 = 1, b_2 = -1, b_3 = 1, b_4 = -1, t = i$, or
- $a = 0, b_1 = 1, b_2 = -\zeta_8^2, b_3 = -1, b_4 = \zeta_8^2, t = \zeta_8$,

which implies that we are in cases $(\mathfrak{S}_4), (C_4), (C_8)$, respectively.

If $a_1 = a_3$ and $b_1 = b_2 = b_3 = b_4$, then after a coordinate change, we may assume that $a_1 = a_3 = 0$. If $a_2 = 0$, then we are in the case (\mathfrak{S}_4) . Finally, if $a_2 \neq 0$, then, scaling x_5 , we may further assume that $a_1 = 1$, so X is given by

$$x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4 + ax_5^3 + b_1x_5^2(x_1 + x_2 + x_3 + x_4) + x_5(x_1x_3 + x_2x_4) = 0,$$

which gives $\text{Aut}(X) = \langle \sigma_1, \sigma_2, \sigma_3 \rangle \cong \mathfrak{D}_4$, so we are in the case (\mathfrak{D}_4) .

To proceed, we may assume that $\text{im}(\phi) \neq C_2$ and $\text{im}(\phi) \neq C_4$. Then, up to a coordinate change, one of the following four cases holds:

- $\text{im}(\phi) = \langle (12)(34), (14)(23) \rangle \simeq C_2^2$,
- $\text{im}(\phi) = \langle (12)(34), (1234) \rangle \simeq \mathfrak{D}_4$,
- $\text{im}(\phi) = \langle (12)(34), (14)(23), (123) \rangle \simeq \mathfrak{A}_4$,
- $\text{im}(\phi) = \langle (12)(34), (14)(23), (1234), (123) \rangle \simeq \mathfrak{S}_4$.

Since $\text{im}(\phi)$ contains $(12)(34)$ or (1234) , the cubic X must be given by one of the equations explicitly described above. Using additional symmetries of X , we conclude that we are in one of the cases $(C_2^2), (\mathfrak{D}_4), (\mathfrak{S}_4)$, or the cubic X is given by (13) with

$$a \neq 0, \quad a_1 = a_2 = a_3 = 0, \quad b_1 = b_3 = 1, \quad b_2 = b_4 = -1,$$

or

$$a_1 = 1, \quad a_2 = \zeta_3, \quad a_3 = \zeta_3^2, \quad b_1 = b_2 = b_3 = b_4 = 0.$$

In the first of the latter two cases, X has eight nodes, and in the last case, the singularities of X are not nodes. This completes the proof of the theorem. □

Birational model

Let $\pi: \tilde{X} \rightarrow X$ be the blowup of the nodes of X . Then there exists an $\text{Aut}(X)$ -equivariant diagram:

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\rho} & \hat{X} \\ \pi \downarrow & & \downarrow \phi \\ X & & Y \end{array}$$

where ρ is a composition of flops in the strict transform of the lines passing through pair of nodes, ϕ is a contraction of the strict transform of the hyperplane section containing four nodes (the surface $X \cap \{x_5 = 0\}$) to a smooth point of the threefold Y , and Y is a smooth divisor in $(\mathbb{P}^1)^4$ of degree $(1, 1, 1, 1)$. Implicitly, the birational map $X \dashrightarrow Y$ has been constructed in the proof of [48, Proposition 4.5]. Note that the anticanonical model of \tilde{X} is a singular Fano threefold with six nodes of degree 16, which can be smoothed to a smooth Fano threefold of degree 16 and Picard rank 1.

Burnside formalism

We realize the situation of Proposition 2.6 in some of the 4-nodal cases.

Example 5.2. Let X be the cubic threefold given in the Case (\mathfrak{D}_4) or (\mathfrak{S}_4) in Theorem 5.1. Consider the group $G \subset \text{Aut}(X)$ where $G = \langle \sigma_2, \sigma_1\sigma_3 \rangle = C_2^2$. Then we are in the situation of Proposition 2.6, and the G -action is not linearizable. In particular, $\sigma_1\sigma_3$ fixes a cubic surface receiving a residual σ_2 -action with a G -fixed elliptic curve on it.

Specialization

One can equivariantly specialize 4-nodal cubic threefolds to an 8-nodal one:

Proposition 5.3. *Let \mathcal{X}_b be the 4-nodal cubic threefold defined by*

$$f_b = x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4 + x_5^2(x_1 + x_2 + b(x_3 + x_4)) = 0.$$

For all $b \in k$, \mathcal{X}_b carries a $G = C_2$ -action generated by

$$\sigma_1 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_1, x_4, x_3, x_5).$$

Then \mathcal{X}_b is not G -stably linearizable for a very general b .

Proof. Let $\mathcal{X} \rightarrow \mathbb{A}_k^1$ be the family given by f_b . The generic fiber \mathcal{X}_b is a 4-nodal cubic of the type (C_2) in Theorem 5.1. The special fiber $\mathcal{X}_{9/4}$ is an 8-nodal cubic, with an **(H1)**-obstruction to stable linearizability of the G -action by Corollary 8.3. The additional four nodes have trivial stabilizer and thus are BG -rational singularities. Applying Proposition 2.9 and Example 2.8, one concludes that a very general member in the family \mathcal{X} is not G -stably linearizable. □

One can also specialize to the Segre cubic threefold:

Proposition 5.4. *Let \mathcal{X}_a be a cubic of type (\mathfrak{S}_4) in Theorem 5.1; that is, \mathcal{X}_a is given by*

$$ax_5^3 + x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4 + x_5^2(x_1 + x_2 + x_3 + x_4) = 0.$$

Consider the subgroup $G = \langle \sigma_1, \sigma_2 \rangle \simeq C_2^2 \subset \text{Aut}(X_a)$. Then, for a very general $a \in k$, the G -action on \mathcal{X}_a is not stably linearizable.

Proof. Let $\mathcal{X} \rightarrow \mathbb{A}_k^1$ be the family consisting of X_a . The special fiber cX_0 is a 10-nodal cubic, with an **(H1)**-obstruction to stable linearizability of the G -action, from computations in [24]. The additional six nodes have C_2 -stabilizers. They are BG -rational singularities, by Example 2.8. Applying Proposition 2.9 and Example 2.8, one concludes that a very general fiber is not G -stably linearizable. □

Remark 5.5. We note that the degeneration of cubics in Proposition 5.4 is equivalent to the degeneration of divisors in $(\mathbb{P}^1)^4$ of degree $(1, 1, 1, 1)$, which was studied in [48, Section 7] and [17]. In particular, the product of projections from four planes in the tetrahedron formed by the four nodes of the cubics gives an G -equivariant birational map from the cubics to divisors in $(\mathbb{P}^1)^4$ of degree $(1, 1, 1, 1)$.

Cubics with a plane

Now we treat the case when the four nodes are contained in a distinguished, G -stable plane Π . This is case (J6) in [37]. Unprojecting from Π , we have a G -equivariant birational map

$$\phi : X \dashrightarrow X_{2,2},$$

where $X_{2,2}$ is a smooth complete intersection in \mathbb{P}^5 of two quadrics with a G -fixed point $P \in X_{2,2}$, and the map ϕ^{-1} is a projection from P . Linearizability of actions on smooth $X_{2,2}$ is determined by existence of invariant lines [42, Theorem 24]. In particular, we have the following:

Proposition 5.6. *The G -action on X is not linearizable if and only if no singular points of X are fixed by G , and X does not contain G -stable lines that are disjoint from Π .*

Proof. We may assume that no singular point of X is G -fixed. If X contains a G -stable line that is disjoint from Π , then the G -action on $X_{2,2}$ is linearizable, by Lemma 1.1. Conversely, if the G -action on $X_{2,2}$ is linearizable, then it follows from [42, Theorem 24] that $X_{2,2}$ contains a G -stable line ℓ . And $P \notin \ell$ because otherwise, the preimage of ℓ on X would be a G -fixed singular point. Similarly, we see that ℓ must be disjoint from the four lines in $X_{2,2}$ containing P . Then ℓ is mapped by ϕ^{-1} to a G -stable line in X that is disjoint from the plane Π . □

Examples of nonlinearizable actions, based on the Burnside formalism [46] or the adaptation to the equivariant context of the *torsors over intermediate Jacobians* formalism from [43], [11], can be found in [42, Sections 8.3 and 8.4]. As a special case, we have the following:

Example 5.7. We may assume that $X_{2,2}$ is given by

$$\sum_{i=1}^6 a_i x_i^2 = \sum_{i=1}^6 x_i^2 = 0.$$

Let $G = \langle \sigma \rangle$, with σ acting diagonally by $(1, 1, 1, 1, -1, -1)$. Then G does not leave invariant any line on $X_{2,2}$, and the action is not linearizable. However, there is a genus 1 curve C fixed by G , obtained by intersecting $X_{2,2}$ with $x_5 = x_6 = 0$. Projecting from any of the points on C , we obtain a singular cubic threefold, generically with four nodes.

Example 5.8. Let $X \subset \mathbb{P}^4_{y_1, \dots, y_5}$ be the 4-nodal cubic given by

$$(y_1 - y_3)y_2y_4 + (y_2 - y_3)y_1y_5 + (y_4 - y_5)y_4y_5 - y_4^3 - y_5^3.$$

The four nodes lie on the unique plane $y_4 = y_5 = 0$. The automorphism group $\text{Aut}(X)$ contains $G = C_2^3$ generated by

$$\iota_1 : (y_1, y_2, y_3, y_4, y_5) \mapsto (-y_1, -y_1 + y_3, -y_1 + y_2, y_4, y_5)$$

$$\iota_2 : (y_1, y_2, y_3, y_4, y_5) \mapsto (y_2 - y_3, y_1 - y_3, -y_3, y_4, y_5).$$

and

$$\iota_3 : (y_1, y_2, y_3, y_4, y_5) \mapsto (y_1, y_2, y_3, -y_4, -y_5).$$

Unprojecting X from the unique plane under the map

$$(y_1, \dots, y_5) \mapsto (y_1y_5, y_2y_5, y_3y_5, y_4y_5, y_5^2, y_1y_2 - y_2y_3 - y_4^2),$$

one sees that X is G -equivariantly birational to a smooth intersection of two quadrics $X_{2,2} \subset \mathbb{P}^5_{x_1, \dots, x_6}$ given by

$$x_1x_2 - x_2x_3 - x_4^2 - x_5x_6 = x_1x_2 - x_1x_3 + x_4^2 - x_4x_5 - x_5^2 + x_4x_6 = 0.$$

The $G = C_2^3$ action on the first five coordinates is the same as that on \mathbb{P}^4 , ι_1 and ι_2 acts trivially on x_6 and ι_3 changes the sign of x_6 . For any subgroup $G' \subset G$, there is a G' -stable line in $X_{2,2}$ if and only if $G' = C_2$ and the character of the G' -representation of the ambient $\mathbb{A}^6_{x_1, \dots, x_6}$ is

$$(6, 0) \quad \text{or} \quad (6, 4).$$

In the first case, G' fixes a singular point of X and thus is linearizable. In the latter case, G' pointwise fixes a smooth intersection of two quadrics in dimension 2 (i.e., a quartic Del Pezzo surface), which contains 16 lines. The other C_2 subgroups have character $(6, 2)$. They fix an elliptic curve but do not leave any line invariant in $X_{2,2}$. Any of the other subgroups of G will contain one of the nonlinearizable C_2 .

6. Five nodes

Now, we suppose that X has five nodes.

Birational model

If the nodes are not in general linear position, then there is a distinguished G -fixed node, and the G -action on X is linearizable. Hence, we may assume that the nodes of X are

$$p_1 = [1 : 0 : 0 : 0 : 0], \quad p_2 = [0 : 1 : 0 : 0 : 0], \quad p_3 = [0 : 0 : 1 : 0 : 0],$$

$$p_4 = [0 : 0 : 0 : 1 : 0], \quad p_5 = [0 : 0 : 0 : 0 : 1].$$

Then $G \subseteq \mathfrak{S}_5$ acts via permutation of coordinates. We may also assume that G does not fix any of the nodes, since otherwise, the G -action is clearly linearizable.

Linearizability

Using the standard Cremona involution

$$\iota : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4,$$

we obtain a G -birational map $\chi : X \dashrightarrow Q$, where $Q \subset \mathbb{P}^4$ is a smooth quadric. For more details of this map, see the proof of Theorem 2.12.

Lemma 6.1. *Suppose that G does not act transitively on $\text{Sing}(X)$. Then the G -action on X is linearizable.*

Proof. Since G does not fix any of the nodes, either $G \simeq C_2 \times \mathfrak{S}_3$ or $G \simeq C_2 \times C_3$. In both cases, we may assume that G preserves the subset $\{p_1, p_2\}$ and $\{p_3, p_4, p_5\}$. Then G pointwise fixes the line $l \subset \mathbb{P}^4$ that passes through the points $[1 : 1 : 0 : 0 : 0]$ and $[0 : 0 : 1 : 1 : 1]$. Observe that $\iota(l) = l$, so that the intersection $l \cap Q$ contains G -fixed points, which implies the assertion. \square

Thus, we may assume that G acts transitively on the nodes of X , and G contains the 5-cycle $(1, 2, 3, 4, 5)$. Then X is defined by

$$x_1x_2x_3 + x_2x_3x_4 + x_1x_2x_5 + x_1x_4x_5 + x_3x_4x_5 + a(x_1x_2x_4 + x_1x_3x_4 + x_1x_3x_5 + x_2x_3x_5 + x_2x_4x_5) = 0,$$

for some a . And Q is defined by

$$x_1x_2 + x_2x_3 + \dots + x_5x_1 + a(x_1x_3 + x_2x_4 + \dots + x_5x_2) = 0. \tag{14}$$

Note that $a \neq -1$, since otherwise, X would be 6-nodal. Then Q is smooth. For the group G , we have the following possibilities:

- (1) $G \simeq C_5$,
- (2) $G \simeq \mathfrak{D}_5$,
- (3) $G \simeq C_4 \rtimes C_5$ and $a = 1$,
- (4) $G \simeq \mathfrak{A}_5$ and $a = 1$,
- (5) $G \simeq \mathfrak{S}_5$ and $a = 1$.

In the first case, $G = C_5$, the group G fixes a point in Q , and the G -action on X is linearizable. In the second case, the action is necessarily of the form in the following lemma:

Lemma 6.2. *Suppose that $G \simeq \mathfrak{D}_5$ acting on $\mathbb{P}^4 = \mathbb{P}(\mathbb{I} \oplus V_2 \oplus V'_2)$, where V_2 and V'_2 are two nonisomorphic 2-dimensional irreducible representations of \mathfrak{D}_5 . Then the G -action on every G -invariant smooth quadric in \mathbb{P}^4 is linearizable.*

Proof. We may assume the G action is generated by

$$\begin{aligned} (x_1, \dots, x_5) &\mapsto (x_4, x_3, x_2, x_1, x_5), \\ (x_1, \dots, x_5) &\mapsto (\zeta x_1, \zeta^2 x_2, \zeta^3 x_3, \zeta^4 x_4, x_5), \end{aligned}$$

where $\zeta = e^{\frac{2\pi i}{5}}$. Smooth G -invariant quadrics $Q_{a,b}$ are given by

$$ax_1x_4 - bx_2x_3 + x_5^2 = 0$$

for $a, b \neq 0$. Notice that each $Q_{a,b}$ is \mathfrak{D}_5 -isomorphic to $Q_{1,1}$ with the same \mathfrak{D}_5 -action under a change of variables

$$x'_1 = \sqrt{a}x_1, x'_2 = \sqrt{b}x_2, x'_3 = \sqrt{b}x_3, x'_4 = \sqrt{a}x_4, x'_5 = x_5.$$

Consider a G -invariant conic


$$C = \{x_2 = x_3 = 0\} \cap Q_{1,1}$$

and a G -invariant twisted cubic curve

$$R = \{x_5 = x_1x_3 - x_2^2 = x_2x_4 - x_3^2 = 0\} \cap Q_{1,1}.$$

The system of quadric hypersurfaces on \mathbb{P}^4 containing both C and R induces a G -equivariant birational map $Q_{1,1} \dashrightarrow \mathbb{P}^3$; see, for example, [3, Section 5.10]. □

Lemma 6.3. *Suppose that $G \simeq C_4 \rtimes C_5$ and $a = 1$. Then Q from (14) contains a G -invariant smooth quintic elliptic curve E , and we have the following G -Sarkisov link:*



where α is a blowup of the curve E , and β is a blowup of a smooth quintic elliptic curve isomorphic to E .

Proof. It is easy to see that $\text{Aut}(Q)$ contains a unique subgroup isomorphic to $C_4 \rtimes C_5$. Thus, we may change coordinates on \mathbb{P}^4 as we need and, in particular, assume that Q is given by

$$\sum_{i=1}^5 x_i^2 + i \sum_{1 \leq i < j \leq 5} x_i x_j = 0,$$

and that the action of G on Q is given by

$$\begin{aligned} (x_1, x_2, x_3, x_4, x_5) &\mapsto (x_2, x_3, x_4, x_5, x_1), \\ (x_1, x_2, x_3, x_4, x_5) &\mapsto (x_1, x_3, x_5, x_2, x_4). \end{aligned}$$

Then Q contains the following smooth quintic elliptic curve:

$$\begin{cases} x_1^2 + i(x_3x_4 + x_2x_5) = 0, \\ x_2^2 + i(x_4x_5 + x_3x_1) = 0, \\ x_3^2 + i(x_5x_1 + x_4x_2) = 0, \\ x_4^2 + i(x_1x_2 + x_5x_3) = 0, \\ x_5^2 + i(x_2x_3 + x_1x_4) = 0. \end{cases}$$

Blowing up Q along this curve, we obtain the claim; cf. [19]. □

If $G \simeq \mathfrak{S}_5$ and $a = 1$, then it follows from [21] that X is G -solid, and the only G -Mori fiber spaces G -birational to X are X and Q . In particular, the G -action is not linearizable. If $G \simeq \mathfrak{A}_5$ and $a = 1$, we also expect that X and Q are the only G -Mori fiber spaces G -birational to X , which would imply that the G -action is not linearizable.

7. Six nodes

Cubics without planes


Let X be the 6-nodal cubic threefold such that the nodes are in general linear position. Then $\text{rk Cl}(X) = 2$, so the defect of X is 1. This is case (J9) in [37]. Note that X does not contain planes, but it contains two families of cubic scrolls (see Remark 7.1 below). Moreover, by [41, Section 3], X can be given by

$$\det(M) = 0$$

for a 3×3 matrix M whose entries are linear forms. Thus, one can define a rational map $X \dashrightarrow \mathbb{P}^2$ that maps $p \mapsto (a, b, c)$, where (a, b, c) is a nonzero solution of the equation

$$M \begin{pmatrix} a \\ b \\ c \end{pmatrix} = 0.$$

This map is dominant, it is undefined at the nodes of X , and its general fiber is a line in X . Similarly, we can define another rational map $X \dashrightarrow \mathbb{P}^2$ using the transpose of the matrix M . Taking resolution of singularities X , we resolve indeterminacy of both of these rational maps, which yields the following commutative diagram:



where f is the standard resolution, q^+ and q^- are small resolutions, h^+ and h^- are birational morphisms such that $h^- \circ (h^+)^{-1}$ is a composition of six Atiyah flops, and both p^+ and p^- are \mathbb{P}^1 -bundles. The diagram (15) is implicitly contained in [44, §7.5], as an illustration of the first row in the table there. Taking a product of morphisms $p^+ \circ h^+$ and $p^- \circ h^-$, we obtain a morphism $\widetilde{X} \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$ that is birational onto its image (a divisor of degree $(2, 2)$ with 15 nodes).

Remark 7.1. Let l be a general line in \mathbb{P}^2 . Set

$$\bar{S} = (q^-)_*(p^-)^*(l) \quad \text{and} \quad \bar{S}' = (q^+)_*(p^+)^*(l).$$

Then \bar{S} and \bar{S}' are smooth cubic scrolls in X that freely generate the class group $\text{Cl}(X)$.

Remark 7.2. Let $G \subseteq \text{Aut}(X)$. Then the commutative diagram (15) is G -equivariant if and only if $\text{rk Cl}^G(X) \neq 1$.

To describe possibilities for $\text{Aut}(X)$, we can assume that the nodes of X are the points

$$\begin{aligned} p_1 &= [1 : 0 : 0 : 0 : 0], & p_2 &= [1 : 1 : 1 : 1 : 1], & p_3 &= [0 : 0 : 0 : 0 : 1], \\ p_4 &= [0 : 0 : 0 : 1 : 0], & p_5 &= [0 : 0 : 1 : 0 : 0], & p_6 &= [0 : 1 : 0 : 0 : 0]. \end{aligned}$$

Fix the \mathfrak{S}_6 -action on \mathbb{P}^4 generated by

$$\begin{aligned} \tau_{(12)} &: (x_1, \dots, x_5) \mapsto (-x_1, -x_1 + x_2, -x_1 + x_3, -x_1 + x_4, -x_1 + x_5), \\ \tau_{(1\dots 6)} &: (x_1, \dots, x_5) \mapsto (-x_1 + x_2, -x_1 + x_3, -x_1 + x_4, -x_1 + x_5, -x_1), \end{aligned} \tag{16}$$

where the indices correspond to the permutation of six nodes. Then $\text{Sing}(X)$ forms an \mathfrak{S}_6 -orbit, but X is not \mathfrak{S}_6 -invariant. Moreover, it follows from a classical construction [30] that there exists the following 4-dimensional \mathfrak{S}_6 -Sarkisov link:

$$\begin{array}{ccc} U & \xrightarrow{\beta} & Y \\ \alpha \downarrow & & \downarrow \gamma \\ \mathbb{P}^4 & \xrightarrow{\chi} & V \end{array}$$

where V is the 10-nodal Segre cubic threefold in \mathbb{P}^4 , χ is given by the linear system of cubic hypersurfaces singular at the points p_1, \dots, p_6 , α is the blowup of p_1, \dots, p_6 , β is a composition of antiflips in the strict transforms of the 15 lines that contain two points among p_1, \dots, p_6 , and γ is a \mathbb{P}^1 -bundle.

Observe that $\text{Aut}(X) \subseteq \mathfrak{S}_6$. Restricting the above \mathfrak{S}_6 -Sarkisov link to X , we obtain the following $\text{Aut}(X)$ -equivariant diagram:

$$\begin{array}{ccc} \tilde{X} & \dashrightarrow & \widehat{X} \\ f \downarrow & & \downarrow \pi \\ X & & S \end{array} \tag{17}$$

where S is a smooth hyperplane section of the Segre cubic V , $\tilde{X} \dashrightarrow \widehat{X}$ is a composition of Atiyah flops in the strict transforms of the 15 lines in X that contains two nodes among p_1, \dots, p_6 , π is a \mathbb{P}^1 -bundle. For more details, see [41].

Our cubic X is given by

$$a_1 f_1 + a_2 f_2 + a_3 f_3 + a_4 f_4 + a_5 f_5 = 0 \tag{18}$$

for some $a_1, a_2, a_3, a_4, a_5 \in k$, where

$$\begin{aligned} f_1 &= x_1x_2x_3 - x_2x_3x_4 - x_2x_3x_5 - x_1x_4x_5 + x_2x_4x_5 + x_3x_4x_5, \\ f_2 &= x_1x_2x_4 - x_2x_3x_4 - x_1x_4x_5 + x_3x_4x_5, \\ f_3 &= x_1x_2x_5 - x_2x_3x_5 - x_1x_4x_5 + x_3x_4x_5, \\ f_4 &= x_1x_3x_4 - x_2x_3x_4 - x_1x_4x_5 + x_2x_4x_5, \\ f_5 &= x_1x_3x_5 - x_2x_3x_5 - x_1x_4x_5 + x_2x_4x_5. \end{aligned}$$

Enumerating $G \subseteq \mathfrak{S}_6$ and searching for G -invariant cubics singular at p_1, \dots, p_6 , we can find all possibilities for $\text{Aut}(X)$. In particular, $\text{Aut}(X) = 1$ for general a_1, \dots, a_5 . Moreover, one has the following:

Proposition 7.3. *Let $X \subset \mathbb{P}^4$ be a 6-nodal cubic threefold given by (18). Assume that none of the nodes of X is fixed by $\text{Aut}(X)$. Then under the \mathfrak{S}_6 -action specified in (16), one of the following holds:*

1. $a_1 + a_2 + a_4 + a_5 = 0$, and

$$\text{Aut}(X) \simeq C_2 = \langle (1, 3)(2, 5)(4, 6) \rangle.$$

2. $a_1 + a_3 = a_2 - a_3 + a_4 + a_5 = 0$, and

$$\text{Aut}(X) \simeq \mathfrak{S}_3 = \langle (1, 3)(2, 5)(4, 6), (1, 4, 5)(2, 6, 3) \rangle.$$

3. $a_1 + a_4 = a_2 + a_5 = a_3 - a_4 = 0$, and

$$\text{Aut}(X) \simeq \mathfrak{S}_4 = \langle (1, 3)(2, 5)(4, 6), (3, 4, 5, 6) \rangle.$$

4. $a_1 + a_4 = a_2 + a_5 = a_3 + a_4 = 0$, and

$$\text{Aut}(X) \simeq \mathfrak{D}_4 = \langle (3, 5), (1, 3, 2, 5)(4, 6) \rangle.$$

5. $a_1 + a_4 + 2a_5 = a_2 - a_5 = 0$, and

$$\text{Aut}(X) \simeq C_2^2 = \langle (1, 2)(3, 5)(4, 6), (1, 3)(2, 5)(4, 6) \rangle.$$

6. $a_1 + a_4 + 2a_5 = a_2 - a_5 = a_3 - a_4 - 2a_5 = 0$, and

$$\text{Aut}(X) \simeq \mathfrak{D}_6 = \langle (1, 3)(2, 5)(4, 6), (1, 6, 5, 2, 4, 3) \rangle.$$

7. $a_1 = a_3 = a_5 = 1, a_2 = a_4 = -1$, and

$$\text{Aut}(X) \simeq \mathfrak{S}_3^2 \rtimes C_2 = \langle (1, 3)(2, 5)(4, 6), (2, 4), (1, 5)(2, 3, 4, 6) \rangle.$$

8. $a_1 = a_4 = 1, a_2 = a_3 = a_5 = -1$, and

$$\text{Aut}(X) \simeq \mathfrak{S}_5 = \langle (1, 3)(2, 5)(4, 6), (1, 2, 5, 6, 4) \rangle.$$

Proof. Enumerating all (conjugacy classes of) subgroups G of \mathfrak{S}_6 which do not fix any point among p_1, \dots, p_6 , and computing all G -invariant cubics singular at p_1, \dots, p_6 , we obtain the list of (families of) 6-nodal cubics whose automorphism groups do not fix any of the nodes. These are the eight families of cubics listed above. Since $\text{Aut}(X) \subset \mathfrak{S}_6$, one can find the full automorphism groups $\text{Aut}(X)$. \square

As in [4], we find two maximal subgroups \mathfrak{S}_5 and $\mathfrak{S}_3^2 \rtimes C_2$ such that (up to conjugation in \mathfrak{S}_6) G and X can be described as follows:

1. $G = \mathfrak{S}_5 = \langle (1, 3)(2, 5)(4, 6), (1, 2, 5, 6, 4) \rangle$ and X is given by

$$x_1x_2x_3 - x_1x_2x_4 + x_1x_3x_4 - x_2x_3x_4 - x_1x_2x_5 - x_1x_3x_5 + x_2x_3x_5 + x_1x_4x_5 + x_2x_4x_5 - x_3x_4x_5 = 0, \tag{19}$$

2. $G = \mathfrak{S}_3^2 \rtimes C_2 = \langle (1, 3)(2, 5)(4, 6), (2, 4), (1, 5)(2, 3, 4, 6) \rangle$ and X is given by

$$x_1x_2x_3 - x_1x_2x_4 - x_1x_3x_4 + x_2x_3x_4 + x_1x_2x_5 + x_1x_3x_5 + x_2x_4x_5 - 3x_2x_3x_5 - x_1x_4x_5 + x_3x_4x_5 = 0. \tag{20}$$

In the first case, $\text{Aut}(X) \simeq \mathfrak{S}_5$, $\text{rk Cl}^{\mathfrak{S}_5}(X) = 1$, and (17) is a \mathfrak{S}_5 -Sarkisov link such that S is the Clebsch diagonal cubic surface. In the second case, $\text{Aut}(X) \simeq \mathfrak{S}_3^2 \rtimes C_2$, $\text{rk Cl}^{\mathfrak{S}_3^2 \rtimes C_2}(X) = 1$, and (17) is a $\mathfrak{S}_3^2 \rtimes C_2$ -Sarkisov link such that S is the Fermat cubic surface.

Lemma 7.4. *Let $X \subset \mathbb{P}^4$ be a 6-nodal cubic threefold such that the nodes are in general linear position. If $\text{Aut}(X)$ contains an involution σ not fixing any node, then $\text{rk Cl}^{\langle \sigma \rangle}(X) = 1$.*

Proof. Since σ does not fix any node, we may assume that

$$\sigma = (1, 3)(2, 5)(4, 6)$$

and X is one of the cases in Proposition 7.3. From the diagram (17), we know that

$$\text{rk Cl}^{\langle \sigma \rangle}(X) + 3 = \text{rk Cl}^{\langle \sigma \rangle}(\tilde{X}) = \text{rk Cl}^{\langle \sigma \rangle}(\hat{X}) = \text{rk Cl}^{\langle \sigma \rangle}(S) + 1,$$

where S is a smooth cubic surface contained in a hyperplane $H \subset \mathbb{P}^4$. By Lefschetz fixed-point theorem, one has [36, Section 6]

$$\begin{aligned} \text{rk Cl}^{\langle \sigma \rangle}(S) &= \frac{1}{2}(7 + \text{Tr}_2(\sigma^*)), \\ \text{Tr}_2(\sigma^*) &= s - 2 + \sum_i (2 - 2g_i), \end{aligned}$$

where $\text{Tr}_2(\sigma^*)$ is the trace of σ^* -action on $H^2(S, \mathbb{C})$, s is the number of isolated σ -fixed points on S , and g_i are the genera of fixed curves. In our case, we compute that the induced σ -action on $H \simeq \mathbb{P}^3$ has weights $(1, 1, 1, -1)$. The fixed locus S^σ consists of one point and a smooth cubic curve. Substituting into the formulae above we obtain

$$\text{rk Cl}^{\langle \sigma \rangle}(S) = 3,$$

which implies $\text{rk Cl}^{\langle \sigma \rangle}(X) = 1$. □

Proposition 7.5. *Let $X \subset \mathbb{P}^4$ be a 6-nodal cubic threefold such that the nodes are in general linear position and $\text{Aut}(X)$ contains an involution σ . Let \tilde{X} be the standard resolution of X . Then the action of $\langle \sigma \rangle \simeq C_2$ on $\text{Pic}(\tilde{X})$ fails (HI) if and only if σ does not fix any node.*

Proof. We know that $\text{Pic}(\tilde{X})$ is generated by the pullback of the hyperplane section H , six exceptional divisors E_1, \dots, E_6 , and the classes of the strict transforms of two cubic scrolls S and S' (see Remark 7.1), subject to the relation

$$2H = S + S' + \sum_{i=1}^6 E_i.$$

There is a short exact sequence of $\text{Aut}(X)$ -modules

$$0 \rightarrow \bigoplus_{i=1}^6 E_i \rightarrow \text{Pic}(\tilde{X}) \rightarrow \text{Cl}(X) \rightarrow 0,$$

giving rise to the long exact sequence of cohomology groups

$$\dots \rightarrow H^1(\langle \sigma \rangle, \bigoplus_{i=1}^6 E_i) \rightarrow H^1(\langle \sigma \rangle, \text{Pic}(\tilde{X})) \rightarrow H^1(\langle \sigma \rangle, \text{Cl}(X)) \rightarrow H^2(\langle \sigma \rangle, \bigoplus_{i=1}^6 E_i) \rightarrow \dots$$

By our assumption, σ permutes the E_i without fixing any E_i . So

$$H^1(\langle \sigma \rangle, \bigoplus_{i=1}^6 E_i) = H^2(\langle \sigma \rangle, \bigoplus_{i=1}^6 E_i) = 0,$$

and

$$H^1(\langle \sigma \rangle, \text{Pic}(\tilde{X})) = H^1(\langle \sigma \rangle, \text{Cl}(X)).$$

If σ does not fix any node, Lemma 7.4 implies that $\text{rk Cl}^{\langle \sigma \rangle}(X) = 1$. So σ acts on $\text{Cl}(X)$ via

$$\sigma(H) = H, \quad \sigma(S) = S' = 2H - S.$$

In another basis of $\text{Cl}(X)$ (namely, H and $H - S$) the action becomes

$$\sigma(H) = H, \quad \sigma(H - S) = -H + S.$$

Then

$$H^1(\langle \sigma \rangle, \text{Pic}(\tilde{X})) = H^1(\langle \sigma \rangle, \text{Cl}(X)) = \mathbb{Z}/2.$$

Conversely, if $\langle \sigma \rangle$ fails **(H1)**, it is not stably linearizable and thus cannot fix any node. □

Example 7.6. Let $X \subset \mathbb{P}^4$ be a 6-nodal cubic threefold in one of the eight cases in Proposition 7.3. Then $\text{Aut}(X)$ contains the involution

$$\sigma = (1, 3)(2, 5)(4, 6),$$

satisfying the conditions in Proposition 7.5. Therefore, the σ -action on any 6-nodal cubic is not stably linearizable.

Cubics with a plane

This is case (J8) in [37]. Four of the six nodes of X are contained in a unique, and thus G -stable plane $\Pi \subset X$. The other two are on a G -stable line ℓ . Note that $\ell \cap \Pi = \emptyset$, since otherwise, the hyperplane containing Π and ℓ would intersect X by three planes. So, the action of $\text{Aut}(X)$ on X is linearizable by Lemma 1.1.


Cubics with three planes

Let X be a cubic threefold in \mathbb{P}^4 with six nodes such that X contains three planes Π_1, Π_2, Π_3 . Then X belongs to a four-parameter family, which is denoted by (J11) in [37]. It follows from [37] that

$\Pi_1 + \Pi_2 + \Pi_3$ is cut out by a hyperplane. Thus, we may assume that this hyperplane is $\{x_1 = 0\}$, and

$$\begin{aligned} \Pi_1 &= \{x_1 = 0, x_2 = 0\}, \\ \Pi_2 &= \{x_1 = 0, x_3 = 0\}, \\ \Pi_3 &= \{x_1 = 0, x_4 = 0\}. \end{aligned}$$

Observe the existence of the following diagram:



where π is the standard resolution, Y is the double cover of $(\mathbb{P}^1)^3$ branched over a singular divisor of degree $(2, 2, 2)$, and η is a birational morphism that contracts the strict transforms of Π_1, Π_2, Π_3 . Note that $\text{Aut}(Y)$ contains a Galois involution of the double cover, and this involution acts biregularly on X , which follows from the following:

Proposition 7.7. *Up to a change of coordinates, X is given by*

$$x_2x_3x_4 + ax_1^3 + x_1^2(b_1x_2 + b_2x_3 + b_3x_4) + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0, \tag{21}$$

for some a, b_1, b_2, b_3 .

Proof. A priori, the cubic X is given by

$$\begin{aligned} x_2x_3x_4 + ax_1^3 + x_1^2(b_1x_2 + b_2x_3 + b_3x_4 + cx_5) + x_1(e_1x_2^2 + e_2x_3^2 + e_3x_4^2 + e_4x_2x_3 + e_5x_2x_4 + e_6x_3x_4) + \\ + x_1(x_5(d_1x_2 + d_2x_3 + d_3x_4) - x_5^2) = 0 \end{aligned}$$

for some $a, b_1, b_2, b_3, c, e_1, e_2, e_3, e_4, e_5, e_6, d_1, d_2, d_3$. Changing x_2, x_3, x_4 , we may assume that $e_4 = e_5 = e_6 = 0$. Moreover, up to scaling, there exists a unique such choice of coordinates x_2, x_3, x_4 that preserves the equations of the planes Π_1, Π_2, Π_3 .

Similarly, changing the coordinate x_5 , we may further assume that $c = d_1 = d_2 = d_3 = 0$. As above, we see that there is a unique such choice for x_5 up to scaling.

Finally, using the fact that X has six nodes, we see that $e_1 \neq 0, e_2 \neq 0, e_3 \neq 0$. Hence, scaling the coordinates x_1, x_2, x_3, x_4, x_5 , we may also assume that $e_1 = e_2 = e_3 = 1$, which completes the proof. \square

Remark 7.8. If we permute b_1, b_2, b_3 in (21), or simultaneously change signs of two of them, we obtain an isomorphic cubic threefold.

From now on, we assume that the cubic threefold X is given by (21). Then the nodes of X are

$$\begin{aligned} p_1 &= [0 : 0 : 0 : 1 : 1], & p_2 &= [0 : 0 : 0 : -1 : 1], \\ p_3 &= [0 : 0 : 1 : 0 : 1], & p_4 &= [0 : 0 : -1 : 0 : 1], \\ p_5 &= [0 : 1 : 0 : 0 : 1], & p_6 &= [0 : -1 : 0 : 0 : 1]. \end{aligned}$$

Remark 7.9. Let S_3 be the cubic surface $\{x_5 = 0\} \cap X$. Then S_3 is smooth. The proof of Proposition 7.7 shows that S_3 is $\text{Aut}(X)$ -invariant.

Remark 7.10. One can find an explicit condition on a, b_1, b_2, b_3 that guarantees that (21) defines a 6-nodal cubic, but it is too bulky to present here. However, if the equation (21) has additional symmetries,

the condition simplifies a lot. For instance, if $b_1 = b_2 = b_3 = b$, then (21) defines a 6-nodal cubic if and only if

$$(4a - b^2 + 8b + 16)(4b^3 + a^2 - 6ab - 3b^2 + 4a) \neq 0.$$

Moreover, in this very special case, we have the following possibilities:

1. if $4b^3 + a^2 - 6ab - 3b^2 + 4a = 0$ and $(a, b) \notin \{(1, 1), (-4, 0), (-8, 28)\}$, then (21) defines a 7-nodal cubic;
2. if $(a, b) = (1, 1)$, then (21) defines a cubic with six nodes and one double non-nodal singularity;
3. if $4a - b^2 + 8b + 16 = 0$ and $(a, b) \notin \{(-4, 0), (-8, 28)\}$, then (21) defines a 9-nodal cubic;
4. if $(a, b) \in \{(-4, 0), (-8, 28)\}$, then (21) defines the Segre cubic.

As we mentioned earlier, $\text{Aut}(X)$ is never trivial since it contains the involution:

$$\iota_5 : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, x_4, -x_5).$$

Moreover, if b_1, b_2, b_3 in (21) are general enough, then $\text{Aut}(X) = \langle \iota_5 \rangle$. In fact, we can say more:

Proposition 7.11. *Suppose $\text{Aut}(X) \neq \langle \iota_5 \rangle$. Then, up to a permutation of coordinates x_2, x_3, x_4 and changing signs of two of them, one of the following holds:*

1. $b_1 \neq b_2, b_2 = b_3, b_1, b_2 \neq 0$, so X is given by

$$x_2x_3x_4 + ax_1^3 + x_1^2(b_1x_2 + b_2(x_3 + x_4)) + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0,$$

and $\text{Aut}(X) \simeq C_2^2$, generated by ι_5 and

$$\sigma_{34} : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_4, x_3, x_5);$$

2. $b_1 \neq 0, b_2 = b_3 = 0$, so X is given

$$x_2x_3x_4 + ax_1^3 + b_1x_1^2x_2 + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0,$$

and $\text{Aut}(X) \simeq C_2^3$, generated by ι_5, σ_{34} and

$$\iota_{34} : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, -x_3, -x_4, x_5);$$

3. $b_1 = b_2 = b_3 \neq 0$, so X is given by

$$x_2x_3x_4 + ax_1^3 + b_1x_1^2(x_2 + x_3 + x_4) + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0,$$

and $\text{Aut}(X) \simeq C_2 \times \mathfrak{S}_3$, generated by ι_5, σ_{34} and

$$\sigma_{234} : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_3, x_4, x_2, x_5);$$

4. $b_1 = b_2 = b_3 = 0$, so X is given by

$$x_2x_3x_4 + ax_1^3 + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0,$$

and $\text{Aut}(X) \simeq C_2 \times \mathfrak{S}_4$, generated by $\iota_5, \sigma_{34}, \sigma_{234}, \iota_{34}$ and

$$\iota_{24} : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, -x_2, x_3, -x_4, x_5).$$

Proof. Permuting the coordinates x_2, x_3, x_4 , we may assume that one of the following four cases hold:

1. $b_1 \neq 0, b_2 \neq 0, b_3 \neq 0$;
2. $b_1 = 0, b_2 \neq 0, b_3 \neq 0$;

- 3. $b_1 = 0, b_2 = 0, b_3 \neq 0$;
- 4. $b_1 = 0, b_2 = 0, b_3 = 0$.

In the first two cases, we may assume b_2 and b_3 have the same sign by changing the signs of two among three variables x_2, x_3 and x_4 .

There is a natural homomorphism $\phi: \text{Aut}(X) \rightarrow \mathfrak{S}_3$ given by the action of $\text{Aut}(X)$ on the planes Π_1, Π_2, Π_3 . Arguing as in the proof of Proposition 7.7, we see that an element in the kernel of ϕ is given by

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, \lambda_1 x_2, \lambda_2 x_3, \lambda_3 x_4, \lambda_4 x_5)$$

for some nonzero $\lambda_1, \lambda_2, \lambda_3, \lambda_4$. Using this, we see that the kernel of ϕ can be described as follows:

- if $b_2 \neq 0, b_3 \neq 0$, then $\ker(\phi) = \langle \iota_5 \rangle \simeq C_2$,
- if $b_1 = b_2 = 0, b_3 \neq 0$, then $\ker(\phi) = \langle \iota_5, \iota_{23} \rangle \simeq C_2^2$,
- if $b_1 = b_2 = b_3 = 0$, then $\ker(\phi) = \langle \iota_5, \iota_{23}, \iota_{24} \rangle \simeq C_2^3$.

Let $G = \text{Aut}(X)$. First, assume $(2, 3) \in \phi(G)$ (i.e., there exists an element $\sigma \in G$ swapping the planes Π_2 and Π_3 and leaving Π_1 invariant). Then σ takes the form

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ s_1 & s_9 & 0 & 0 & 0 \\ s_2 & 0 & 0 & s_{10} & 0 \\ s_3 & 0 & s_{11} & 0 & 0 \\ s_4 & s_5 & s_6 & s_7 & s_8 \end{pmatrix}$$

for parameters s_1, \dots, s_{11} . Note that σ^2 is contained in the kernel of ϕ , which implies that $s_8 = \pm 1$. Moreover, we may assume that $s_8 = 1$ by replacing σ by $\sigma \circ \iota_5$. Furthermore, the fact that σ leaves X invariant imposes relations on the parameters. Solving for the equations, we obtain solutions

- $b_2 = b_3, s_1 = \dots = s_7 = 0, s_{10} = s_{11} = s_9 = 1$,
- $b_2 = -b_3, s_1 = \dots = s_7 = 0, s_{10} = s_{11} = -1, s_9 = 1$.

Similarly, if $(1, 2, 3) \in \phi(G)$ (i.e., there exists an element $\sigma \in G$ translating three planes). As above, we see that σ takes the form

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ s_1 & 0 & 0 & s_9 & 0 \\ s_2 & s_{10} & 0 & 0 & 0 \\ s_3 & 0 & s_{11} & 0 & 0 \\ s_4 & s_5 & s_6 & s_7 & 1 \end{pmatrix}$$

In this case, we obtain four solutions

- $b_1 = b_2 = -b_3, s_1 = \dots = s_7 = 0, s_9 = s_{11} = -1, s_{10} = 1$,
- $b_1 = -b_2 = -b_3, s_1 = \dots = s_7 = 0, s_9 = s_{10} = -1, s_{11} = 1$,
- $b_1 = -b_2 = b_3, s_1 = \dots = s_7 = 0, s_9 = 1, s_{10} = s_{11} = -1$,
- $b_1 = b_2 = b_3, s_1 = \dots = s_7 = 0, s_9 = s_{10} = s_{11} = 1$.

Combining these solutions with symmetries, we obtain the result. □

Linearization

Let X be the 6-nodal cubic given by (21). In this subsection, we solve the linearizability problem for subgroups in $\text{Aut}(X)$, almost completely. We use notation introduced in the previous subsection, and let S_3 be the cubic surface $\{x_5\} \cap X$. Then S_3 is smooth by Remark 7.9, which implies the following:

Lemma 7.12. *Let $G = \langle \iota_5 \rangle$. Then the G -action on X is linearizable.*

Proof. The surface S_3 is pointwise fixed by ι_5 , and $\Pi_1 \cap S_3$ is a line. Since S_3 is smooth, it contains another line l disjoint from $\Pi_1 \cap S_3$. Hence, l is disjoint from Π_1 . Since Π_1 is G -invariant, the G -action is linearizable by Lemma 1.1. \square

Similarly, we prove the following:

Lemma 7.13. *Suppose that $b_2 = b_3 = 0$, so X is given by*

$$x_2x_3x_4 + ax_1^3 + b_1x_1^2x_2 + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0.$$

Let $G = \langle \iota_5, \iota_{34} \rangle \simeq C_2^2$. Then the G -action on X is linearizable.

Proof. Note that G leaves invariant the planes Π_1, Π_2, Π_3 . As in the proof of Lemma 7.12, we see that S_3 contains a G -invariant line that is disjoint from one of these planes. Indeed, if r is a root of $r^2 + b_1r + a = 0$, then S_3 contains the reducible conic

$$x_5 = x_2 - rx_1 = rx_3x_4 + x_3^2 + x_4^2 = 0,$$

and its irreducible components are G -invariant lines disjoint from Π_2 and Π_3 . Hence, the G -action is linearizable by Lemma 1.1. \square

Now, let us discuss nonlinearizable actions. We start with

Lemma 7.14. *Suppose that $b_2 = b_3$, and let $G = \langle \iota_5, \sigma_{34} \rangle \cong C_2^2$. Then the G -action on X is not linearizable.*

Proof. The involution ι_5 pointwise fixes the G -invariant surface S_3 , while the involution σ_{34} pointwise fixes the cubic curve

$$C = \{x_3 = x_4\} \cap S_3 \subset S_3.$$

One can check that a singular point on C is also a singular point of X . Then C is necessarily smooth since the six nodes on X are away from S_3 . Therefore, C is a genus 1 curve, and by Proposition 2.6, the G -action is not linearizable. \square

Remark 7.15. The same argument shows that the following two G -actions on X (when they act) are not linearizable:

$$G = \langle \iota_5, \sigma_{34}\iota_{34} \rangle \simeq C_2^2, \quad \text{and} \quad G = \langle \sigma_{234}, \sigma_{34} \rangle \simeq \mathfrak{S}_3.$$

In each of the two cases, there is a cubic surface in X fixed by a subgroup $C_2 \subset G$ and receiving a G/C_2 -residual action which fixes an elliptic curve. Therefore, Proposition 2.6 is also applicable to these cases.

Using Proposition 7.11 and Lemma 7.12, we obtain the following:

Corollary 7.16. *The action of $\text{Aut}(X)$ on X is linearizable if and only if $\text{Aut}(X) = \langle \iota_5 \rangle$.*

We proceed to the actions of other subgroups of the full automorphism groups from Proposition 7.11.

Lemma 7.17. *Let $\mathcal{X}_a \subset \mathbb{P}^4$ be a 6-nodal cubic threefold given by*

$$ax_1^3 + b(x_2 + x_3 + x_4)x_1^2 + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) + x_2x_3x_4 = 0, \tag{22}$$

for some $b \notin \{0, 28\}$, and let $G \simeq C_3$ be a group acting on \mathbb{P}^4 by

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_4, x_3, x_2, x_5).$$

Then X is G -invariant, and the G -action on \mathcal{X}_a is not stably linearizable for a very general $a \in k$.

Proof. Fixing $b \notin \{0, 28\}$, consider the family $\mathcal{X} \rightarrow \mathbb{A}_k^1$ whose fiber over $a \in k$ is the cubic given by (22). From Remark 7.10, we know that the generic fiber \mathcal{X}_a is 6-nodal if a is very general. However, the special fiber \mathcal{X}_ϵ is 9-nodal when $\epsilon = \frac{b^2}{4} - 2b - 4$ and the additional three nodes have trivial stabilizer.

Now set $\epsilon = \frac{b^2}{4} - 2b - 4$. Then the G -action on planes in \mathcal{X}_ϵ has a G -orbit of length 3, consisting of three planes

$$\Pi_4 = \{x_2 - 2x_1 = (bx_1 + 2x_3 + 2x_4 + 2x_5) = 0\},$$

$$\Pi_5 = \{x_3 - 2x_1 = (bx_1 + 2x_4 + 2x_2 + 2x_5) = 0\},$$

$$\Pi_6 = \{x_4 - 2x_1 = (bx_1 + 2x_2 + 2x_3 + 2x_5) = 0\}.$$

Moreover, $\Pi_4 + \Pi_5 + \Pi_6$ is a Cartier divisor on \mathcal{X}_ϵ . Then by Remark 9.1, the G -action on \mathcal{X}_ϵ is not stably linearizable. Applying Proposition 2.9, we conclude that the G -action on \mathcal{X}_a is not stably linearizable for a very general $a \in k$. □

Lemma 7.18. *Let $\mathcal{X}_a \subset \mathbb{P}^4$ be a cubic threefold given by*

$$ax_1^3 + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) + x_2x_3x_4 = 0. \tag{23}$$

Consider a group $G \simeq C_2^2$ acting on \mathbb{P}^4 via

$$t_{24} : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, -x_2, x_3, -x_4, x_5),$$

$$t_{34} : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, -x_3, -x_4, x_5).$$

Then \mathcal{X}_a is G -invariant, and the G -action on \mathcal{X}_a is not stably linearizable for a very general $a \in k$.

Proof. Consider the family $\mathcal{X} \rightarrow \mathbb{A}_k^1$ whose fiber over $a \in k$ is the cubic given by (23). From Remark 7.10, we know that \mathcal{X}_a is 6-nodal for a very general a . However, the special fiber $\mathcal{X}'_{a'}$, when $a' = -4$, is the Segre cubic with 10 nodes. The G -action on \mathcal{X}_{-4} leaves invariant three planes – namely,

$$\Pi_i = \{x_i = x_i = 0\}, \quad i = 2, 3, 4.$$

The action has an orbit of nodes of length 4 and three orbits of nodes of length 2. By [24, Section 6], the G -action on \mathcal{X}_{-4} does not satisfy **(H1)** and is not stably linearizable. Moreover, the four additional nodes are in one G -orbit and are BG -rational singularities. Applying Proposition 2.9, one concludes that a very general member in the family \mathcal{X} is not G -stably linearizable. □

Remark 7.19. The same argument shows that for the same family of cubics, the action on \mathcal{X}_a of the group

$$G = \langle t_5, t_{23}\sigma_{34} \rangle \simeq C_2 \times C_4$$

for a very general a is not stably linearizable. The action specializes to the unique $C_2 \times C_4$ -action on the Segre cubic \mathcal{X}_{-4} . This action on \mathcal{X}_{-4} does not satisfy **(SP)**. The four additional nodes have stabilizer C_2 , and they are BG -rational singularities; see Example 2.8.

Let us summarize what is left using the notation of Proposition 7.11.

1. When $\text{Aut}(X) \simeq C_2^2$, we are left with $\langle t_5\sigma_{34} \rangle \simeq C_2$,

2. When $\text{Aut}(X) \simeq C_2^3$, we are left with five groups:

Group	Generators	Orbit of nodes	Orbit of planes
C_2	$\iota_5\sigma_{34}$	$2 + 2 + 2$	$1 + 2$
C_2	$\iota_5\sigma_{34}\iota_{34}$	$2 + 2 + 2$	$1 + 2$
C_2^2	$\iota_5\sigma_{34}, \iota_{34}$	$2 + 4$	$1 + 2$
C_2^2	$\iota_5\sigma_{34}, \iota_{34}\sigma_{34}$	$2 + 2 + 2$	$1 + 2$
C_2^2	$\iota_5\iota_{34}, \sigma_{34}$	$2 + 2 + 2$	$1 + 2$

3. When $\text{Aut}(X) \simeq C_2 \times \mathfrak{S}_3$ and a, b_1 are very general, we are left with $\langle \iota_5\sigma_{34} \rangle \simeq C_2$.

4. When $\text{Aut}(X) \simeq C_2 \times \mathfrak{S}_4$, for very general a , we are left with

Group	Generators	Orbit of nodes	Orbit of planes
C_2	$\iota_5\sigma_{34}$	$2 + 2 + 2$	$1 + 2$
C_3	σ_{234}	$3 + 3$	3
C_2^2	$\iota_5\sigma_{34}, \iota_{34}$	$2 + 4$	$1 + 2$
C_4	$\iota_{23}\sigma_{34}$	$2 + 4$	$1 + 2$
C_2^2	$\iota_5\sigma_{34}, \iota_{34}\sigma_{34}$	$2 + 2 + 2$	$1 + 2$
C_6	$\iota_5\sigma_{234}$	6	3
\mathfrak{S}_3	$\iota_5\sigma_{34}, \sigma_{234}$	$3 + 3$	3
\mathfrak{D}_4	$\iota_5\iota_{24}, \iota_{23}\sigma_{34}$	$2 + 4$	$1 + 2$

Specialization in Lemma 7.17 does not apply to the second row.

We also note that in each of the remaining cases, the construction in Lemma 1.1 does not apply. In particular, every G -invariant line in these cases intersects with the G -invariant plane (when it exists) at one point.

8. Eight nodes

The 8-nodal cubic threefolds form a two-parameter family, which is denoted by (J13) in [37]. Let X be one such cubic, and $G = \text{Aut}(X)$. Then $\text{Cl}(X) = \mathbb{Z}^4$, and X contains five planes Π_1, \dots, Π_5 that form a very special configuration [37]. If p_1, \dots, p_8 are the nodes of X , then

$$\begin{aligned} \Pi_1 &\supset \{p_1, p_2, p_6, p_8\}, \\ \Pi_2 &\supset \{p_1, p_2, p_5, p_7\}, \\ \Pi_3 &\supset \{p_5, p_6, p_7, p_8\}, \\ \Pi_4 &\supset \{p_3, p_4, p_5, p_6\}, \\ \Pi_5 &\supset \{p_3, p_4, p_7, p_8\}. \end{aligned}$$

From this configuration, there are two distinguished sets of nodes

$$\{p_1, p_2, p_3, p_4\} \quad \text{and} \quad \{p_5, p_6, p_7, p_8\}. \tag{24}$$

The planes Π_1, Π_2, Π_3 form one tetrahedron (without a face), and Π_3, Π_4, Π_5 form another one. In particular, Π_3 is distinguished, and must be G -invariant.

Unprojecting from Π_3 , we obtain a G -equivariant birational map

$$X \dashrightarrow X_{2,2} = Q \cap Q' \subset \mathbb{P}^5$$

to a singular complete intersection of two quadrics Q and Q' in \mathbb{P}^5 that are singular along lines. The threefold $X_{2,2}$ has four nodes: $Q \cap \text{Sing}(Q')$ and $Q' \cap \text{Sing}(Q)$, and G fixes a point in $X_{2,2}$ – the inverse map $X_{2,2} \dashrightarrow X$ is just a projection from this point. Thus, one could study the geometry of X using $X_{2,2}$ as in Section 5.

Standard form

We change the coordinates in \mathbb{P}^4 so that

$$\Pi_3 = \{x_4 = x_5 = 0\}$$

and

$$p_1 = [0 : 0 : 0 : 1 : 0], \quad p_3 = [0 : 0 : 0 : 0 : 1];$$

this is possible since the line through p_1 and p_3 is disjoint from Π_3 . Changing the variables x_1, x_2, x_3 , we may assume that

$$\begin{aligned} p_5 &= [1 : 1 : 1 : 0 : 0], & p_6 &= [-1 : 1 : 1 : 0 : 0], \\ p_7 &= [1 : -1 : 1 : 0 : 0], & p_8 &= [1 : 1 : -1 : 0 : 0]. \end{aligned}$$

This specifies the equations of the planes:

$$\begin{aligned} \Pi_1 &= \{x_1 + x_3 = x_5 = 0\}, \\ \Pi_2 &= \{x_1 - x_3 = x_5 = 0\}, \\ \Pi_3 &= \{x_4 = x_5 = 0\}, \\ \Pi_4 &= \{x_2 - x_3 = x_4 = 0\}, \\ \Pi_5 &= \{x_2 + x_3 = x_4 = 0\}. \end{aligned}$$

A cubic threefold containing Π_1, \dots, Π_5 and singular at $p_1, p_3, p_5, \dots, p_8$ has the form

$$(a_{22}x_1 + a_{12}x_2 + a_6x_3)x_4x_5 + a_9(x_3^2 - x_1^2)x_4 + a_8(x_3^2 - x_2^2)x_5 = 0,$$

for some $a_6, a_8, a_9, a_{12}, a_{22}$. Since X is 8-nodal, we have

$$a_8, a_9, a_{12}, a_{22} \neq 0.$$

Scaling coordinates, we may assume that $a_8 = a_9 = a_{12} = 1$, and we let $a_{22} = a$ and $a_6 = b$. Thus, $X = X_{a,b}$ is given by

$$(ax_1 + x_2 + bx_3)x_4x_5 + x_4(x_3^2 - x_1^2) + x_5(x_3^2 - x_2^2) = 0, \tag{25}$$

for parameters a, b , where $a \neq 0$. Notice that a and b are defined up to ± 1 . For very general a and b , (25) defines an 8-nodal cubic with nodes at p_1, \dots, p_8 , where $p_1, p_3, p_5, \dots, p_8$ are described above, and

$$p_2 = [0 : 1 : 0 : 1 : 0], \quad \text{and} \quad p_4 = [a : 0 : 0 : 0 : 1].$$

For special parameters a and b , (25) defines a cubic with additional singularities, for instance, the Segre cubic, when $b = 0$ and $a = 1$.

Automorphisms

Recall that $G = \text{Aut}(X)$ and Π_3 is G -invariant. Let l_{12} be the line passing through p_1 and p_2 , and l_{34} the line through p_3 and p_4 . Then the curve $l_{12} + l_{34}$ is G -invariant. However, we have a group homomorphism

$$\phi : G \rightarrow \text{PGL}_3(k),$$

arising from the action of G on Π_3 . Since $\phi(G)$ permutes the points p_5, p_6, p_7, p_8 , we see that $\phi(G) \subseteq \mathfrak{S}_4 \subset \text{PGL}_3(k)$, permuting the coordinates x_1, x_2, x_3 and changing signs of these variables. Moreover, the set $(l_{12} + l_{34}) \cap \Pi_3$ is $\phi(G)$ -invariant, which implies that $\phi(G)$ is contained in $\mathfrak{D}_4 \subset \mathfrak{S}_4$ generated by

$$\begin{aligned} (x_1, x_2, x_3) &\mapsto (-x_2, x_1, x_3), \\ (x_1, x_2, x_3) &\mapsto (-x_1, x_2, x_3). \end{aligned}$$

Lemma 8.1. *The kernel $\ker(\phi)$ of ϕ is nontrivial if and only if $b = 0$. Moreover, if $b = 0$, then $\ker(\phi) \simeq C_2$, generated by*

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 - ax_5, x_2 - x_4, x_3, -x_4, -x_5).$$

Proof. An element $\tau \in \ker(\phi)$ preserves Π_3 , the points p_6, p_7, p_8, p_9 , and each line l_{12} and l_{34} . Moreover, since τ leaves the subsets $\{p_1, p_2\}, \{p_3, p_4\}$ invariant, we have the following three possibilities:

1. $\tau(p_1) = \tau(p_1), \tau(p_2) = \tau(p_2), \tau(p_3) = \tau(p_4), \tau(p_4) = \tau(p_3),$
2. $\tau(p_1) = \tau(p_2), \tau(p_2) = \tau(p_1), \tau(p_3) = \tau(p_3), \tau(p_4) = \tau(p_4),$
3. $\tau(p_1) = \tau(p_2), \tau(p_2) = \tau(p_1), \tau(p_3) = \tau(p_4), \tau(p_4) = \tau(p_3).$

These impose linear conditions on τ . Solving them, we see that τ is one of the following linear transformations:

1. $(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2 - x_4, x_3, -x_4, x_5),$
2. $(x_1, x_2, x_3, x_4, x_5) \mapsto (-ax_5 + x_1, x_2, x_3, x_4, -x_5)$
3. $(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 - ax_5, x_2 - x_4, x_3, -x_4, -x_5).$

However, (25) must be preserved by τ , which implies that only the third case is possible, and only in the case when $b = 0$. □

We are ready to classify all possibilities for $G = \text{Aut}(X)$.

Proposition 8.2. *Let $X \subset \mathbb{P}^4$ be an 8-nodal cubic threefold given by*

$$(ax_1 + x_2 + bx_3)x_4x_5 + x_4(x_3^2 - x_1^2) + x_5(x_3^2 - x_2^2) = 0$$

and $G = \text{Aut}(X)$. Then one of the following holds:

- $b \neq 0$ and $a \neq \pm 1, G \simeq C_2^2 = \langle \iota_1, \iota_2 \rangle$, generated by

$$\begin{aligned} \iota_1 : (x_1, x_2, x_3, x_4, x_5) &\mapsto (ax_5 - x_1, x_2, x_3, x_4, x_5), \\ \iota_2 : (x_1, x_2, x_3, x_4, x_5) &\mapsto (x_1, x_4 - x_2, x_3, x_4, x_5); \end{aligned}$$

- $b \neq 0$ and $a = \pm 1, G \simeq \mathfrak{D}_4$, generated by ι_2 and

$$\sigma_1 : (x_1, x_2, x_3, x_4, x_5) \mapsto (\pm x_2, x_5 \mp x_1, x_3, x_5, x_4);$$

- $b = 0, a \neq \pm i, G \simeq C_2^3$, generated by ι_1, ι_2 and

$$\tau : (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 - ax_5, x_2 - x_4, x_3, -x_4, -x_5);$$

◦ $b = 0, a = \pm i, G \simeq C_2 \cdot \mathfrak{D}_4 \simeq C_2^2 \rtimes C_4$, generated by τ, ι_2 and

$$\sigma_2 : (x_1, x_2, x_3, x_4, x_5) \mapsto (\mp x_2, \pm x_1, x_3, ix_5, ix_4).$$

Proof. Observe that G always contains ι_1 and ι_2 , and $\langle \iota_1, \iota_2 \rangle \cong C_2^2$, which shows that $\phi(G)$ is at least C_2^2 . Moreover, if $b = 0$, then it follows from Lemma 8.1 that G also contains the involution τ , so together, the involutions ι_1, ι_2, τ generate a subgroup C_2^3 in this case. If $\phi(G) \simeq C_2^2$, this gives us all possibilities for G . To complete the proof, we have to find all a and b such that $\phi(G) \simeq \mathfrak{D}_4$ and describe G in these cases. This can be done explicitly.

If $\phi(G) \simeq \mathfrak{D}_4$, then G contains a σ such that $\phi(\sigma)$ is given by

$$(x_1, x_2, x_3) \mapsto (x_2, -x_1, x_3),$$

which implies that σ is given by the matrix

$$\begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & s_{25} \\ 0 & 0 & 1 & 0 & 0 \\ s_{14} & 0 & 0 & 0 & s_{54} \\ 0 & s_{25} & 0 & s_{45} & 0 \end{pmatrix}$$

for some $s_{14}, s_{25}, s_{45}, s_{54}$. Since σ preserves (25), we obtain constraints on these entries, which result in the following possibilities:

1. $a = 1, s_{14} = 0, s_{25} = 1, s_{45} = 1, s_{54} = 1$;
2. $a = -1, s_{14} = -1, s_{25} = 0, s_{45} = 1, s_{54} = 1$;
3. $b = 0, a = -i, s_{14} = 0, s_{25} = 0, s_{45} = i, s_{54} = i$;
4. $b = 0, a = i, s_{14} = -1, s_{25} = i, s_{45} = i, s_{54} = i$.

Using them, we obtain all possibilities for G listed above. □

Cohomology

Let $\tilde{X} \rightarrow X$ be the standard resolution, let E_1, \dots, E_8 be exceptional divisors over p_1, \dots, p_8 , and let $\tilde{\Pi}_1, \dots, \tilde{\Pi}_5$ be the strict transforms of the planes Π_1, \dots, Π_5 on \tilde{X} , respectively. Then $\text{Pic}(\tilde{X})$ is generated by $E_1, \dots, E_8, \tilde{\Pi}_1, \dots, \tilde{\Pi}_5$. These are subject to the relation

$$\tilde{\Pi}_1 + \tilde{\Pi}_2 - \tilde{\Pi}_4 - \tilde{\Pi}_5 = E_1 + E_2 - E_3 - E_4.$$

Notice that this presentation of the lattice $\text{Pic}(\tilde{X})$ is independent of the equation of the cubic threefold X . To compute the (H1)-obstruction on $\text{Pic}(\tilde{X})$, for generality, we may work with the maximal symmetry group appearing in Proposition 8.2. Let $G = C_2^2 \rtimes C_4$, as defined above. Then G acts on nodes via permutation of indices

$$\begin{aligned} \tau &: (1, 2)(3, 4), \\ \iota_1 &: (3, 4)(5, 6)(7, 8), \\ \iota_2 &: (1, 2)(5, 7)(6, 8), \\ \sigma_2 &: (1, 3)(2, 4)(5, 7, 8, 6). \end{aligned}$$

There is a unique (conjugacy class of) $C_2 = \langle \iota_1 \iota_2 \rangle$ contributing to

$$H^1(C_2, \text{Pic}(\tilde{X})) = \mathbb{Z}/2.$$

Indeed, this C_2 acts on nodes via the permutation of indices

$$(1, 2)(3, 4)(5, 8)(6, 7).$$

Under the basis

$$\tilde{\Pi}_1 - \tilde{\Pi}_5 - E_2 + E_4, \tilde{\Pi}_1, \dots, \tilde{\Pi}_5, E_3, \dots, E_8,$$

one can see that C_2 acts on $\tilde{\Pi}_1 - \tilde{\Pi}_5 - E_2 + E_4$ by -1 , and on the rest as a permutation module.

Note that this C_2 is contained in $\text{Aut}(X) = C_2^2$ for generic X in Proposition 8.2 (i.e., when $b \neq 0$ and $a \neq \pm 1$).

Linearization

Let X be an 8-nodal cubic threefold. The classification in Proposition 8.2 implies

- $b \neq 0$ and $a \neq \pm 1$: a subgroup of $\text{Aut}(X) \simeq C_2^2$ is linearizable if and only if it fixes a singular point; otherwise, it fails **(H1)**.
- $b \neq 0$ and $a = \pm 1$: a subgroup of $\text{Aut}(X) \simeq \mathfrak{D}_4$ is linearizable if and only if it fixes a singular point; otherwise, it fails **(H1)**.
- $b = 0$ and $a \neq \pm i$: Excluding subgroups failing **(H1)** or with a fixed singular point, we are left with the following (classes of) subgroups

1. $C_2^2 = \langle (3, 4)(5, 7)(6, 8), (1, 2)(3, 4) \rangle$, acting via

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 + ix_5, -x_2, x_3, -x_4, -x_5),$$

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 + ix_5, x_2 - x_4, x_3, -x_4, -x_5).$$

2. $C_2^2 = \langle \iota_1, \tau \rangle = \langle (3, 4)(5, 6)(7, 8), (1, 2)(3, 4) \rangle$, acting via

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (-x_1 - ix_5, x_2, x_3, x_4, x_5),$$

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 + ix_5, x_2 - x_4, x_3, -x_4, -x_5).$$

- $b = 0$ and $a = \pm i$: Excluding groups with an **(H1)**-obstruction or with a fixed singular point, we are left with

1. $C_2^2 = \langle (3, 4)(5, 6)(7, 8), (1, 2)(3, 4) \rangle$, acting via

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (-x_1 - ix_5, x_2, x_3, x_4, x_5),$$

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_1 + ix_5, x_2 - x_4, x_3, -x_4, -x_5),$$

2. $C_4 = \langle (1, 3)(2, 4)(5, 7, 8, 6) \rangle$, acting via

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, -x_1, x_3, ix_5, ix_4).$$

We turn to linearization constructions for subgroups unobstructed by cohomology and not fixing singular points. Consider the maximal symmetry group, in the case $b = 0$ and $a = \pm i$. We have two unobstructed cases:

- $G = C_2^2 = \langle (3, 4)(5, 6)(7, 8), (1, 2)(3, 4) \rangle$. The group G swaps the planes Π_1 and Π_2 and preserves Π_4 and Π_5 . The line passing through p_1 and p_2 is a G -invariant line disjoint from Π_4 . Then the G -action on X is linearizable by Lemma 1.1.
- $G = C_4 = \langle (1, 3)(2, 4)(5, 7, 8, 6) \rangle$. In this case, G swaps the planes Π_1, Π_2 and swaps Π_4, Π_5 . But Π_3 is G -invariant. The line passing through p_1 and p_3 is a G -invariant line disjoint from Π_3 . The G -action on X is linearizable by Lemma 1.1.

Note that the constructions above only depend on the group actions on singular points and planes. One can also establish the same linearization results for the two unobstructed C_2^2 in the case when $b = 0$ and $a \neq \pm i$. We summarize this section by the following:

Corollary 8.3. *Let X be an 8-nodal cubic threefold and $G \subseteq \text{Aut}(X)$. The G -action on X is linearizable if and only if it satisfies **(HI)**, if and only if G does not contain a subgroup isomorphic to C_2 which does not fix any nodes of X ; in particular, if it is not linearizable then it is not stably linearizable.*

9. Nine nodes

Standard form

We follow [52]: 9-nodal cubic threefolds X_a are given in \mathbb{P}^5 by equations

$$x_1x_2x_3 - x_4x_5x_6 = a(x_1 + x_2 + x_3) + x_4 + x_5 + x_6 = 0, \quad a^3 \neq 0, -1.$$

Their automorphisms depend on the parameter a as follows:

$$\text{Aut}(X_a) = \begin{cases} \mathfrak{S}_3^2 & \text{when } a^3 \neq 1, \\ \mathfrak{S}_3^2 \rtimes C_2 & \text{otherwise.} \end{cases}$$

These groups act via \mathfrak{S}_3 -permutations of two sets of coordinates: x_1, x_2, x_3 , and x_4, x_5, x_6 . When $a = 1$, the additional C_2 switches $x_i \leftrightarrow x_{3+i}$, $i = 1, 2, 3$. In both cases, the nine nodes are given by

$$\{x_{i_1} = x_{i_2} = x_{j_1} = x_{j_2} = 0, \quad x_{j_3} + ax_{i_3} = 0\},$$

where

$$i_1 \neq i_2 \neq i_3 \in \{1, 2, 3\}, \quad j_1 \neq j_2 \neq j_3 \in \{4, 5, 6\}.$$

There are also nine distinguished planes, given by

$$\Pi_{i,j} = \{x_i = x_{3+j} = 0\} \cap X, \quad i, j \in \{1, 2, 3\}.$$

The G -action on X_a fixes a singular point if and only if G is a 2-group.

Fixed point obstruction

Let $G = C_3^2$ be the group generated by

$$(x_1, x_2, x_3, x_4, x_5, x_6) \mapsto (x_3, x_1, x_2, x_4, x_5, x_6),$$

$$(x_1, x_2, x_3, x_4, x_5, x_6) \mapsto (x_1, x_2, x_3, x_6, x_4, x_5).$$

Then $X_a^G = \emptyset$, for all a such that $a^3 \neq 0, -1$. By Lemma 2.1, the G -action on X is not linearizable.

Cohomology

Let $\tilde{X}_a \rightarrow X_a$ be the blowup of X_a at nine nodes. Then $\text{Pic}(\tilde{X}_a)$ is generated by $E_i, i = 1, \dots, 9$, the exceptional divisors over the nine nodes, the pullbacks $\tilde{\Pi}_{i,j}$ of $\Pi_{i,j}$, and H , the pullback of the hyperplane section. They are subject to relations

$$\begin{aligned}
 H &= \tilde{\Pi}_{1,1} + \tilde{\Pi}_{1,2} + \tilde{\Pi}_{1,3} + E_2 + E_3 + E_5 + E_6 + E_8 + E_9, \\
 H &= \tilde{\Pi}_{1,1} + \tilde{\Pi}_{2,1} + \tilde{\Pi}_{3,1} + E_1 + E_2 + E_3 + E_4 + E_5 + E_6, \\
 H &= \tilde{\Pi}_{1,2} + \tilde{\Pi}_{2,2} + \tilde{\Pi}_{3,2} + E_1 + E_2 + E_3 + E_7 + E_8 + E_9, \\
 H &= \tilde{\Pi}_{1,3} + \tilde{\Pi}_{2,3} + \tilde{\Pi}_{3,3} + E_4 + E_5 + E_6 + E_7 + E_8 + E_9, \\
 H &= \tilde{\Pi}_{2,1} + \tilde{\Pi}_{2,2} + \tilde{\Pi}_{2,3} + E_1 + E_3 + E_4 + E_6 + E_7 + E_9, \\
 H &= \tilde{\Pi}_{3,1} + \tilde{\Pi}_{3,2} + \tilde{\Pi}_{3,3} + E_1 + E_2 + E_4 + E_5 + E_7 + E_8.
 \end{aligned}$$

When $a^3 \neq 1$, computation yields two minimal classes of groups contributing to nonvanishing cohomology:

$$H^1(G', \text{Pic}(\tilde{X}_a)) = \mathbb{Z}/3,$$

for $G' = C_3 = \langle(1, 2, 3)\rangle$ or $\langle(4, 5, 6)\rangle$, realized as permutations of indices of the coordinates. When $a = 1$, these two classes of C_3 are conjugate in $\text{Aut}(X_1)$, and thus, we found a unique class of groups contributing to nonvanishing cohomology:

$$H^1(G', \text{Pic}(\tilde{X}_a)) = \mathbb{Z}/3,$$

for $G' = C_3 = \langle(1, 2, 3)\rangle$. Any subgroup of $\text{Aut}(X_a)$ containing those classes has **(H1)**-obstructions to stable linearizability.

Remark 9.1. One can characterize geometrically the C_3 -action contributing to **(H1)**-obstructions as follows: let X_a be a 9-nodal cubic threefold, and $G = C_3 \subseteq \text{Aut}(X_a)$. Then the G -action on X_a does not satisfy **(H1)** if and only if there exists a G -orbit of planes of length 3, which forms a Cartier divisor.

Excluding $G \subseteq \text{Aut}(X_a)$ with **(H1)**-obstruction or with G -fixed singular points, one is left with

- o When $a^3 = 1$, the unobstructed groups are

$$\mathcal{D}_6, \mathfrak{S}_3, \mathfrak{S}'_3, C_6, C_3, \tag{26}$$

where \mathcal{D}_6 acts on $\{x_1, x_2, x_3\}$ and $\{x_4, x_5, x_6\}$ via diagonal \mathfrak{S}_3 permutations and C_2 swapping them (i.e., $x_i \leftrightarrow x_{3+i}, i = 1, 2, 3$). The other groups are all subgroups of \mathcal{D}_6 .

- o When $a^3 \neq 1$, we are left with

$$\mathfrak{S}_3 \quad \text{and} \quad C_3,$$

where \mathfrak{S}_3 is the diagonal permutation and C_3 its subgroup.

Next, we show that the actions of these unobstructed groups on X_a are equivariantly birational to actions on a smooth quadric threefold. In particular, the actions of cyclic groups C_3 and C_6 are linearizable.

Linearization

Consider the family of degree (1, 1) divisors in $(\mathbb{P}^2)^2$

$$W_b \subset \mathbb{P}^2_{t_1, t_2, t_3} \times \mathbb{P}^2_{z_1, z_2, z_3}, \quad b \in \mathbb{C} \setminus \{0, -1, \zeta_3, \zeta_3^2\},$$

given by

$$(-t_1 z_2 + t_2 z_1) + b(-t_1 z_2 + t_3 z_3) = 0,$$

with a $G = \mathfrak{S}_3$ -action generated by

$$\iota : t_1 \leftrightarrow z_2, \quad t_2 \leftrightarrow z_1, \quad t_3 \leftrightarrow z_3$$

and

$$\sigma : t_1 \mapsto \zeta t_1, \quad t_2 \mapsto \zeta^2 t_2, \quad z_1 \mapsto \zeta z_1, \quad z_2 \mapsto \zeta^2 z_2.$$

Let $p_1, p_2, p_3 \in W_b$ be the points

$$[1 : 1 : 1] \times [1 : 1 : 1], \quad [\zeta : \zeta^2 : 1] \times [\zeta : \zeta^2 : 1], \quad [\zeta^2 : \zeta : 1] \times [\zeta^2 : \zeta : 1],$$

where $\zeta = e^{\frac{2\pi i}{3}}$. Note that $\{p_1, p_2, p_3\}$ forms one G -orbit. The linear system

$$|H - p_1 - p_2 - p_3|$$

consisting of hyperplanes on \mathbb{P}^4 containing points p_1, p_2 and p_3 has projective dimension 4. Under a chosen basis, it gives a birational map to a 9-nodal cubic hypersurface $Y_b \subset \mathbb{P}^4$, with equation

$$y_1 y_2 y_3 + y_1 y_5^2 - y_2^2 y_4 + y_2 y_4^2 - y_1^2 y_5 - \frac{b}{b+1} y_1 y_3 y_4 + \frac{b}{(b+1)^2} y_3^3 - \frac{b}{b+1} y_2 y_3 y_5 - \frac{1}{b+1} y_3 y_4 y_5 = 0.$$

Up to a change of variables by

$$(y_1 \ y_2 \ y_3 \ y_4 \ y_5) \cdot \begin{pmatrix} 1 & \zeta & \zeta^2 & \frac{-\zeta b + \zeta^2}{b - \zeta^2} & \frac{-\zeta^2 b + 1}{b - \zeta^2} \\ 1 & \zeta^2 & \zeta & \frac{-b + \zeta}{b - \zeta^2} & \frac{-\zeta^2 b + 1}{b - \zeta^2} \\ \frac{\zeta b - \zeta^2}{b+1} & \frac{\zeta b - \zeta^2}{b+1} & \frac{\zeta b - \zeta^2}{b+1} & \frac{-\zeta b + \zeta^2}{b+1} & \frac{-\zeta b + \zeta^2}{b+1} \\ \zeta & 1 & \zeta^2 & \frac{-\zeta^2 b + 1}{b - \zeta^2} & \frac{-\zeta b + \zeta^2}{b - \zeta^2} \\ \zeta & \zeta^2 & 1 & \frac{-b + \zeta}{b - \zeta^2} & \frac{-\zeta b + \zeta^2}{b - \zeta^2} \end{pmatrix},$$

Y_b is G -isomorphic to

$$\{y_1 y_2 y_3 + \lambda_b y_4 y_5 (y_1 + y_2 + y_3 + y_4 + y_5) = 0\} \subset \mathbb{P}^4, \quad \lambda_b = -\left(\frac{b - \zeta^2}{b - \zeta}\right)^3;$$

that is,

$$X_a = \{x_1 x_2 x_3 - x_4 x_5 x_6 = a(x_1 + x_2 + x_3) + x_4 + x_5 + x_6 = 0\} \subset \mathbb{P}^5,$$

where

$$a = -\frac{b - \zeta^2}{b - \zeta}.$$

The G -action on X_a is given by the diagonal permutation of coordinates x_1, x_2, x_3 and x_4, x_5, x_6 . When $b \neq 0, -1, \zeta, \zeta^2$, i.e., $a^3 \neq -1, 0$, one sees that W_b (and thus X_a) is G -equivariantly birational to

$$Q_b = \{(b+1)t_1 z_2 - t_2 z_1 + b z^2 = 0\} \subset \mathbb{P}^4_{t_1, t_2, z_1, z_2, z^2}, \tag{27}$$

realized as the equivariant compactification of the affine chart of W_b given by

$$\{t_3 \neq 0, z_3 \neq 0\} \subset W_b,$$

with the natural action of ι and σ (acting trivially on z).

When $a^3 = 1$ (i.e. $b = 1, -2$ or $-\frac{1}{2}$), there is extra symmetry on W_b and Q_b . For example, when $b = -2$, W_b and Q_b are invariant under the additional involution

$$\tau : t_1 \leftrightarrow t_2, \quad z_1 \leftrightarrow z_2.$$

The group $G' = \langle \iota, \sigma, \tau \rangle$ is isomorphic to \mathfrak{D}_6 . The corresponding G' -action on X_1 is generated by the diagonal \mathfrak{S}_3 -permutation and by swapping two sets of coordinates $\{x_1, x_2, x_3\}$ and $\{x_4, x_5, x_6\}$. We do not know whether or not this action is linearizable.

Corollary 9.2. *Let X_a be a 9-nodal cubic threefold as above. The C_3 -action on X_a via permutation of coordinates*

$$(x_1, x_2, x_3, x_4, x_5, x_6) \mapsto (x_3, x_1, x_2, x_6, x_4, x_5)$$

is linearizable for all $a^3 \neq 0, 1$. When $a = 1$, the C_6 -action on X_1 via

$$(x_1, x_2, x_3, x_4, x_5, x_6) \mapsto (x_6, x_4, x_5, x_3, x_1, x_2).$$

is linearizable.

Proof. By constructions above, these actions are equivariantly birational to actions on the corresponding smooth quadric Q_b , necessarily with fixed points. Projection from a fixed point on Q_b gives linearizations. □

Birational rigidity

Let X be the 9-nodal cubic threefold in $\mathbb{P}^4 \subset \mathbb{P}^5$ given by

$$x_1x_2x_3 - x_4x_5x_6 = x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 0,$$

and let $G = \text{Aut}(X) = \mathfrak{S}_3^2 \rtimes C_2$. We claim that X is G -birationally super-rigid. We start with several preliminary results.

Lemma 9.3. *If Σ is a G -orbit in X of length < 12 , then $|\Sigma| \in \{6, 9\}$.*

Proof. Left to the reader. □

Set

$$S = \{x_1 + x_2 + x_3 - x_4 - x_5 - x_6\} \cap X.$$

Then S is the unique G -invariant hyperplane section of X . Moreover, the cubic surface S is smooth, and G acts faithfully on it. This implies that S is isomorphic to the Fermat cubic surface [35, 36]. Consider

$$\alpha_G(S) = \sup \left\{ \lambda \in \mathbb{Q} \left| \begin{array}{l} \text{the pair } (S, \lambda D) \text{ is log canonical for every} \\ \text{effective } G\text{-invariant } \mathbb{Q}\text{-divisor } D \sim_{\mathbb{Q}} -K_S \end{array} \right. \right\}.$$

Lemma 9.4 (cf. [16, 25]). *One has $\alpha_G(S) = 2$.*

Proof. One can check that $\text{Pic}^G(S) = \mathbb{Z}[-K_S]$. Note that the group G is missed in [36, Theorem 6.14]. Note also that the linear system $| -K_S |$ does not contain G -invariant divisors, but $| -2K_S |$ contains a G -invariant divisor. Applying Lemma 9.3 and [16, Lemma 5.1], we obtain $\alpha_G(S) = 2$. \square

Lemma 9.5. *Let $C \subset X$ be a G -irreducible curve of degree < 12 . Then $C \subset S$.*

Proof. Assume $C \not\subset S$. Set $d = \text{deg}(C)$. Intersecting C with S , we immediately obtain $d = 6$ or $d = 9$, by Lemma 9.3. Moreover, we also see that

$$|S \cap C| = d,$$

so that C is smooth at every point in $S \cap C$, and S intersects C transversally. Hence, if C is irreducible and $C \not\subset S$, then G acts faithfully on C , which implies that the stabilizer of any point in $C \cap S$ is cyclic, which is impossible, since G does not have cyclic subgroups of index 6 and 9.

To complete the proof, we may assume that C is reducible. Let r be the number of its irreducible components. Write

$$C = C_1 + \dots + C_r,$$

where each C_i is an irreducible component of C . Set $d_1 = \text{deg}(C_1)$, and let H_1 be the stabilizer of the component C_1 in G . Then $d = d_1 r$, and since G does not have subgroups of index 3, we have one of the following cases:

- (1) $d = 9, r = 9, d_1 = 1, H_1 \cong \mathfrak{D}_4$,
- (2) $d = 6, r = 6, d_1 = 1, H_1 \cong \mathfrak{D}_6$,
- (3) $d = 6, r = 2, d_1 = 3, H_1 \cong \mathfrak{S}_3^2$ or $H_1 \cong \mathfrak{S}_3 \rtimes C_4$.

We exclude these cases one by one. In Case (1), there is a unique class of subgroups isomorphic to \mathfrak{D}_4 , and the \mathfrak{D}_4 -linear representation decomposes as

$$\mathbb{P}(\mathbb{I} \oplus \chi^2 \oplus V),$$

that is, a sum of the trivial representation \mathbb{I} , two copies of a nontrivial 1-dimensional subrepresentation χ , and an irreducible 2-dimensional representation V . By Schur’s lemma, V is the unique irreducible 2-dimensional representation in the ambient space of X . The projectivization $\mathbb{P}(V)$ defines an invariant line contained in S ,

$$l = \{x_1 + x_2 = x_3 = x_5 = 0\} \subset S.$$

The plane $\mathbb{P}(\mathbb{I} \oplus \chi^2) \subset \mathbb{P}^4$ intersects X along an irreducible cubic curve and contains no line. It follows that l is the only H_1 -invariant line in X , and thus, $C \subset S$.

In Case (2), there are two classes of subgroups isomorphic to \mathfrak{D}_6 . In one class, the \mathfrak{D}_6 -linear representation is

$$\mathbb{P}(\mathbb{I}^2 \oplus \chi \oplus V),$$

that is, the sum of two copies of the trivial 1-dimensional representation \mathbb{I} , a nontrivial 1-dimensional representation χ , and an irreducible 2-dimensional representation V . Again, V is the unique irreducible 2-dimensional representation. But in this case, the line $\mathbb{P}(V)$ is not contained in X . And the plane $\mathbb{P}(\mathbb{I}^2 \oplus \chi)$ intersects X along an irreducible cubic curve. Therefore, there is no H_1 -invariant line. The other class of \mathfrak{D}_6 decomposes as representation as

$$\mathbb{P}(\chi \oplus V_1 \oplus V_2),$$

that is, the sum of a nontrivial 1-dimensional representation χ and two nonisomorphic irreducible 2-dimensional representations V_1 and V_2 . Here, $\mathbb{P}(V_1)$ defines a line contained in S :

$$l = \{x_1 - x_4 = x_2 - x_4 = x_3 + x_4 + x_5 = 0\} \subset S,$$

while $\mathbb{P}(V_2)$ is not contained in X . In this case, we also have $C \subset S$.

In Case (3), suppose that $d = 6, r = 2, d_1 = 3$. Then the hyperplane

$$\{x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 0\}$$

is the unique H_1 -invariant hyperplane, and every H_1 -invariant plane in \mathbb{P}^4 is contained in this hyperplane. This implies that $C \subset S$. □

Theorem 9.6. *The Fano threefold X is G -birationally super-rigid.*

Proof. Suppose that X is not G -birationally super-rigid. Then it follows from the equivariant version of the Noether–Fano inequality [23] that there exists a G -invariant nonempty mobile linear system \mathcal{M} on X such that the singularities of the log pair $(X, \lambda\mathcal{M})$ are not canonical for $\lambda \in \mathbb{Q}_{>0}$ such that $\lambda\mathcal{M} \sim_{\mathbb{Q}} -K_X$. We seek a contradiction.

First, we claim that the singularities of the log pair $(X, \lambda\mathcal{M})$ are canonical away from finitely many points. Indeed, if this is not the case, then there exists a G -irreducible curve $C \subset X$ such that

$$\text{mult}_C(\mathcal{M}) > \frac{1}{\lambda},$$

which immediately implies that the degree of C is less than 12, which implies that $C \subset S$ by Lemma 9.5, so that the log pair $(S, \lambda\mathcal{M}|_S)$ is not log canonical, which contradicts Lemma 9.4, since $\lambda\mathcal{M}|_S \sim_{\mathbb{Q}} -2K_S$.

Next, we claim that the log pair $(X, \lambda\mathcal{M})$ is canonical at every singular point of X . Indeed, let $f: \tilde{X} \rightarrow X$ be the blow up of all singular points of X , let E_1, \dots, E_9 be the f -exceptional surfaces, let $\tilde{\mathcal{M}}$ be the strict transform on \tilde{X} of the linear system \mathcal{M} , and let \tilde{M} be a general surface in $\tilde{\mathcal{M}}$. Then, since $\text{Sing}(X)$ forms one G -orbit, we have

$$\lambda\tilde{M} \sim_{\mathbb{Q}} f^*(-K_X) - a \sum_{i=1}^9 E_i,$$

for some integer $a > 1$, by [26, Theorem 1.7.20] or [32, Theorem 3.10]. Recall that X contains nine planes

$$\Pi_{i,j} = \{x_i = 0, x_{3+j} = 0\} \subset \mathbb{P}^4,$$

and each of them contains four singular points of X . Let Π be one of the planes, C_2 a general conic in Π that contains $\Pi \cap \text{Sing}(X)$, and \tilde{C}_2 its strict transform on \tilde{X} . Then $\tilde{C}_2 \not\subset \tilde{M}$, so that

$$\begin{aligned} 0 \leq \lambda\tilde{M} \cdot \tilde{C}_2 &= \left(f^*(-K_X) - a \sum_{i=1}^9 E_i \right) \cdot \tilde{C}_2 \\ &= 4 - a \sum_{i=1}^9 E_i \cdot \tilde{C}_2 = 4 - 4a < 0, \end{aligned}$$

which is absurd.

Let P be a point in X such that the log pair $(X, \lambda\mathcal{M})$ is not canonical at P . Then $(X, \lambda\mathcal{M})$ is canonical in a punctured neighborhood of P , and it follows from [21, Remark 3.6] that the log pair $(X, \frac{3\lambda}{2}\mathcal{M})$ is not log canonical at P . Arguing as in the proof of [21, Proposition 3.5], we obtain a contradiction. □

Acknowledgements. We are grateful to Anton Mellit for his help with computations in Proposition 3.2 and to Olivier Wittenberg for his comments.

Competing interest. The authors have no competing interest to declare.

Financial support. The first author was partially supported by the Leverhulme Trust grant RPG-2021-229. The second author was partially supported by NSF grant 2301983.

References

- [1] J. D. Achter, S. Casalaina-Martin and Ch. Vial, ‘On descending cohomology geometrically’, *Compos. Math.* **153**(7) (2017), 1446–1478.
- [2] D. Allcock, ‘The moduli space of cubic threefolds’, *J. Algebr. Geom.* **12**(2) (2003), 201–223.
- [3] C. Araujo, A.-M. Castravet, I. Cheltsov, K. Fujita, A.-S. Kaloghiros, J. Martínez-García, C. Shramov, H. Süß, and N. Viswanathan, *The Calabi Problem for Fano Threefolds* vol. 485 (London Mathematical Society Lecture Note Series) (Cambridge University Press, Cambridge, 2023).
- [4] A. Avilov, ‘Automorphisms of singular cubic threefolds and the Cremona group’, *Math. Notes* **100**(3) (2016) 482–485.
- [5] A. Avilov, ‘Automorphisms of singular three-dimensional cubic hypersurfaces’, *Eur. J. Math.* **4**(3) (2018), 761–777.
- [6] A. Avilov, ‘On the forms of the Segre cubic’, *Mat. Zametki* **107**(1) (2020), 3–10.
- [7] L. Bayle and A. Beauville, ‘Birational involutions of \mathbf{P}^2 ’, *Asian J. Math.* **4**(1) (2000), 11–17. Kodaira’s issue.
- [8] A. Beauville, ‘Non-rationality of the symmetric sextic Fano threefold’, in *Geometry and Arithmetic* (European Mathematical Society (EMS), Zürich, 2012), 57–60. Based on the conference, Island of Schiermonnikoog, Netherlands, September 2010.
- [9] A. Beauville, ‘Non-rationality of the \mathfrak{S}_6 -symmetric quartic threefolds’, *Rend. Semin. Mat., Univ. Politec. Torino* **71**(3–4) (2013), 385–388.
- [10] O. Benoist and O. Wittenberg, ‘The Clemens–Griffiths method over non-closed fields’, *Algebr. Geom.* **7**(6) (2020), 696–721.
- [11] O. Benoist and O. Wittenberg, ‘Intermediate Jacobians and rationality over arbitrary fields’, *Ann. Sci. Éc. Norm. Supér. (4)* **56**(4) (2023), 1029–1084.
- [12] J. Blanc, ‘Elements and cyclic subgroups of finite order of the Cremona group’, *Comment. Math. Helv.* **86**(2) (2011), 469–497.
- [13] J. Blanc, I. Cheltsov, A. Duncan and Yu. Prokhorov, ‘Finite quasisimple groups acting on rationally connected threefolds’, *Math. Proc. Camb. Philos. Soc.* **174**(3) (2023), 531–568.
- [14] F. Bogomolov and Yu. Prokhorov. ‘On stable conjugacy of finite subgroups of the plane Cremona group, I’, *Cent. Eur. J. Math.* **11**(12) (2013), 2099–2105.
- [15] S. Casalaina-Martin, S. Grushevsky, K. Hulek and R. Laza, ‘Complete moduli of cubic threefolds and their intermediate Jacobians’, *Proc. Lond. Math. Soc. (3)* **122**(2) (2021), 259–316.
- [16] I. Cheltsov, ‘Log canonical thresholds of del Pezzo surfaces’, *Geom. Funct. Anal.* **18**(4) (2008), 1118–1144.
- [17] I. Cheltsov, M. Fedorchuk, K. Fujita, A.-S. Kaloghiros, K-moduli of pure states of four qubits, Preprint, 2024, [arXiv:2412.19972](https://arxiv.org/abs/2412.19972).
- [18] I. Cheltsov and J. Park, ‘Sextic double solids’, in *Cohomological and Geometric Approaches to Rationality Problems* (Progr. Math.) vol. 282 (Birkhäuser Boston, Boston, MA, 2010), 75–132.
- [19] I. Cheltsov and P. Pokora, ‘On K-stability of \mathbf{P}^3 blown up along a quintic elliptic curve’, *Ann. Univ. Ferrara Sez. VII Sci. Mat.* **70**(3) (2024), 1165–1192.
- [20] I. Cheltsov and A. Sarikyan, ‘Equivariant pliability of the projective space’, *Selecta Math. (N.S.)* **29**(5) (2023), Paper No. 71, 84.
- [21] I. Cheltsov, A. Sarikyan, and Z. Zhuang. ‘Birational rigidity and alpha invariants of Fano varieties’, Preprint, 2023, [arXiv:2304.11333](https://arxiv.org/abs/2304.11333).
- [22] I. Cheltsov and C. Shramov, *Cremona Groups and the Icosahedron* (Monographs and Research Notes in Mathematics) (CRC Press, Boca Raton, FL, 2016).
- [23] I. Cheltsov and C. Shramov, *Cremona Groups and the Icosahedron* (Monogr. Res. Notes Math.) (CRC Press, Boca Raton, FL, 2016).
- [24] I. Cheltsov, Yu. Tschinkel, and Zh. Zhang, ‘Equivariant geometry of the Segre cubic and the Burkhardt quartic’, 2023, To appear in *Selecta Math. (N.S.)*.
- [25] I. Cheltsov and A. Wilson, ‘Del Pezzo surfaces with many symmetries’, *J. Geom. Anal.* **23**(3) (2013), 1257–1289.
- [26] I. A. Cheltsov, ‘Birationally rigid Fano varieties’, *Russ. Math. Surv.* **60**(5) (2005), 875–965.
- [27] C. H. Clemens and P. A. Griffiths, ‘The intermediate Jacobian of the cubic threefold’, *Ann. Math. (2)* **95** (1972), 281–356.
- [28] J.-L. Colliot-Thélène and A. Pirutka, ‘Three-dimensional quartic threefolds: stable non-rationality’, *Ann. Sci. Éc. Norm. Supér. (4)* **49**(2) (2016), 371–397.
- [29] J.-L. Colliot-Thélène, J.-J. Sansuc and P. Swinnerton-Dyer, ‘Intersections of two quadrics and Châtelet surfaces. I’, *J. Reine Angew. Math.* **373** (1987), 37–107.
- [30] D. F. Coray, D. J. Lewis, N. I. Shepherd-Barron and P. Swinnerton-Dyer, ‘Cubic threefolds with six double points’, in *Number Theory in Progress, Vol. 1 (Zakopane-Kościelisko, 1997)* (de Gruyter, Berlin, 1999), 63–74.

- [31] A. Corti, 'Factoring birational maps of threefolds after Sarkisov. Appendix: Surfaces over nonclosed fields', *J. Algebr. Geom.* **4**(2) (1995), 223–254, appendix 248–254.
- [32] A. Corti, 'Singularities of linear systems and 3-fold birational geometry', in *Explicit Birational Geometry of 3-Folds* (Cambridge University Press, Cambridge, 2000), 259–312.
- [33] S. Cynk, 'Defect of a nodal hypersurface', *Manuscripta Math.* **104**(3) (2001), 325–331.
- [34] T. de Fernex, 'On planar Cremona maps of prime order', *Nagoya Math. J.* **174** (2004), 1–28.
- [35] I. V. Dolgachev, *Classical Algebraic Geometry. A Modern View* (Cambridge University Press, Cambridge, 2012).
- [36] I. V. Dolgachev and V. A. Iskovskikh, 'Finite subgroups of the plane Cremona group', in *Algebra, Arithmetic, and Geometry: In Honor of Yu. I. Manin. Vol. I* (Progr. Math.) vol. 269 (Birkhäuser Boston, Boston, MA, 2009), 443–548.
- [37] H. Finkelberg and J. Werner, 'Small resolutions of nodal cubic threefolds', *Nederl. Akad. Wetensch. Indag. Math.* **51**(2) (1989), 185–198.
- [38] V. González-Aguilera, A. Liendo and P. Montero, 'On the liftability of the automorphism group of smooth hypersurfaces of the projective space', *Israel J. Math.* **255**(1) (2023), 283–310.
- [39] Chr. D. Hacon and J. McKernan, 'The Sarkisov program', *J. Algebr. Geom.* **22**(2) (2013), 389–405.
- [40] B. Hassett, A. Kresch and Yu. Tschinkel, 'Stable rationality and conic bundles', *Math. Ann.* **365**(3–4) (2016), 1201–1217.
- [41] B. Hassett and Yu. Tschinkel, 'Flops on holomorphic symplectic fourfolds and determinantal cubic hypersurfaces', *J. Inst. Math. Jussieu* **9**(1) (2010), 125–153.
- [42] B. Hassett and Yu. Tschinkel, 'Equivariant geometry of odd-dimensional complete intersections of two quadrics', *Pure Appl. Math. Q.* **18**(4) (2022), 1555–1597.
- [43] B. Hassett and Yu. Tschinkel, 'Torsors and stable equivariant birational geometry', *Nagoya Math. J.* **250** (2023), 275–297.
- [44] P. Jahnke, Th. Peternell and I. Radloff, 'Threefolds with big and nef anticanonical bundles. II', *Cent. Eur. J. Math.* **9**(3) (2011), 449–488.
- [45] A. Kresch and Yu. Tschinkel, 'Cohomology of finite subgroups of the plane Cremona group', 2022, To appear in *Algebraic Geometry and Physics*.
- [46] A. Kresch and Yu. Tschinkel, 'Equivariant birational types and Burnside volume', *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **23**(2) (2022), 1013–1052.
- [47] A. Kuribayashi and K. Komiya, 'On Weierstrass points and automorphisms of curves of genus three', in *Algebraic Geometry (Proc. Summer Meeting, Univ. Copenhagen, Copenhagen, 1978)* (Lecture Notes in Math.) vol. 732 (Springer, Berlin, 1979), 253–299.
- [48] A. Kuznetsov and Yu. Prokhorov, 'Rationality over nonclosed fields of Fano threefolds with higher geometric Picard rank', *J. Inst. Math. Jussieu* **23**(1) (2024), 207–247.
- [49] K. Lauter, 'Geometric methods for improving the upper bounds on the number of rational points on algebraic curves over finite fields. Appendice by J. -P. Serre', *J. Algebr. Geom.* **10**(1) (2001), 19–36.
- [50] A. Pinaridin, A. Sarikyan, E. Yasinsky, 'Linearization problem for finite subgroups of the plane Cremona group', [arXiv:2412.12022](https://arxiv.org/abs/2412.12022).
- [51] Z. Reichstein and B. Youssin, 'Essential dimensions of algebraic groups and a resolution theorem for G -varieties', *Canad. J. Math.* **52**(5) (2000), 1018–1056. With an appendix by János Kollár and Endre Szabó.
- [52] N. I. Shepherd-Barron, 'The Hasse principle for 9-nodal cubic threefolds', Preprint, 2012, [arXiv:1210.5178](https://arxiv.org/abs/1210.5178).
- [53] Y. Tschinkel, K. Yang and Z. Zhang, 'Equivariant birational geometry of linear actions', *EMS Surv. Math. Sci.* **11**(2) (2024), 235–276.
- [54] N. Tziolas, 'Terminal 3-fold divisorial contractions of a surface to a curve. I', *Compos. Math.* **139**(3) (2003), 239–261.
- [55] S. Viktorova, 'On the classification of singular cubic threefolds', Preprint, 2023, [arXiv:2304.10452](https://arxiv.org/abs/2304.10452).
- [56] C. Voisin, 'Unirational threefolds with no universal codimension 2 cycle', *Invent. Math.* **201**(1) (2015), 207–237.
- [57] L. Wei and X. Yu, 'Automorphism groups of smooth cubic threefolds', *J. Math. Soc. Japan* **72**(4) (2020), 1327–1343.