

CrossMark

The 13th European Nutrition Conference, FENS 2019, was held at the Dublin Convention Centre, 15–18 October 2019

Intestinal permeability modulation through a polyphenol-rich dietary pattern in older subjects: MaPLE project outcomes and perspectives

Stefano Bernardi¹, Cristian Del Bo^{'1}, Simone Guglielmetti¹, Giorgio Gargari¹, Antonio Cherubini², Paul Kroon³, Benjamin Kirkup³, Nicole Hidalgo Liberona^{4,5}, rio Peron^{4,5}, Raul Gonzalez Dominguez^{4,5}, Cristina Andres-Lacueva^{4,5} and Patrizia Riso¹ Gregorio Peron⁴,

¹Department of Food, Environmental and Nutritional Sciences (DeFENS), Università degli Studi di Milano, Milan,

Italv.

²IRCCS INRCA, Geriatria, Accettazione Geriatrica e Centro di ricerca per l'invecchiamento, Ancona, Italy, ³Ouadram Institute Bioscience, Norwich Research Park, Norwich, United Kingdom,

⁴Biomarkers and Nutrimetabolomics Laboratory; Department of Nutrition, Food Sciences and Gastronomy;

Food Technology Reference Net (XaRTA); Nutrition and Food Safety Research Institute (INSA); Faculty of

Pharmacy and Food Sciences; University of Barcelona, Barcelona, Spain and

⁵CIBER Fragilidad y Envejecimiento Saludable (CIBERfes), Instituto de Salud Carlos III, Barcelona, Spain

Abstract

NS Proceedings of the Nutrition Society

In recent years, research has been focusing on strategies to counteract inflammatory processes and age-related diseases⁽¹⁾. During ageing, a low-grade systemic inflammation is often associated to an altered intestinal permeability (IP) a condition that has been shown to promote inflammation possibly through the translocation of dietary and bacterial factors into the blood stream that activates the immune system⁽²⁾. In this regard, dietary pattern and environmental factors could play a fundamental role because of their potential ability to modulate inflammation, IP and the gut microbial ecosystem (GME). Moreover, it has been hypothesized that bioactive compounds such as polyphenols may affect IP and GME⁽³⁾. The MaPLE project (Microbiome mAnipulation through Polyphenols for managing gut Leakiness in the Elderly) aimed to investigate the hypothesis that a polyphenol-rich diet can improve IP condition in a target population with beneficial changes at intestine and systemic level. To this aim, a randomised, controlled, crossover dietary intervention study (8-week polyphenol-rich diet versus 8-week control diet, separated by a wash-out period) was carried out in a group of older subjects (> 60 years) living in a well-controlled setting (i.e. nursing home). Markers related with IP, inflammation, oxidative stress, vascular function and intestinal microbial ecosystem were investigated in serum, urine and/or fecal samples. Moreover, blood bacteria DNAemia, and serum/urine metabolomics has been assessed. Moreover, a consistent nutritional evaluation of the standard menu (provided by the nursing home) and of weighed food diaries was performed, providing also data on actual polyphenol intake during the intervention. The results show there were higher levels of IP in the older subjects, and that the polyphenol-enriched diet changed the levels of serum zonulin, a marker of IP. In addition, an association between zonulin and blood bacterial load was demonstrated. Ongoing in vitro and in vivo experiments are exploring the potential effects of different polyphenols on IP and the mechanisms involved. The MaPLE project will generate new data to improve the understanding on the role of polyphenols in the modulation of intestinal microbiome and its interactions with the host.

Funding

JPI-HDHL http://www.healthydietforhealthylife.eu/; Mipaaft(D.M.8245/7303/2016); MINECO (PCIN2015-238); BBSRC (BB/ R012512/1)

Conflict of Interest

There is no conflict of interest

References

- Franceschi et al. (2000).
- 2. Bischoff et al. (2014).
- Guerriero et al. (2018).