

Original Article

Cite this article: Talbot E, Bruggeman J, Hauton C, Widdicombe S (2019). Uncovering the environmental drivers of short-term temporal dynamics in an epibenthic community from the Western English Channel. *Journal of the Marine Biological Association of the United Kingdom* **99**, 1467–1479. <https://doi.org/10.1017/S0025315419000663>

Received: 28 February 2019

Revised: 21 June 2019

Accepted: 5 July 2019

First published online: 5 August 2019


Key words:

Benthic community; benthic epifauna; bottom-water temperature; natural variability; phytodetrital input; supply side ecology; time series; Western English Channel

Author for correspondence:

Elizabeth Talbot, E-mail: sat@pml.ac.uk

Uncovering the environmental drivers of short-term temporal dynamics in an epibenthic community from the Western English Channel

Elizabeth Talbot^{1,2} , Jorn Bruggeman¹, Chris Hauton²
and Stephen Widdicombe¹

¹Plymouth Marine Laboratory, Plymouth PL1 3DH, UK and ²School of Ocean and Earth Science, University of Southampton, National Oceanography Centre Southampton, Waterfront Campus, Southampton SO14 3ZE, UK

Abstract

Benthic communities, critical to the health and function of marine ecosystems, are under increasing pressure from anthropogenic impacts such as pollution, eutrophication and climate change. In order to refine predictions of likely future changes in benthic communities resulting from these impacts, we must first better constrain their responses to natural seasonality in environmental conditions. Epibenthic time series data (July 2008–May 2014) have been collected from Station L4, situated 7.25 nautical miles south of Plymouth in the Western English Channel. These data were analysed to establish patterns in community abundance, wet biomass and composition, and to link any observed patterns to environmental variables. A clear response to the input of organic material from phytoplankton blooms was detected, with sediment surface living deposit feeders showing an immediate increase in abundance, while predators and scavengers responded later, with an increase in biomass. We suggest that this response is a result of two factors. The low organic content of the L4 sediment results in food limitation of the community, and the mild winter/early spring bottom water temperatures allow the benthos to take immediate advantage of bloom sedimentation. An inter-annual change in community composition was also detected, as the community shifted from one dominated by the anomuran *Anapagurus laevis* to one dominated by the gastropod *Turitella communis*. This appeared to be related to a period of high larval recruitment for *T. communis* in 2013/2014, suggesting that changes in the recruitment success of one species can affect the structure of an entire community.

Introduction

Benthic epifaunal communities are under growing pressure from anthropogenic activities in coastal seas, the strength and range of which have increased significantly over the last century with the growth in global industrialization and urbanization (Daan *et al.*, 1996; Halpern *et al.*, 2008). As a result, impacts such as pollution, eutrophication and the effects of climate change are of major concern to conservationists and resource managers alike (Capasso *et al.*, 2010). In order to assess long-term changes in epibenthic communities as a result of these impacts, it is critical that we gain an improved understanding of the short-term temporal variability in the responses of community abundance, biomass and composition to environmental parameters (Reiss & Kröncke, 2004; Chikina *et al.*, 2014; Włodarska-Kowalczyk *et al.*, 2016).

While descriptions of the large-scale spatial distribution and structure of temperate, shelf-sea epibenthic communities in connection with environmental factors are relatively common, only a few studies have investigated the temporal variability of these communities. Many of those focused on the North Sea, and concluded that sea surface temperature (SST) is a dominant factor influencing the temporal variability of epibenthic communities, particularly in the shallow, well-mixed areas of the south-eastern North Sea (Reiss & Kröncke, 2004; Neumann *et al.*, 2008, 2009b), which are characterized by strong seasonal fluctuations in temperature (Neumann *et al.*, 2008). The influence of SST appears to be less dominant in the deeper, stratified areas of the northern North Sea however. Neumann *et al.* (2009a) found no correlation between overall epibenthic community structure and changing SST, although relationships were found between SST and the abundance and biomass of some individual species, in some cases with a one year lag.

Shallow-water communities are generally thought to have access to high quality, if temporally variable, food (Pearson & Rosenberg, 1986), and as a result, the influence of organic input on structuring the benthos may be secondary to other physical and biological factors (Quijón *et al.*, 2008). Again, there are comparatively few studies which focus on the responses of benthic epifauna to bloom sedimentation, but a number have investigated macro-infaunal community structures and responses to phytodetrital inputs. The trophic structure of North Sea macrofauna communities was found to reflect differences in the relative quality of organic matter received (Dauwe *et al.*, 1998; Wiekling & Kröncke, 2005), and between 55% and 84% of year to year variability in benthic infaunal abundance off the coast of Northumberland was explained by changes in primary production (Buchanan, 1993). A marked increase in macrofaunal abundance in the same area in the 1980s was attributed to increases in phytodetrital



input (Frid *et al.*, 1996), as were decadal-scale variations in taxonomic composition (Frid *et al.*, 2009a, 2009b; Clare *et al.*, 2017). Josefson *et al.* (1993) showed that the abundance, biomass and growth of macro-infaunal species were closely related to bloom sedimentation in the Skagerrak–Kattegat region, while macrofaunal deposit feeders were found to increase in abundance immediately following bloom sedimentation in the Western English Channel, while other trophic groups responded more slowly, primarily with an increase in biomass (Zhang *et al.*, 2015). However, not all studies found a clear response to organic input. Quijón *et al.* (2008) found that the effects of phytodetrital input were short-term, and were minor in comparison to the seasonal differences observed in the macrofaunal community, and studies of the infauna of the western Baltic (Graf *et al.*, 1982) and of the epifauna in the German Bight area of the North Sea (Reiss & Kröncke, 2004) failed to find any response to bloom sedimentation at all.

In this study, the seasonal and inter-annual variability of the epibenthic community at Station L4 in the Western English Channel was investigated from July 2008 until May 2014. Since little is known about the ecology and biology of the epibenthos in the Western English Channel, these data provide valuable information on the short-term variation of several epibenthic groups. The purpose of this study was to (1) describe the seasonal and inter-annual variability in diversity, abundance and biomass of the epibenthos at Station L4 and (2) to identify and discuss environmental drivers in accordance with faunal patterns.

Materials and methods

The L4 sampling station

Station L4 is part of the Western Channel Observatory (WCO, <http://www.westernchannelobservatory.org.uk>), and represents a long-term oceanographic and marine biodiversity time series,

having been routinely sampled since 1988. In 2008, a benthic series was added – an element often missing from other observatories (Smyth *et al.*, 2015). Situated in the Western English Channel, 7.25 nautical miles south of Plymouth, UK (50°15.00'N 4°13.02'W), it can be considered representative of a temperate coastal environment (Tait *et al.*, 2015). Water column depth is ~53 m, and the station is well mixed during the winter and stratified over the summer (Figure 1A). Bottom water temperature varies from ~8–16°C (Figure 1A). Chlorophyll-*a* concentration is at its lowest during winter, and higher during the characteristic spring and autumn phytoplankton blooms (Figure 1B, note that these are fluorescence measurements, a proxy for chlorophyll-*a*). Phytoplankton cells can sink very quickly at L4, with high chlorophyll-*a* and phytoplankton carbon values measured at the seabed within two weeks of the start of a bloom (Tait *et al.*, 2015; Zhang *et al.*, 2015).

Animal collection and processing

Using a 60 cm wide Naturalist dredge with a 22 mm mesh, three replicate samples were taken every other month during the period July 2008–May 2014 from Station L4. For each replicate the dredge was lowered to the seabed and then towed for ~2 min at a speed of ~0.3 knots. Total towed distance was calculated for each replicate using the formula:

$$\cos^{-1}(\cos \text{lat}_S \times \cos \text{lat}_E + \sin \text{lat}_S \times \sin \text{lat}_E \times \cos(\text{lon}_S - \text{lon}_E)) \times 6371$$

Where: lat_S is start latitude (in radians), lat_E is end latitude (in radians), lon_S is start longitude (in radians), lon_E is end longitude (in radians) and 6371 is the approximate radius of the Earth (in km).




Fig. 1. Monthly mean water temperature (A), between July 2008 and May 2014 and fluorescence (B) between Jan 2009 and May 2014 at station L4. Data were collected weekly (weather permitting) using a SeaBird SBE 19+CTD deployed from the RV ‘Plymouth Quest’. Dotted vertical lines denote periods of thermal stratification.

All organisms collected in the dredge were fixed in 10% formalin solution. Prior to analysis, samples were rinsed at 4 mm and epifaunal individuals carefully picked out. All individuals were identified to species level, wherever possible, using a stereo microscope (Leica M32 Kombistereio). Individuals within each of the identified taxonomic groups were counted, carefully blotted dry and weighed on a Sartorius R220D microbalance (± 0.01 mg, European Instruments). Standardized values for abundance and biomass for each time point were calculated by dividing the total abundance (or biomass) in three replicate samples by the total area covered by the dredge. Those species which were only ever represented by a single individual across the course of the time series ($N = 23$) were excluded from further analysis.

Ancillary data

During the study period (July 2008–May 2014), a suite of environmental and biological data were collected from L4 every week (weather permitting) from the RV 'Plymouth Quest'. Vertical profiles of temperature and fluorescence were measured using a SeaBird SBE 19+ CTD. Water samples for phytoplankton analysis were collected from a depth of 10 m using 10 L Niskin bottles attached to the CTD rosette, and zooplankton were collected in two vertical WP2 net hauls (mesh size = 200 μm , mouth aperture = 57 cm diameter) taken from the seabed to the surface (UNESCO, 1968).

Phytoplankton analysis

Paired water-bottle samples were preserved with 2% Lugol's iodine solution (Thronsdén, 1978) and 4% buffered formaldehyde. Between 10 and 100 ml of sample (depending on cell density) were settled for at least 48 h (Widdicombe *et al.*, 2010). Cell volumes were calculated according to the equations of Kovalá & Larrance (1966) and converted to carbon (pgC cell^{-1}) (Menden-Deuer & Lessard, 2000) and then expressed per unit volume of seawater (mgC m^{-3}).

Meroplankton analysis

Haul samples were preserved and stored in 5% formalin. Two subsamples were extracted using a Folsom splitter and a Stempel pipette, to identify large and small organisms separately, then counted and identified under a microscope. Abundances in the two hauls were averaged to reduce the variability related to the sampling, and counts were converted to individuals per m^3 (John *et al.*, 2001). Due to the difficulties in larval identification and because different analysts have worked on the data set over the years, meroplankton are only identified to major taxonomic groups. These groups are: Decapoda, Brachyura, Cirripedia, Bivalvia, Gastropoda, Echinodermata and Polychaeta. These groups provide an overall picture of the seasonal changes in the meroplankton assemblage at L4. For this study, all groups except Cirripedia were considered, because although Cirripede larvae can dominate the meroplankton at L4 (Highfield *et al.*, 2010), mature animals are rarely present in the epibenthic faunal samples.

Statistical analysis

All statistical analyses were conducted in R statistical software. Time series of epibenthic abundance and wet biomass per square metre between July 2008 and May 2014 were compiled. Missing data were interpolated using the 'zoo' package in R (Zeileis *et al.*, 2018). Data for each sampling month (January, March, May, July, September and November) were pooled across the whole time series and overall means of community, major phyla and dominant species

abundance and wet biomass were calculated to establish the structure of the community. Average individual body mass of the whole community and each phylum was calculated by dividing the overall mean wet biomass by the overall mean abundance for each sampling month. To establish whether responses to environmental drivers were more easily identifiable when considering functional groups rather than taxa, species were grouped into one of five feeding guilds (predator/scavenger, omnivore, surface-deposit feeder, subsurface-deposit feeder, suspension feeder). Information on polychaete feeding mode was retrieved from Jumars *et al.* (2015). Information on feeding mode for all other phyla was retrieved from the Marine Life Information Network's biological traits catalogue (MarLIN, 2006). Where a species exhibited more than one feeding method, it was classified by the preferred or most frequently documented method. While we appreciate that the 'fuzzy coding' method (Chevene *et al.*, 1994; Neumann & Kröncke, 2011), which uses positive scores to describe the affinity of species to trait categories, would reflect a wider range of ecological function than the method adopted here, the aim of the present study was to provide a broad overview of the structure of the community and its responses to environmental variables, rather than an in-depth analysis of biological traits. Data on meroplanktonic larval abundance, water temperature and phytoplankton carbon for the duration of the time series were also collected and monthly means calculated.

The 'vegan' package in R (Oksanen *et al.*, 2017) was used to construct a species accumulation curve to determine whether the full diversity of Station L4 had been captured, and to calculate rarefied species richness as an indicator of changes in diversity over the course of the time series. Rarefied species richness was chosen as the measure of diversity as the total area covered by the dredge varied between sampling dates. As a consequence, it is likely that the number of species (and individuals) collected in each sample was a function of the area sampled. Rarefaction techniques can correct for this difference in sampling effort by generating the expected number of species in a small collection of n individuals drawn at random from a larger pool of N individuals (Gotelli & Colwell, 2001).

In order to identify any correlation between the biological (community, feeding guild and phyla abundance and biomass) and the environmental (water temperature, phytoplankton carbon) data series, cross correlation analysis was performed (Olden & Neff, 2001; Probst *et al.*, 2012) in R using the 'astsa' package (Stoffer, 2016). Where relationships between data series were found, linear regressions were used to quantify the relationship for a particular time lag. Cross correlation analysis was also performed on the phyla abundance and larval abundance data series. Prior to this analysis, all data series were checked for homoscedasticity by fitting a simple ordinary least squares regression model and checking the residuals against the fitted values. The community abundance and biomass series, and the larval abundance data series were found to be heteroscedastic and were log- n transformed to achieve homoscedasticity. All data series were differenced to remove any trends or seasonal effects (Probst *et al.*, 2012), and then checked for stationarity using an Augmented Dickey–Fuller test in the 'tseries' package for R (Trapletti *et al.*, 2017). Finally, to ensure that estimates of cross correlations were not inflated by any temporal autocorrelation, series were checked for any autocorrelation by generating ACF and PACF plots (Olden & Neff, 2001).

Results

Variations in the epibenthic community 2008–2014

The total area covered by the dredge across the three replicate samples varied from 326.1 m^2 in March 2009 to 1170.3 m^2 in

September 2012. A total of 169 species were recorded over the course of the series, 143 of which were used for analysis. The number of species represented by only a single individual in each sample ranged from 4 (17% of total species recorded in the sample) in January 2012 to 23 (38% of total species recorded in the sample) in July 2009. The species accumulation curve for Station L4 started to level off after ~20 samples (corresponding to a sampling period of 2.5 years) had been collected (Figure 2A). Rarefied species richness varied across the course of the series (Figure 2B) with values ranging from 7–24 species. Spring/summer values were generally higher than values in the preceding winter. Rarefied richness reached a maximum in spring/summer 2011, and declined steadily to the end of the series.

Community abundance over the course of the series varied from 0.13 individuals m^{-2} in May 2010, to 3.93 individuals m^{-2} in May 2014 (Figure 3A). Community wet biomass ranged from 0.21 $g m^{-2}$ in September 2008 to 5.99 $g m^{-2}$ in May 2014 (Figure 3B). The peak in community abundance seen in summer 2009 was largely attributable to a peak in crustacean abundance (Figure 3E). The increase in abundance in autumn 2013 was driven by increases in crustaceans (Figure 3E), molluscs (Figure 3E) and to a lesser extent, echinoderms (Figure 3C). The abundance maximum in spring 2014 was due predominantly to an increase in mollusc numbers (Figure 3E). All but one of the observed peaks in wet biomass were driven by increases in echinoderm biomass (Figure 3F). The very high wet biomass maximum in spring 2014 can be attributed to increases in biomass of molluscs and crustaceans (Figure 3F). The increase in polychaete biomass in spring 2014 (Figure 3D) was due to the presence of a single large *Aphrodita aculeata* (Linnaeus 1758) in the sample. The species which contribute to these peaks in community abundance and biomass are listed in Table 1.

Benthic larvae were always present in the water column over the course of the time series. Abundances ranged from 12 (± 8) individuals m^{-3} in December 2009 to 2080 (± 3656) individuals m^{-3} in July 2010. In 2009, 2010 and 2011, summer abundances of benthic larvae were very high, reaching more than 1000 individuals m^{-3} (Figure 4A). In 2009, the majority of the benthic larvae recorded were gastropod molluscs (Figure 4C), while in 2010 and

2011, echinoderm larvae were the primary contributors to the observed peaks in abundance (Figure 4D).

Both the abundance and wet biomass of suspension and deposit feeders was relatively stable over the course of the time series, although suspension feeder abundance increased sharply in summer 2013 and spring 2014 (Figure 5A). Likewise, the wet biomass of suspension and deposit feeders reached a peak in spring 2014 (Figure 5B). In terms of abundance, omnivores were the most dominant feeding guild at L4 (Figure 5C), while the predator/scavenger guild dominated in terms of wet biomass (Figure 5F). The predators/scavengers (Figure 5E) exhibited an increase in abundance in spring 2014, although it was not as dramatic as that recorded for the suspension feeders. The number of taxa mapped into each feeding guild is shown in Table 2. The majority of taxa recorded were predators/scavengers, while sub-surface deposit feeders were represented by only a small number of taxa.

Overall structure of the epibenthic community

There was some structure apparent in the overall means of the community abundance and wet biomass series. Abundance increased from January to reach a peak in May, before declining again over the summer. There was a second, smaller increase apparent in September, and abundance then decreased steadily through the winter (Figure 6A). Wet biomass peaked in March, and remained fairly high over the spring before declining from May onward. There was a second, smaller increase in community wet biomass in September (Figure 6B). The first peak in abundance (Figure 6E) which, when added to an already high abundance of crustaceans (Figure 6I), raised community abundance to ~1.2 individuals m^{-2} . While mollusc abundance dropped sharply after the May maximum (Figure 6E), the decline in community abundance was more gradual, due to an increase in the abundance of crustaceans (Figure 6I), and numbers of polychaetes and echinoderms remaining relatively high (Figure 6C and G). This increase in crustacean abundance, which reached its maximum in September, was the primary contributor to the second community abundance peak (Figure 6A). The biomass maximum in




Fig. 2. Species accumulation curve (A) and rarefied species richness (B) over the course of the L4 epibenthic time series, July 2008–May 2014. The species accumulation curve started to level off after ~20 samples (representing a sampling period of 2.5 years) were collected. Grey shading denotes standard deviation from the mean curve, generated from 1000 random permutations of the data. In terms of diversity, the series is characterized by a period of high species richness in late 2010/2011, with periods of lower richness at the beginning and end of the series. Dotted vertical lines denote January of each year.




Fig. 3. Abundance of the whole L4 epibenthic community (A), polychaetes and echinoderms (C) and molluscs and crustaceans (E) over the study period July 2008–May 2014. Biomass of the whole community (B), polychaetes (D) and echinoderms, molluscs and crustaceans (F) over the same period. Dotted vertical lines denote January of each year.

March was predominantly caused by a sharp increase in biomass of echinoderms (Figure 6H), and polychaetes (Figure 6D). Both molluscs and crustaceans (Figure 6F and J) reached a biomass peak in May, ensuring that community biomass remained high throughout the spring (Figure 6B).

Average individual body mass of the whole community reached a maximum in March, and then decreased steadily over the summer, before increasing again from September (Figure 7A). Polychaete (Figure 7A) individual body mass exhibited a similar pattern, although the decline after March was much steeper. Peaks in body mass for crustaceans and molluscs (Figure 7A) were reached in May, again with a decrease over the summer, and an increase beginning in September. Echinoderms (Figure 7B) reached a maximum in March, and body mass remained high into May before declining. There was a subsequent increase in echinoderm body mass although it started earlier than in other taxa, in July.

Overall monthly means of larval abundance show that the highest numbers of benthic larvae are recorded in the water column in July/August (Figure 8A). While all four major phyla contribute to this peak in abundance, it is largely attributable to high abundances of molluscs (Figure 8C) and echinoderms (Figure 8D). Different classes of mollusc appear to have different

spawning times at L4, with gastropod larvae abundances (grey dotted line, Figure 8C) in the water column peaking slightly earlier than bivalve larvae (grey dashed line, Figure 8C). Polychaete larvae exhibited two peaks, in March and June (Figure 8B), while decapod larvae abundances peaked in March/April (Figure 8E), and remained relatively high throughout the summer, before declining steadily from July. Much of the initial peak in decapod larval abundance can be attributed to brachyuran larvae (grey dotted line, Figure 8E), although this declines after April.

Drivers of variation in epibenthic community structure

NMDS ordination revealed no clear seasonal pattern over the course of the time series in either the abundance or wet biomass data, but there were some inter-annual differences in the abundance data. Analysis of community abundance identified three clusters (Figure 9A). Cluster 1 consists of the years 2008, 2013 and 2014. Cluster 2 contains the years 2011 and 2012, and cluster 3 contains the years 2009 and 2010. The differences between these three clusters were driven by differences in the relative abundances of the two dominant species. In cluster 1, the gastropod *T. communis* was dominant, whereas in cluster 3 the anomuran crab *Anapagurus laevis* was dominant. Cluster 2 was characterized by

Table 1. Species contributing to the observed community and/or wet biomass peaks

Date	Peak in abundance/wet biomass?	Contributing species	Taxonomic authority
Summer 2009	Abundance	<i>Anapagurus laevis</i>	Bell, 1845
Autumn 2013	Abundance	<i>A. laevis</i>	–
		<i>Galathea nexa</i>	Embleton, 1834
		<i>Macropodia</i> spp.	–
		<i>Turitella communis</i>	Risso, 1826
		<i>Astropecten irregularis</i>	Pennant, 1777
Spring 2014	Abundance & biomass	<i>Ophiothrix fragilis</i>	Abildgaard, 1789
		<i>T. communis</i> (abundance & biomass)	–
		<i>Liocarcinus depurator</i> (biomass)	Linnaeus, 1758
March 2010	Biomass	<i>Atelecyclus rotundatus</i> (biomass)	Olivi, 1792
		<i>Marthasterias glacialis</i>	Linnaeus, 1758
		<i>Luidia ciliaris</i>	Philippi, 1837
September 2012	Biomass	<i>Marthasterias glacialis</i>	–

a more even community structure, with no single species dominant. There were no clear inter-annual patterns identified in the biomass data, with most data points falling into a single cluster (Figure 9B). The only months to fall outside this cluster were January 2010 and January 2011. This appears to be due to the fact that during these months, the asteroid *M. glacialis*, which dominated the biomass over the course of the time series, was not recorded.

All significant results of the cross correlation analysis are shown in Table 3. There was no significant relationship between total community abundance and any of the explanatory environmental variables. Community wet biomass correlated with both phytoplankton carbon and temperature, with a 6 month lag in both cases. If both carbon and temperature are left in the linear model as explanatory variables, the model fit improves and both terms remain significant. There was no significant interaction effect detected between carbon and temperature. Of the four major phyla, only echinoderm biomass exhibited significant relationships with any of the explanatory environmental variables, correlating with temperature with a 6 month lag. Both mollusc and echinoderm abundance were correlated with larval abundance. Mollusc abundance was correlated with total mollusc larval abundance and gastropod larval abundance with a 4.5 year lag in each case. Echinoderm abundance is correlated with echinoderm larval abundance with a lag of 3 years.

Analysis of the community when grouped into feeding guilds (suspension feeders, surface and sub-surface deposit feeders, omnivores and predators/scavengers) showed a relationship between predator/scavenger biomass and phytoplankton carbon with a 6 month lag (Table 3). As with community wet biomass,

this group also exhibited a relationship with temperature, again with a 6 month lag. If both of these terms are left in the model, they remain significant and the overall fit improves (Table 3). There was no significant interaction effect detected however. This analysis also showed a relationship between phytoplankton carbon and surface-deposit feeder abundance, with an immediate response from the surface-deposit feeders to phytodetrital input (Table 3).

Discussion

Time series data (collected every other month during the period July 2008–May 2014) for benthic epifauna at Station L4 were analysed to establish patterns in community abundance, wet biomass and composition, and to link any observed patterns to environmental variables. A clear response to the input of organic material from phytoplankton blooms was detected, with sediment surface living deposit feeders showing an immediate increase in abundance, while predators and scavengers responded later, with an increase in biomass. An inter-annual change in community composition was also detected, as the community shifted from one dominated by the anomuran *Anapagurus laevis* to one dominated by the gastropod *Turitella communis*.

There is some evidence for benthic-pelagic coupling at Station L4, demonstrated by the correlation between surface-deposit feeder abundance and phytoplankton carbon. This pattern has been previously observed in the macro-infauna at L4, with deposit feeders rapidly responding to phytodetrital input with an increase in abundance, while predators and scavengers responded more slowly with an increase in biomass (Zhang *et al.*, 2015). While many studies have concluded that benthic communities can be structured by phytodetrital input over both short-term and decadal scales (Buchanan, 1993; Josefson *et al.*, 1993; Dauwe *et al.*, 1998; Wiekling & Kröncke, 2005; Frid *et al.*, 2009a, 2009b; Clare *et al.*, 2017) clear responses to organic input from benthic fauna can be difficult to detect (Graf *et al.*, 1982; Reiss & Kröncke, 2004). The ‘food bank’ hypothesis suggests that large reserves of labile organic matter in sediments can sustain benthic communities at constant levels of abundance on a year round basis, and clear responses to phytodetrital input are difficult to detect as a consequence (Mincks *et al.*, 2005; Kędra *et al.*, 2012; Włodarska-Kowalczyk *et al.*, 2016). This appears not to be the case at Station L4, which is fairly impoverished in terms of organic matter content, with organic carbon contributing only 0.4% to total sediment mass (Zhang *et al.*, 2015). It is possible that this comparatively low sediment carbon content results in the epibenthic community at L4 being food limited, and so the seasonal pulses of phytodetrital input elicit measurable responses. Furthermore, spring bloom sedimentation in temperate areas can often occur when bottom water temperatures are low, and benthic faunal responses are limited as a result. Weeks can pass before water temperature increases enough to allow for macrofaunal feeding (Lopez & Levinton, 2011). It is possible that the particular hydrographic conditions in the Western English Channel, where bottom water temperatures fluctuate less than in other temperate systems, result in early spring temperatures high enough for the surface-deposit feeders in the L4 community to respond immediately.

Interestingly, there was no apparent decrease in diversity associated with the sedimentation of the spring bloom. In macrofaunal communities, enriched sediments are typically rapidly colonized by a few opportunist, fast-reproducing species (Widbom & Frithsen, 1995) which can take advantage of the fresh organic matter, generally resulting in a reduction in diversity (Chamberlain *et al.*, 2001; Widdicombe & Austen, 2001). As noted above, epibenthic surface-deposit feeders did show an




Fig. 4. Abundance of benthic larvae in the water column for (A) the four major benthic phyla, (B) polychaetes, (C) molluscs, (D) echinoderms and (E) decapod crustaceans during the study period July 2008–May 2014. The grey dotted line in panel (C) is the abundance of gastropod larvae present, while the grey dashed line is the abundance of bivalve larvae. The grey dotted line in panel (E) is the abundance of brachyuran larvae present. Grey shading represents standard deviation from the mean calculated for the phyla.

immediate increase in abundance with the arrival of phytodetritus, but rarefied richness values for May (post sedimentation) are generally equal to or higher than values for March (pre-bloom). It is possible that these values are indirect evidence of predation. Predation is thought to play a key role in marine sedimentary systems, due in part to the lack of clear evidence for competitive exclusion (Peterson, 1979; Woodin, 1999). While detection of predation is challenging, and numerous studies have found no consistent regulatory role (Thrush, 1999), it has been suggested that epibenthic predators can equalize numbers and increase evenness by preying preferentially on numerically dominant species (Quijón & Snelgrove, 2005). Given the fact that L4 community wet biomass is predominantly represented by predators and scavengers, there is a possibility that opportunistic deposit feeders are prevented from becoming dominant after sedimentation of the spring bloom by the feeding of the predator/scavenger group. This pattern in the regulation of benthic community structure has been noted before (Posey *et al.*, 1995), with those authors concluding that the presence or absence of predation may alter the visible response of the benthos to organic enrichment. While there was no direct evidence of predator–prey interactions (e.g. a clear relationship between deposit-feeder and predator/scavenger abundance or biomass, as defined by Lotka–Volterra type models) detected in this study, the patterns in

species richness observed would seem to support the proposal that epibenthic predators can be of major influence in benthic communities (Quijón & Snelgrove, 2005), and may diminish or counterbalance the changes in prey species that result from phytodetrital input.

A relationship between community wet biomass and both bottom-water temperature and phytoplankton carbon was detected at Station L4, although there was no significant interaction between the two predictors and their effects on biomass. This leads us to propose that temperature and phytoplankton carbon primarily influence biomass at different times of the year. Community wet biomass peaks in March/May, driven predominantly by an increase in biomass of echinoderms and molluscs. Individual body mass curves for these two phyla show an identical pattern, with a maximum also being reached in March/May. It is possible that this is representative of the development of the gonads in preparation for spawning. Several studies have found that ripe gonads in these two phyla can make a significant contribution to body mass (Barker & Nichols, 1983; Nichols & Barker, 1984a, 1984b; Berthelin *et al.*, 2000; Freeman *et al.*, 2001; Alunno-Bruscia *et al.*, 2011). This view would appear to be supported by the increase in benthic larvae (of which mollusc and echinoderm larvae are recorded in the highest numbers) in the water column from May onwards, while community, mollusc




Fig. 5. Abundance (left hand side of the panel) and wet biomass (right hand side of the panel) of the main feeding guilds found at L4. (A) and (B) are suspension and deposit feeders, (C) and (D) are omnivores and (E) and (F) are predators/scavengers. Deposit feeders were split into surface and sub-surface feeders for the purposes of analysis, but were combined for plotting.

and echinoderm biomass decreases after May, perhaps indicating spent individuals. We suggest that this pre-spawning biomass is influenced by temperature. Several studies have noted the role of temperature in triggering gonad development in marine invertebrate species (Sastry, 1966; Sastry & Blake, 1971; Aktaş *et al.*, 2003; Herrmann *et al.*, 2009; Balogh *et al.*, 2018), and it is possible that gonad development at L4 is initiated by the high water

Table 2. The number of taxa mapped into each feeding guild and their % contribution to overall species richness

Feeding guild	Number of taxa	% of total number of species recorded
Suspension feeders	39	27.2
Surface deposit feeders	19	13.3
Sub-surface deposit feeders	8	5.6
Omnivores	25	17.5
Predators/scavengers	52	36.4
Total	143	100

Sub-surface deposit feeders are likely to be under-represented in samples due to the sampling method.

temperatures recorded in September, with full maturation and spawning occurring the following spring. Gonad development and maturation in some temperate echinoderm and mollusc species has been recorded to take up to 6 months, which would be in keeping with the 6-month lag between peaks in temperature and biomass identified in this study (Bowner, 1982; Sköld & Gunnarsson, 1996; Kim *et al.*, 2016). Although there was no significant interaction between temperature and phytoplankton carbon and their effects on biomass detected in this study, food availability will clearly affect gonad development as it dictates the nutritional status of an individual (Nunes & Jangoux, 2004), and the autumn bloom characteristic of Station L4, along with the carbon from seaweed detritus which contributes to winter organic matter in the area (Queirós *et al.*, 2019) is likely to help fuel gonad development over the winter. In contrast to maximum temperatures, maximum phytoplankton carbon values are generally recorded in April/May, with a response in community biomass seen 6 months later. It is possible that the relationship between phytoplankton carbon and biomass is indicative of somatic growth, which occurs after spawning has taken place in the spring. The seasonal prioritization of either sexual or somatic growth in benthic fauna is well documented, particularly in echinoderms (Greenwood, 1980; Peterson & Fegley, 1986; Guillou & Michel, 1993; Lozano *et al.*, 1995; Coma *et al.*, 1998). This shift in energetic prioritization is often related to reproductive effort




Fig. 6. Overall means for community and major phyla abundance (left hand side of panel) and wet biomass (right hand side of panel), calculated from data for each sampling month (Jan, Mar, May, Jul, Sept, Nov) pooled across the whole time series. Shaded grey area represents standard deviation of the mean.




Fig. 7. Average individual body mass of the whole epibenthic community, molluscs, polychaetes and crustaceans (A), and echinoderms (B).

being concentrated at a time favourable to the survival of offspring, e.g. spawning prior to or coincident with a phytoplankton bloom (Giangrande *et al.*, 1994). The same lagged relationship between biomass, temperature and phytoplankton carbon was

also recorded in the predator/scavenger group. The biomass of this feeding guild is dominated by echinoderms (70%), so the postulated relationships outlined above could also be driving the responses of this group.




Fig. 8. Overall monthly mean abundances of benthic larvae in the water column for (A) the four major benthic phyla combined, (B) polychaetes, (C) molluscs, (D) echinoderms and (E) decapod crustaceans. Means were calculated for each month from data pooled across the whole time series. The grey dotted line in panel (C) is the abundance of gastropod larvae present, while the grey dashed line is the abundance of bivalve larvae. The grey dotted line in panel (E) is the abundance of brachyuran larvae present. Grey shading represents standard deviation from the mean calculated for the phyla.




Fig. 9. NMDS ordination of community abundance (A) and wet biomass (B) data over the course of the time series July 2008–May 2014. Although there is no seasonal pattern evident in either the abundance or wet biomass data, there is some inter-annual variation in the abundance data, predominantly due to the relative variations in abundance of the dominant species.

Table 3. Significant models identifying relationships between the epibenthic community and environmental variables. In those models where two predictors were included, the significance value for the whole model has been given in the column for R^2 , and the significance value for individual predictors has been given in the coefficients column

Response variable	Predictor	lag	R^2 (P)	Coefficient (P)
Community biomass	Phytoplankton carbon	6 months	0.4 (0.0001)	0.1
Community biomass	Temperature	6 months	0.43 (0.0001)	0.18
Community biomass	Carbon + temperature	6 months	0.6 (<0.0001)	Carbon = 0.1(0.003) Temperature = 0.14 (0.002)
Echinoderm biomass	Temperature	6 months	0.37 (0.0003)	0.11
Predator/scavenger biomass	Phytoplankton carbon	6 months	0.32 (0.0006)	0.1
Predator/scavenger biomass	Temperature	6 months	0.38 (0.0003)	0.13
Predator/scavenger biomass	Carbon + temperature	6 months	0.53 (<0.0001)	Carbon = 0.09 (0.02) Temperature = 0.12 (0.002)
Surface-deposit feeder abundance	Phytoplankton carbon	0 months	0.42 (0.02)	0.09
Mollusc abundance	Mollusc larvae	4.5 years	0.81 (0.001)	0.21
Mollusc abundance	Gastropod larvae	4.5 years	0.61 (0.01)	0.17
Echinoderm abundance	Echinoderm larvae	3 years	0.58 (0.002)	0.1

The role of larval supply as a determinant of the structure and dynamics of marine populations (i.e. supply side ecology) has long been discussed (Thorson, 1950; De Wolf, 1973; Lewin, 1986; Underwood & Fairweather, 1989), and there is much evidence to suggest that variations in recruitment can contribute to patterns of abundance and demographics in adult populations of fish (Williams, 1980; Doherty & Fowler, 1994), barnacles (Gaines & Roughgarden, 1985; Sutherland, 1990; Scrosati & Ellrich, 2017), mussels (Scrosati & Ellrich, 2017) and bryozoans (Hughes, 1990). We propose that larval recruitment of dominant species is also a key influence on benthic community structure and composition at Station L4. The dramatic increase in community and suspension feeder abundance and biomass in May 2014, and the shift in community structure (from one dominated by *Anapagurus laevis* in 2009 to one dominated by *T. communis* in 2013/2014) are likely due to the sieve recruitment (the point at which individuals recruited to the population reach a size where they would be retained on the sieve mesh) of the high numbers of gastropod larvae present in the plankton in 2009. Previous studies of benthic recruitment have stressed that sieve recruitment can be far removed in time from actual settlement (Buchanan & Moore, 1986), as many benthic macrofaunal settlers are of meio-faunal size. The lag of 4.5 years identified between mollusc abundance and gastropod larval abundance likely reflects the fact that any newly settled animal needs to reach a size both big enough to be collected by the dredge, and to be retained on the 4 mm sieve used in this study.

Analysis of the first six years of the epibenthic time series at Station L4 reveals some temporal structure in community abundance and wet biomass, apparently influenced by both bottom water temperature and seasonal phytodetrital input. We suggest that the spring phytoplankton bloom fuels somatic growth, while gonad development and maturation is triggered by warmer water temperature in the autumn, resulting in a pre-spawning biomass peak evident in early spring. Different functional groups within the community were found to respond to the bloom in specific ways, a result that is in keeping with previous studies of the L4 macro-benthos. While benthic faunal responses to changes in water temperatures have been previously recorded in other temperate systems, clear responses to phytodetrital input as seen here are less common. We suggest that the reason we can detect this response is a combination of two factors. (1) The relative

impoverishment of the L4 sediment in terms of organic content, indicating a food-limited community, and (2) the comparatively small range of bottom water temperatures, resulting in relatively mild winter/early spring conditions and a community that is able to take immediate advantage of bloom sedimentation.

Acknowledgements. We are grateful to the crew of the RV 'Plymouth Quest', Sarah Dashfield, Joana Nunes, Amanda Beesley and Louise McNeill for benthic sample collection; Adrien Lowenstein, Laura Briers, Bianca Maria Torre, Harry Powell, Alessandro Brunzini, Craig Dornan, Olli Ford and Jemma French for their contributions to the taxonomic analysis of the benthic epifaunal samples; Clare Widdicombe for providing the phytoplankton data, and Andrea McEvoy and Angus Atkinson for providing the mero-plankton data. We would also like to thank the two anonymous reviewers whose comments improved the manuscript considerably.

Financial support. This work was supported by the National Environment Research Council via the SPITFIRE studentship [E.T., grant number NE/L002531/1] and through its National Capability Long-term Single Centre Science Programme, Climate Linked Atlantic Sector Science [grant number NE/R015953/1], and is a contribution to Theme 1.3 – Biological Dynamics.

References

- Aktaş M, Kumlu M and Eroldogan O (2003) Off-season maturation and spawning of *Penaeus semisulcatus* by eyestalk ablation and/or temperature–photoperiod regimes. *Aquaculture* **228**, 361–370.
- Alunno-Bruscia M, Bourlès Y, Maurer D, Robert S, Mazurié J, Gangnery A, Gouletquer P and Pouvreau S (2011) A single bio-energetics growth and reproduction model for the oyster *Crassostrea gigas* in six Atlantic ecosystems. *Journal of Sea Research* **66**, 340–348.
- Balogh R, Wolfe K and Byrne M (2018) Gonad development and spawning of the vulnerable commercial sea cucumber, *Stichopus herrmanni*, in the southern Great Barrier Reef. *Journal of the Marine Biological Association of the United Kingdom*. doi: <https://doi.org/10.1017/S0025315418000061>
- Barker MF and Nichols D (1983) Reproduction, recruitment and juvenile ecology of the starfish, *Asterias rubens* and *Marthasterias glacialis*. *Journal of the Marine Biological Association of the United Kingdom* **63**, 745–765.
- Berthelin C, Kellner K and Mathieu M (2000) Storage metabolism in the Pacific oyster (*Crassostrea gigas*) in relation to summer mortalities and reproductive cycle (West Coast of France). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **125**, 359–369.
- Bowner T (1982) Reproduction in *Amphiura filiformis* (Echinodermata: Ophiuroidea): seasonality in gonad development. *Marine Biology* **69**, 281–290.

- Buchanan JB** (1993) Evidence of benthic pelagic coupling at a station off the Northumberland coast. *Journal of Experimental Marine Biology and Ecology* **172**, 1–10.
- Buchanan JB and Moore JJ** (1986) Long-term studies at a benthic station off the coast of Northumberland. *Hydrobiologia* **142**, 121–127.
- Capasso E, Jenkins SR, Frost M and Hinz H** (2010) Investigation of benthic community change over a century-wide scale in the western English Channel. *Journal of the Marine Biological Association of the United Kingdom* **90**, 1161–1172.
- Chamberlain J, Fernandes TF, Read P, Nickell TD and Davies IM** (2001) Impacts of biodeposits from suspended mussel (*Mytilus edulis* L.) culture on the surrounding surficial sediments. *ICES Journal of Marine Science* **58**, 411–416.
- Chevène F, Dolédec S and Chessel D** (1994) A fuzzy coding approach for the analysis of long-term ecological data. *Freshwater Biology* **31**, 295–309.
- Chikina MV, Spiridonov VA and Mardashova MV** (2014) Spatial and temporal variability of coastal benthic communities in the Keretsky Archipelago area and in the Velikaya Salma Strait (Karelian coast, the White Sea). *Oceanology* **54**, 54–65.
- Clare DS, Spencer M, Robinson LA and Frid CLJ** (2017) Explaining ecological shifts: the roles of temperature and primary production in the long-term dynamics of benthic faunal composition. *Oikos* **126**, 1123–1133.
- Coma R, Ribes M, Gili J-M and Zabala M** (1998) An energetic approach to the study of life-history traits of two modular colonial benthic invertebrates. *Marine Ecology Progress Series* **162**, 89–103.
- Daan N, Richardson K and Pope JG** (1996) Changes in the North Sea ecosystem and their causes: Aarhus 1975 revisited—Introduction. *ICES Journal of Marine Science* **53**, 879–883.
- Dauwe B, Herman PMJ and Heip HR** (1998) Community structure and bio-turbation potential of macrofauna at four North Sea stations with contrasting food supply. *Marine Ecology Progress Series* **173**, 67–83.
- De Wolf P** (1973) Ecological observations on the mechanisms of dispersal of barnacle larvae during planktonic life and settling. *Netherlands Journal of Sea Research* **6**, 1–129.
- Doherty P and Fowler A** (1994) Demographic consequences of variable recruitment to coral reef fish populations: a congeneric comparison of two damselfishes. *Bulletin of Marine Science* **54**, 297–313.
- Freeman SM, Richardson CA and Seed R** (2001) Seasonal abundance, spatial distribution, spawning and growth of *Astropecten irregularis* (Echinodermata: Asteroidea). *Estuarine, Coastal and Shelf Science* **53**, 39–49.
- Frid CLJ, Buchanan JB and Garwood PR** (1996) Variability and stability in benthos: twenty-two years of monitoring off Northumberland. *ICES Journal of Marine Science* **53**, 978–980.
- Frid CLJ, Garwood PR and Robinson LA** (2009a) The North Sea benthic system: a 36 year time-series. *Journal of the Marine Biological Association of the United Kingdom* **89**, 1–10.
- Frid CLJ, Garwood PR and Robinson LA** (2009b) Observing change in a North Sea benthic system: a 33 year time series. *Journal of Marine Systems* **77**, 227–236.
- Gaines S and Roughgarden J** (1985) Larval settlement rate: a leading determinant of structure in an ecological community of the marine intertidal zone. *Proceedings of the National Academy of Sciences USA* **82**, 3707–3711.
- Giangrande A, Geraci S and Belmonte G** (1994) Life-cycle and life-history diversity in marine invertebrates and the implications in community dynamics. *Oceanography and Marine Biology: An Annual Review* **32**, 305–333.
- Gotelli NJ and Colwell RK** (2001) Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* **4**, 379–391.
- Graf G, Bengtsson W, Diesner U, Schulz R and Theede H** (1982) Benthic response to sedimentation of a spring phytoplankton bloom: process and budget. *Marine Biology* **67**, 201–208.
- Greenwood PJ** (1980) Growth, respiration and tentative energy budgets for two populations of the sea urchin *Parechinus angulosus* (Leske). *Estuarine and Coastal Marine Science* **10**, 347–367.
- Guillou M and Michel C** (1993) Reproduction and growth of *Sphaerechinus granularis* (Echinodermata: Echinoidea) in southern Brittany. *Journal of the Marine Biological Association of the United Kingdom* **73**, 179–192.
- Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D'Agrosa C, Bruno JF, Casey KS, Ebert C, Fox HE, Fujita R, Heinemann D, Lenihan HS, Madin EM, Perry MT, Selig ER, Spalding M, Steneck R and Watson R** (2008) A global map of human impact on marine ecosystems. *Science* **319**, 948–952.
- Herrmann M, Alfaya JEF, Lepore ML, Penchaszadeh PE and Laudien J** (2009) Reproductive cycle and gonad development of the Northern Argentinean *Mesodesma mactroides* (Bivalvia: Mesodesmatidae). *Helgolander Marine Research* **63**, 207.
- Highfield JM, Eloire D, Conway DVP, Lindeque PK, Attrill MJ and Somerfield PJ** (2010) Seasonal dynamics of meroplankton assemblages at station L4. *Journal of Plankton Research* **32**, 681–691.
- Hughes TP** (1990) Recruitment limitation, mortality, and population regulation in open systems: a case study. *Ecology* **71**, 12–20.
- John EH, Batten SD, Harris RP and Hays GC** (2001) Comparison between zooplankton data collected by the Continuous Plankton Recorder survey in the English Channel and by WP-2 nets at station L4, Plymouth (UK). *Journal of Sea Research* **46**, 223–232.
- Josefson AB, Jensen JN and Ærtebjerg G** (1993) The benthos community structure anomaly in the late 1970s and early 1980s – a result of a major food pulse? *Journal of Experimental Marine Biology and Ecology* **172**, 31–45.
- Jumars PA, Dorgan KM and Lindsay SM** (2015) Diet of worms emended: an update of polychaete feeding guilds. *Annual Review of Marine Science* **7**, 497–520.
- Kędra M, Kuliński K, Walkusz W and Legeżyńska J** (2012) The shallow benthic food web structure in the high Arctic does not follow seasonal changes in the surrounding environment. *Estuarine, Coastal and Shelf Science* **114**, 183–191.
- Kim H, Kim BH, Son MH, Jeon MA, Lee YG and Lee JS** (2016) Gonadal development and reproductive cycle of cultured abalone, *Haliotis discus hannai* (Gastropoda: Haliotidae) in Korea: implications for seed production. *Journal of Shellfish Research* **35**, 653–659.
- Kovala PE and Larrance JD** (1966) Computation of phytoplankton cell numbers, cell volume, cell surface and plasma volume per litre, from microscopic count. *University of Washington*, no. 38.
- Lewin R** (1986) Supply-side ecology: existing models of population structure and dynamics of ecological communities have tended to ignore the effect of the influx of new members into the communities. *Science* **234**, 25–27.
- Lopez GR and Levinton JS** (2011) Particulate organic detritus and detritus feeders in coastal food webs. In Wolanski E and McLusky DS (eds), *Treatise on Estuarine and Coastal Science*, **6**. Waltham, MA: Academic Press, pp. 5–21.
- Lozano J, Galera J, López S, Turon X, Palacín C and Morera G** (1995) Biological cycles and recruitment of *Paracentrotus lividus* (Echinodermata: Echinoidea) in two contrasting habitats. *Marine Ecology Progress Series* **122**, 179–191.
- MarLIN** (2006) *BIOTIC – Biological Traits Information Catalogue*. Plymouth: Marine Biological Association of the United Kingdom.
- Menden-Deuer S and Lessard EJ** (2000) Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. *Limnology and Oceanography* **45**, 569–579.
- Mincks SL, Smith CR and DeMaster DJ** (2005) Persistence of labile organic matter and microbial biomass in Antarctic shelf sediments: evidence of a sediment ‘food bank’. *Marine Ecology Progress Series* **300**, 3–19.
- Neumann H and Kröncke I** (2011) The effect of temperature variability on ecological functioning of epifauna in the German Bight. *Marine Ecology* **32**(s1), 49–57.
- Neumann H, Ehrich S and Kröncke I** (2008) Effects of cold winters and climate on the temporal variability of an epibenthic community in the German Bight. *Climate Research* **37**, 241–251.
- Neumann H, Ehrich S and Kröncke I** (2009a) Variability of epifauna and temperature in the northern North Sea. *Marine Biology* **156**, 1817–1826.
- Neumann H, Reiss H, Rakers S, Ehrich S and Kröncke I** (2009b) Temporal variability in southern North Sea epifauna communities after the cold winter of 1995/1996. *ICES Journal of Marine Science* **66**, 2233–2243.
- Nichols D and Barker MF** (1984a) A comparative study of reproductive and nutritional periodicities in two populations of *Asterias rubens* (Echinodermata: Asteroidea) from the English Channel. *Journal of the Marine Biological Association of the United Kingdom* **64**, 471–484.
- Nichols D and Barker MF** (1984b) Reproductive and nutritional periodicities in the starfish, *Marthasterias glacialis*, from Plymouth Sound. *Journal of the Marine Biological Association of the United Kingdom* **64**, 461–470.
- Nunes CDAP and Jangoux M** (2004) Reproductive cycle of the spatangoid echinoid *Echinocardium cordatum* (Echinodermata) in the southwestern North Sea. *Invertebrate Reproduction & Development* **45**, 41–57.
- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O'Hara RB, Simpson GL, Solynes P, Stevens MHH,**

- Szoecs E and Wagner H** (2017) Vegan Community Ecology Package. Available at <https://cran.r-project.org>.
- Olden JD and Neff BD** (2001) Cross-correlation bias in lag analysis of aquatic time series. *Marine Biology* **138**, 1063–1070.
- Pearson TH and Rosenberg R** (1986) Feast and famine: structuring factors in marine benthic communities. In Gee JHR and Gill PS (eds), *Organization of Communities*. Oxford: Blackwell Scientific, pp. 373–398.
- Peterson CH** (1979) Predation, competitive exclusion, and diversity in the soft-sediment benthic communities of estuaries and lagoons. In Livingston RJ (ed.), *Ecological Processes in Coastal and Marine Systems*. Boston, MA: Springer, pp. 233–264.
- Peterson CH and Fegley SR** (1986) Seasonal allocation of resources to growth of shell, soma, and gonads in *Mercenaria mercenaria*. *Biological Bulletin* **171**, 597–610.
- Posey M, Powell C, Cahoon L and Lindquist D** (1995) Top down vs. bottom up control of benthic community composition on an intertidal tideflat. *Journal of Experimental Marine Biology and Ecology* **185**, 19–31.
- Probst WN, Stelzenmüller V and Fock HO** (2012) Using cross-correlations to assess the relationship between time-lagged pressure and state indicators: an exemplary analysis of North Sea fish population indicators. *ICES Journal of Marine Science* **69**, 670–681.
- Queirós AM, Stephens N, Widdicombe S, Tait K, McCoy SJ, Ingels J, Rühl S, Aïrs R, Beesley A, Carnovale G, Cazenave P, Dashfield S, Hua E, Jones M, Lindeque P, McNeill CL, Nunes J, Parry H, Pascoe C, Widdicombe C, Smyth T, Atkinson A, Krause-Jensen D and Somerfield PJ** (2019) Connected macroalgal-sediment systems: blue carbon and food webs in the deep coastal ocean. *Ecological Monographs*, e01366.
- Quijón PA and Snelgrove PVR** (2005) Predation regulation of sedimentary faunal structure: potential effects of a fishery-induced switch in predators in a Newfoundland sub-Arctic fjord. *Oecologia* **144**, 125–136.
- Quijón PA, Kelly MC and Snelgrove PVR** (2008) The role of sinking phyto-detritus in structuring shallow-water benthic communities. *Journal of Experimental Marine Biology and Ecology* **366**, 134–145.
- Reiss H and Kröncke I** (2004) Seasonal variability of epibenthic communities in different areas of the southern North Sea. *ICES Journal of Marine Science* **61**, 882–905.
- Sastry AN** (1966) Temperature effects in reproduction of the bay scallop, *Aequipecten irradians* Lamarck. *Biological Bulletin* **130**, 118–134.
- Sastry AN and Blake NJ** (1971) Regulation of gonad development in the bay scallop, *Aequipecten irradians* Lamarck. *Biological Bulletin* **140**, 274–283.
- Scrosati RA and Ellrich JA** (2017) Recruitment and abundance of intertidal barnacles and mussels along the Atlantic Canadian coast: pelagic influences and relationships with predator abundance. *Ecosphere*, e02229.
- Sköld M and Gunnarsson JSG** (1996) Somatic and germinal growth of the infaunal brittle stars *Amphiura filiformis* and *A. chiajei* in response to organic enrichment. *Marine Ecology Progress Series* **142**, 203–214.
- Smyth T, Atkinson A, Widdicombe S, Frost M, Allen I, Fishwick J, Queirós AM, Sims D and Barange M** (2015) The Western Channel observatory. *Progress in Oceanography* **137**(Part B), 335–341.
- Stoffer D** (2016) *astsa* – Applied Statistical Time Series Analysis, R package. Available at <https://github.com/nickpoison/astsa>.
- Sutherland JP** (1990) Recruitment regulates demographic variation in a tropical intertidal barnacle. *Ecology* **71**, 955–972.
- Tait K, Aïrs RL, Widdicombe CE, Tarran GA, Jones MR and Widdicombe S** (2015) Dynamic responses of the benthic bacterial community at the Western English Channel observatory site L4 are driven by deposition of fresh phytodetritus. *Progress in Oceanography* **137**, 546–558.
- Thorson G** (1950) Reproductive and larval ecology of marine bottom invertebrates. *Biological Reviews* **25**, 1–45.
- Thronsen J** (1978) Productivity and abundance of ultra- and nanoplankton in Oslofjorden. *Sarsia* **63**, 273–284.
- Thrush SF** (1999) Complex role of predators in structuring soft-sediment macrobenthic communities: implications of changes in spatial scale for experimental studies. *Austral Ecology* **24**, 344–354.
- Trapletti A, Hornik K and LeBaron B** (2017) *tseries* – Time Series Analysis, R package. Available at <https://cran.r-project.org/package=tseries>.
- Underwood AJ and Fairweather PG** (1989) Supply-side ecology and benthic marine assemblages. *Trends in Ecology & Evolution* **4**, 16–20.
- UNESCO** (1968) *Monographs on Oceanographic Methodology: Zooplankton Sampling*. Paris: UNESCO.
- Widbom B and Frithsen JB** (1995) Structuring factors in a marine soft bottom community during eutrophication – an experiment with radio-labelled phytodetritus. *Oecologia* **101**, 156–168.
- Widdicombe S and Austen MC** (2001) The interaction between physical disturbance and organic enrichment: an important element in structuring benthic communities. *Limnology and Oceanography* **46**, 1720–1733.
- Widdicombe CE, Eloire D, Harbour D, Harris RP and Somerfield PJ** (2010) Long-term phytoplankton community dynamics in the Western English Channel. *Journal of Plankton Research* **32**, 643–655.
- Wiekling G and Kröncke I** (2005) Is benthic trophic structure affected by food quality? The Dogger Bank example. *Marine Biology* **146**, 387–400.
- Williams DMCB** (1980) Dynamics of the Pomacentrid community on small patch reefs in One Tree Lagoon (Great Barrier Reef). *Bulletin of Marine Science* **30**, 159–170.
- Włodarska-Kowalczyk M, Górska B, Deja K and Morata N** (2016) Do benthic meiofaunal and macrofaunal communities respond to seasonality in pelagic processes in an Arctic fjord (Kongsfjorden, Spitsbergen)? *Polar Biology* **39**, 2115–2129.
- Woodin SA** (1999) Shallow water benthic ecology: a North American perspective of sedimentary habitats. *Australian Journal of Ecology* **24**, 291–301.
- Zeileis A, Grothendieck G, Ryan JA, Ulrich JM and Andrews F** (2018) *Zoo: S3 Infrastructure for Regular and Irregular Time Series (Z's Ordered Observations)*, R package. Available at <http://zoo.r-forge.r-project.org/>.
- Zhang Q, Warwick RM, McNeill CL, Widdicombe CE, Sheehan A and Widdicombe S** (2015) An unusually large phytoplankton spring bloom drives rapid changes in benthic diversity and ecosystem function. *Progress in Oceanography* **137**(Part B), 533–545.