Article: 2848

Topic: 46 - Eating Disorders

SECOND GENERATION ANTIPSYCHOTIC-INDUCED WEIGHT ALTERATION

N. Ron Wan¹, A. Silva², M.d.L. Pequeno², J. Ropero Peláez³, J. Gonçalves², **S. Taniguchi**⁴

¹São Luis Hospital, ²Albert Einstein Hospital, ³UFABC, ⁴Basics Sciences, Albert Einstein Hospital, São Paulo, Brazil

Introduction: Atypical antipsychotics are frequently associated with increased risk of weight gain as side effect.

Objective: To compare body weigh alteration in psychotic disorder patients undergoing atypical neuroleptics therapy.

Methods: This study included 37 patients with mean age of 35,14±1,57 years old, enrolled in a public mental health service with psychotic disorders related to the use of illicit drugs and schizophrenia.

Results: Patients treated with atypical neuroleptics such as olanzapine (11) 0,28mg/kg/day, clozapine (7) 1,56 mg/kg/day and risperidone (10) 0,031 mg/kg/day, were evaluated. Haloperidol (9) 0,072 mg/kg/day, as typical neuroleptics was also studied. Neuroleptics' treatment period was $126,97\pm20,93$ days and the patients inicial body weight mean was $67,63\pm2,13$ kg. Increase of body weight of $1,79\pm0,72$ kg was observed for patients receiving haloperidol, compared with olanzapine $10,37\pm1,74$ kg, clozapine $3,6\pm2,10$ kg and risperidone $1,21\pm0,90$ kg in the end of evaluation period.

These increase of body weight calculated in g/day were 31,68±1,83 g/day for haloperidol, 99,54±25,55 g/day for olanzapine, 27,44±13,35 g/day for clozapine and 13,11±1,40g/day for risperidone.

Increase of body weight index (BWI) at the end of the period of analysis was 0,66±0,25 for haloperidol, 3,59±0,63 for olanzapine, 1,52±0,90 for clozapine and 0,40±0,35 for risperidone.

These BWI increases identified per day were 0,011±0,0038 for haloperidol, 0,034±0,0085 for olanzapine, 0,011±0,0057 for clozapine and 0,0039±0,0055 for risperidone.

Conclusions: Although haloperidol, olanzapine, clozapine and risperidone act blocking serotonin receptors conducting to body weight gain, they probably act with different intensity degree of blockage.