## **Index of Subjects**

Airborne infection in a fully air-conditioned hospital, I, 15; II, 31; III, 45; IV, 445

Antibiotic therapy and the faecal excretion of Salmonella typhimurium by experimentally infected chickens, 275

Autoclaves and their dangers, and safety in laboratories, 475

Bacteriological tests as indices for the development of off-flavours in cream, 113

Brucella spp., Yersinia enterocolitica serotype IX and Salmonella serotypes: serological relationship, 151

Camelpox virus from Kenya, characteristics of, 381

Citrobacter, koseri. I. An extended antigenic scheme for, 121; II. Serological and biochemical examination of, 129

Clostridium botulinum in the lakes and waterways of London, 371

Contamination of fluids from a hospital pharmacy, 87

Corynebacterium diphtheriae strains isolated in Victoria from 1962 to 1971, numerical analysis of the characteristics of, 413

Cream, bacteriological tests as indices for the development of off-flavours in, 113

Dairy factories: sludge from as a potential hazard to animal health, 143

Diphtheria in Victoria, 1952–1971. Numerical analysis of the characteristics of the C. diphtheriae strains isolated, 413

Endemic infection in surgical wards, 185

Endotoxins detected by the Limulus test in burned and unburned mice infected with different species of gram-negative bacteria, 99

Epidemiology of ephemeral fever in Kenya, 231

Epidemiology of Rift Valley fever in Kenya, 219

Faecal flora of children in the United Kingdom, 135

Feeding diets containing permitted antibiotics and the faecal excretion of Salmonella typhimurium by experimentally infected chickens, 293

Foot-and-mouth disease vaccination, hypersensitivity in cattle: response to hydroxy-propylmethylcellulose, 79

Foot-and-mouth disease virus type A, variation among strains in the Eastern Mediterranean region, 1964–1972, 387

Haemophilus (acute) infections of children: investigation of the family background of, 315 Hospital pharmacy, contamination of fluids from, 87

Hydatid cysts in man, serologic evidence of the postoperative persistence of, 333

Hypersensitivity in cattle after foot-and-mouth disease vaccination: response to hydroxy-propylmethylcellulose, 79

Influenza surveillance, a collaborative programme for, 1

Influenza virus, a new surface-antigen-adsorbed vaccine, I, 341; II, 353

Influenza virus vaccines, live attenuated and detergent split, comparative trials of, 425

Leprosy, multiple skin testing in, 57

Limulus test, detection of endotoxins with, in burned and unburned mice infected with different species of gram-negative bacteria, 99

Myxoma virus strains of differing virulence grades, differential transmissibility of by the rabbit flea Spilopsyllus cuniculi (Dale), 237

Naegleria fowleri, the effect of thermal pollution on the distribution of, 7

Persistence of antibody induced by rubella vaccine after six years, 407

Persistence, postoperative, of hydatid cysts in man, 333

Pseudomonas aeruginosa infection in hospital: a comparison between 'infective' and 'environmental strains', 195

Removal of total organisms and pathogens from the skin, 259

Rhinovirus colds through indirect contact, possible transmission of, 249

R factors, transfer in vivo between Escherichia coli strains inoculated into the rumen of sheep, 363

Rubella, serological investigation of cases and contacts of, 91

Rubella vaccine (Wistar RA 27/3 strain), persistence of antibody induced by, after six years, 407

Salmonellas, isolation of, from British pork sausages and sausage meat, 173

Salmonella typhimurium in C57BL/6 mice, experimental model of oral antityphoid vaccination with live streptomycin-dependent, 215

Salmonella typhimurium, effects of antibiotic therapy on the faecal excretion by experimentally infected chickens, 275, 293

Salmonella vaccines, live streptomycin-dependent: the problem of reversion to a virulent form, 203

Serological investigation of cases and contacts of rubella, 91

Serological and biochemical examination of Citrobacter koseri, 129

Serological relationship between Brucella spp., Yersinia enterocolitica serotype IX and Salmonella serotypes of Kauffmann-White group N, 151

Skin, removal of total organisms and pathogens from, 259

Skin testing of Kenyan schoolchildren with a series of new tuberculins, 303

Skin testing in leprosy, 57

Sludge from dairy factories: a potential hazard to animal health, 143

Staphylococcus aureus and its nasal acquisition by patients in a fully air-conditioned hospital, 445

Streptococcal infection in young pigs, IV, 69

Thermal pollution and the distribution of Naegleria fowleri, 7

Transfer in vivo of R factors between Escherichia coli strains inoculated into the rumen of sheep, 363

Transmissibility of myxoma virus strains of differing virulence grades by the rabbit flea Spilopsyllus cuniculi (Dale), 237

Vaccine, a new surface-antigen-adsorbed, against influenza virus, I, 341; II, 353

Vaccine, rubella, persistence of antibody induced by, after six years, 407

Vaccines, comparative trials of live attenuated and detergent split influenza virus, 425

Variation among strains of type A foot-and-mouth disease virus in the Eastern Mediterranean region 1964–1972, 387

Vibrio cholerae, type specific action of vibriocidal antibody on, 399

Viruses: camelpox, 381; ephemeral fever, 231; foot-and-mouth, 79, 387; influenza, 341, 353, 425; myxoma, 237; rhinovirus, 249; Rift Valley fever, 219