

## INTERPERSONAL PSYCHOTHERAPY FOR TREATING PSYCHOLOGICAL DISTURBANCES IN ADOLESCENTS WHO EXPERIENCED BULLIED EXPERIENCES: RANDOMIZED CASE CONTROL STUDY

*T.-C. Tang*<sup>1</sup>, *S.-Y. Huang*<sup>2</sup>

<sup>1</sup>Psychiatry, Kaohsiung Medical University Chung-Ho Memorial Hospital, <sup>2</sup>Psychiatry, Kaohsiung Normal University, Kaohsiung, Taiwan R.O.C.

**Background:** This case-control study aimed to assess the intervention effects of six-session interpersonal psychotherapy (IPT-A) on reducing the severity of anxiety and depression in adolescent victims.

**Methods:** A total of 30 adolescents who had clinical significant level after experiencing bullied experiences were allocated to a six-session course of IPT-A (N =15) or to treatment as usual (TAU) (N = 15). T test was performed to examine the effect of IPT-A on reducing the severity of anxiety and depression related to the bullied events.

**Results:** Pre-intervention age, sex, anxiety and depression showed no significant difference between two groups. As the pre-intervention severity of two groups were no significant different, results showed the IPT-A group to have significantly lower post-intervention severity levels of anxiety and depression ( $p < .05$ ) than the TAU group. Effective size showed moderate to high level between IPT-A and TAU.

**Conclusions:** The results of this study support the effectiveness of the IPT-A in improving anxiety symptoms and depression in adolescents experiencing traumatic bullied experiences.