# S22 Structure quality of mental health care systems in Europe

# Recent changes in the Hungarian mental health care system.

### Füredy, Janos (Budapest/Hungary)

Hungary is with ist ca 10 millions of inhabitants one of the smaller European country with a glorious past. Allthoug according to its image as a rural country more than the half of the inhabitants live in the five biggest towns. Before the political change during the last decade because of the communistic ideology the mental health area was widely neglected by the Hungarian Government. Inpatient care was provided by bigger state hospitals, whereas the outpatients care has been provided by governmental outpatients clinic. Since the political change and increasing number of psychiatrists begin to work in private practice and postgraduate training is improving. However, because of its famous psychoanalytical tradition for instance Ferency and Balint have been of Hungarian offspringpsychotherapeutic thinking has maintained in our county. The orientation of Hungarian psychiatry has been directed more to Germany than to other countries. Now decentralization of hospital care as one form of reforms in Hungarian psychiatry is advancing, though the lack of finances is tremendous and the main obstacle to promote reforms quicker. The main part of psychiatrists is concentrated in the towns with the result of poor outpatient facility equipment in the rural areas. A concept of improving the situation will be presented.

# S23 How to teach psychosocial skills to general practitioners? TEACHING PSYCHOSOMATIC MEDICINE - A TWO YEAR

# CURRICULUM FOR POST-GRADUATE CLINICIANS

### Claus Buddeberg, Zurich, Switzerland.

25 to 30% of the patients seen in a general practitioner's office suffer from psychosomatic disorders. In Switzerland there is no structured teaching of a psychosomatic approach to illness, neither in medical school nor during postgraduate training. In this presentation experiences with a two year training program in psychosomatic medicine for post-graduate clinicians are reported. The training program has three parts: Part one consists of 12 training days over a two year period given in seminars with lectures and case presentations using videotapes, one way mirror as well as role plays. Part two is case supervision in small groups. In part three participants are taught and get experiences in a self relaxation technique. In our opinion the two year curriculum serves as a model and example for a certified training in "Psychomatic Medicine". The participants can benefit from their training in various ways: with regard to the organization and atmosphere of their practice, in respect of improved communication skills when treating patients and advanced small group interaction skills. We have just finished the second course and we received excellent feed-back from most of the participants.

# S23 How to teach psychosocial skills to general practitioners?

BEHAVIORAL SCIENCE CURRICULA FOR GENERAL INTERNAL MEDICINE: LEADING THE PEARFUL AND PREACHING TO THE CONVERTED

### S. R. Hahn.

The goal of teaching psychiatry and bahavioral sciencies to primary care phyricians is wisque in both is importance and in the challenges it presents. The gap between need and actual proction in the delivery of psychiatric treatment in the general medicine secting is wish. Medical renderiis and faculty's skills and knowledge and willingness to learn about psychiatry and bahavior sciencies are very heterogeneous renging from "the fearful" or "the convented". However, the spainty of case that will be delivered in general medicine depends on enhancing the skills and knowledge of the majority of processors, not merely a handful of anthusiasts. The magnitude of this task closuse that behavioral science and psychiatry curricula be taught by the majority of primary care faculty and not merely is handful of quasi-subspecialized internists or mental health specialists. The behavioral sciences' curriculum for our large general medicine internists or mental health specialists. The behavioral sciences' curriculum for our large general medicine faculty. Our primary care-trained faculty field comfortable with the clinical management of most psychiatric psychiaters demandishes programs was designed to be taught primaryly by our "regular" general medicine faculty. Our primary care-trained faculty field comfortable with the clinical management of most psychiatric psychiatric design as or subscalable of the production of psychopathology, family and medical problems. "Simulated-patients' are miscriword by resolution in a small group setting using a name of inaching sechiaques that this advantage of the smellation. The simulated-patients sessions are supplemented by presentations on specific techniques or topor such as small group or quasi-subspecialized faculty. By courrast, the certiculum for our resulting (12 medients) "Primary Care" medicines programs which traditionally astronic sussessment that are presented by a small group of quasi-subspecialized faculty. By courrast, the certiculum for our resulting (11 hulle disceptionar

## S23 How to teach psychosocial skills to general practitioners?

GETTING MEDICAL SPECIALISTS INVOLVED IN PSYCHOSOCIAL CARE: A ROLE FOR GENERAL INTERNISTS?

P. Cathebras Service de Medicine Interne, Hôpitaux de St Etienne, 42055 St Etienne, France

Internal medicine holds a special position in France. Confined to tertiary care settings, it is considered as a speciality among other medical specialities. In the meantime, internists favour a holistic perspective very similar to the GP's approach, and opposed to most specialists point of view. Since consultation-liaison psychiatry is poorly developed in many hospitals, internists sometimes have to act as consultants for clinical problems situated outside the organ specialities domain, including psychosomatic problems. This means that internists have a special role in educating their fellow specialists on emotional factors in physical disease, attention to psychosocial context, patient-centred approach and emphasis on "illness' vs. disease perspective in the management of symptoms. Our experience as internists as psychosocial specialists in the general hospital environment will be briefly discussed.