

Are Dumbbell Brightest Cluster Members Signposts to Galaxy Cluster Activity?

K. A. Pimblet^A

^A Department of Physics, University of Queensland, Brisbane, Queensland, QLD 4072, Australia.
Email: pimblet@physics.uq.edu.au

Received 2008 June 30, accepted 2008 August 13

Abstract: We assemble a sample of galaxy clusters whose brightest members are dumbbell galaxies and compare them with a control sample in order to investigate if they are the result of recent mergers. We show that the dumbbell sample is no more likely than other clusters to exhibit subclustering. However, they are much more likely to have at least one dumbbell component possessing a significant peculiar velocity with respect to the parent cluster than a non-dumbbell brightest cluster member. We interpret this in the context of seeing the clusters at various stages of post-merger relaxation.

Keywords: galaxies: clusters: general — galaxies: elliptical and lenticular, cD — galaxies: evolution

1 Introduction

Brightest cluster member (BCM) galaxies have historically held a special place in both the theoretical and observational study of galaxy evolution and formation since they are believed to be directly connected with the conditions in the cluster that they reside in. Consider, for instance, a BCM that is located at the bottom of the gravitational well of a galaxy cluster: there, a process such as cannibalism (e.g. Ostriker & Tremaine 1975) would be at maximum efficiency there and we would be likely to observe BCMs with multiple cores (e.g. Laine et al. 2003; Yamada et al. 2002; Oegerle & Hill 2001; Dubinski 1998; Lauer 1988; Hoessel & Schneider 1985). However, if a more hierarchical paradigm is correct, then the BCM should have formed through multiple minor merger events in a sub-group that was originally outside the cluster centre. It would have then subsequently merged with the rest of the cluster (Merritt 1985), yielding several observational road-signs such as high peculiar velocities for the BCM and cluster substructure (see Woudt et al. 2008; Pimblet et al. 2006; Oegle & Hill 2001; Pinkney et al. 1996; Quintana et al. 1996; Valentijn & Casertano 1988).

Cases of dumbbell galaxies as BCMs (e.g. Gregorini et al. 1994, 1992) are doubly interesting for these reasons — these galaxies should be an indicator of the pre-virialization scenario and should therefore be accompanied by significant substructure indicative of recent cluster merger activity. Indeed, Quintana et al. (1996) present evidence for a dumbbell system where the two components of the BCM *each* belong to major sub-groups that are undergoing a merger (see also de Souza & Quintana 1990).

What is unclear is whether dumbbell BCM clusters in general are special? Does the presence of a dumbbell BCM point toward observable cluster activity such

as subclustering (Quintana et al. 1996) and large BCM peculiar velocities (Pimblet et al. 2006) as would be found during a merger event? If so, then at what stage is the merger event? In order to make a pass at answering these questions, this work assembles a modest sample of dumbbell BCMs from Gregorini et al. (1992) and a control sample derived from the 2 degree field galaxy redshift survey (2dFGRS; Colless et al. 2001).

The format of this paper is as follows. In Section 2 we fully describe our derived dumbbell sample and control sample. In Section 3 we ask whether clusters with dumbbell BCMs are more likely to have substructuring than any other clusters and then in Section 4 we examine the incidence of significant peculiar velocities in our samples are different. We summarize our findings in Section 5. Throughout this work, we adopt a cosmological concordance model with values of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$.

2 The Samples

The Gregorini et al. (1992) sample claims to be a volume limited & homogeneous sample of dumbbell BCM galaxies which should be ideal for our purposes. It is based on the Abell catalogue (Abell 1958; Abell et al. 1989) and is complete out to a comoving distance of $210h^{-1} \text{ Mpc}$ (Gregorini et al. 1992). In addition, they state there are no possible selection biases present in this sample that would have prevented the successful detection and identification of dumbbell galaxies for this sample (Gregorini et al. 1992; see also Scaramella et al. 1991). There are a couple of debatable clusters that could have been added to this sample (e.g. Abell 3323; Gregorini et al. 1994), but were left out. We do not view the exclusion of these clusters as having a significant impact on the ensemble. However, our review of the available literature redshifts

demonstrates that this sample does contain several clusters with $z > 0.1$ (Table 1) due to earlier cluster redshifts being under-estimated. We do not regard this as a major impediment to the present investigation either, since the time evolved between $z \sim 0.07$ and $z \sim 0.10$ is much smaller (i.e. < 0.5 Gyr) than the expected time it would take for sub-clusters to fully merge (Lacey & Cole 1993). For this work, we restrict ourselves to the complete sample of dumbbell galaxies — Table 1 from Gregorini et al. (1992, herein the dumbbell sample) — with one exception: Abell 3653. Abell 3653 is listed by Gregorini et al. (1992) as only a ‘possible’ dumbbell BCM. However, a combination of observations made by Postman & Lauer (1995) and Pimblet et al. (2006) show that this cluster should also be considered to be a confirmed dumbbell BCM galaxy (Figure 1) and we include it in the present dumbbell sample.

Redshift information for the dumbbell sample is initially obtained by downloading all redshifts within 30 arcmin of each cluster from the NASA Extra-Galactic Database (NED). At $z \sim 0.1$, this corresponds to a radius of 1.6 Mpc from the cluster centre. Several clusters in the resultant dumbbell sample (Abell 2824, 3098, 3368, 3397 & 3740) generate very few members ($N < 20$) within 30 arcmin (\approx an Abell radius) and are eliminated from the final sample at this stage since it is likely that any subclustering or cluster velocity dispersion measurement would be unreliable (Girardi et al. 1993). For the remaining sample, we compute the mean cluster velocity and velocity dispersion (σ) using the ‘gapping’ procedure of Zabludoff et al. (1990, 1993) and tabulate these values in Table 1. The final sample yields 13 dumbbell BCM galaxy clusters for us to work with that have a range of velocity dispersions consistent with large and massive clusters (see also Ebeling et al. 2007).

Our control sample is obtained from the 2dFGRS cluster sample of De Propriis et al. (2002; see Colless et al. 2001 for a description of this survey). This is essentially a complete catalogue at $cz < \sim 35\,000$ km s⁻¹ of Abell (Abell 1958; Abell et al. 1989) clusters. In order to attempt to match the relatively high velocity dispersions in the dumbbell sample, we restrict the control sample to only those clusters with a high X-ray luminosity ($L_X > 0.5 \times 10^{44}$ erg s⁻¹, Ebeling et al. 1996; Cruddace et al. 2002). We also trim from this sample any clusters with poor completeness levels (i.e. less than 20 galaxies within an Abell radius of the cluster centre). Our final control sample consists of 14 clusters and is presented in Table 2 with mean velocities and velocity dispersions sourced from De Propriis et al. (2002). Although the control sample has a smaller absolute range of σ (597–1038 km s⁻¹) than the dumbbell cluster sample (455–1376 km s⁻¹), the median values (783.5 km s⁻¹ and 825 km s⁻¹ respectively) are very similar. The average redshift of the control sample ($cz = 26\,010$ km s⁻¹) is also somewhat higher than the dumbbell sample ($cz = 18\,854$ km s⁻¹). In look-back time this is barely 0.3 Gyr, so again we do not regard this as significant for

this work due to the comparative time taken for clusters to violently relax after a merger event (Lacey & Cole 1993). With the control sample being sourced from 2dFGRS, we ensure that all the constituent galaxies are very homogeneously sampled down to $b_J = 19.45$ (Colless et al. 2001); albeit not at 100% completeness levels (see De Propriis et al. 2002). Conversely, our dumbbell sample is a collection of diverse redshifts from multiple sources each with different purposes and associated selection limits imposed (see Table 1). We also eyeball the BCMs in the control sample to ensure that they are not dumbbells themselves that are outside the Gregorini et al. (1992) selection limits. Only two of them give us cause for concern: Abell 2811 has a nearby faint companion, but would not be considered a dumbbell in the definition of Gregorini et al. (1992); Abell S1136 has multiple galaxies near to the BCM that may be interacting with it, but again it is not a dumbbell and its removal from the control sample does not affect our conclusions.

3 Substructure

In order to place the clusters onto a common scale and perform a meaningful comparison of subclustering, we use a virial radius estimator derived by Girardi et al. (1998): $R_v = 0.002\sigma h_{100}^{-1}$ Mpc. Cluster members are then defined to be those galaxies whose velocity is within 3σ of the cluster velocity. We add the caveat that although this approximation’s validity for semi-virialized systems may not be ideal, it is sufficient for our purposes of placing the clusters on to an approximately common scale.

For the cluster centres, we use the quoted NED cluster centres. This choice of cluster centre is somewhat arbitrary, but it will not dramatically affect the final subclustering result since we are sampling the cluster members from a relatively large radius away from this centre. The resultant number of galaxies and values for R_v for each of the clusters are given in Tables 1 and 2.

There are a number of statistical tools available to assess the degree of subclustering in each cluster, ranging from 2 dimensional searches for asymmetry (e.g. West et al. 1988) and bimodality (e.g. Fitchett & Webster 1987) to more complex 3 dimensional tests (e.g. Dressler & Shectman 1988; among others). Given all these different tools, Pinkney et al. (1996) made extensive tests to determine the relative merits of these tools and concluded that the Dressler & Shectman (1988) Δ test is the best one to use to find substructure in arbitrary cases. Its only real limitation is an insensitivity to equal mass mergers in the plane of the sky and superpositions of sub-groups (Pinkney et al. 1996). Both of these situations would require special and unusual lines of sight to the cluster and are therefore considered to be rare events. Importantly for this study, the DS test will be able to detect substructure in 3:1 mergers with reasonable confidence (circa 95%) down to a sample size of even 30 galaxies (Figure 27 in Pinkney et al. 1996). At a sample size of > 60 , this confidence rapidly grows to $> 99\%$.

Table 1. The dumbbell-cluster sample^a

Cluster	RA (J2000)	Dec (J2000)	$N(30')^b$	cz (km s^{-1})	σ (km s^{-1})	R_V (Mpc)	N_{R_V}	Δ_{R_V}	$P(\Delta)_{R_V}$	N_{2R_V}	Δ_{2R_V}	$P(\Delta)_{2R_V}$
Abell 2860	1 ^h 4 ^m 20.60 ^s	-39°29'61.4"	59	31 738 ± 97	741 ⁺⁸⁰ ₋₆₀	1.48	28	76	0.990	45	148	0.985
Abell 2911	1 ^h 26 ^m 12.87 ^s	-37°33'81.8"	50	24 110 ± 102	720 ⁺⁸⁵ ₋₆₃	1.44	34	30	0.720	50	41	0.959
Abell 3151	3 ^h 40 ^m 27.73 ^s	-28°25'43.0"	68	19 992 ± 124	1013 ⁺¹⁰¹ ₋₇₈	2.03	60	115	0.009	103	230	0.016
Abell 3266	4 ^h 31 ^m 11.92 ^s	-61°14'42.6"	286	17 852 ± 81	1376 ⁺⁶¹ ₋₅₄	2.75	248	277	0.313	399	524	<0.001
Abell 0533	5 ^h 1 ^m 30.79 ^s	-22°22'02.4"	27	13 932 ± 229	1191 ⁺²⁰⁸ ₋₁₃₆	2.38	35	92	<0.001	52	146	<0.001
Abell 3391	6 ^h 26 ^m 15.43 ^s	-53°24'52.3"	104	16 070 ± 127	1294 ⁺¹⁰¹ ₋₈₇	2.59	131	207	<0.001	241	440	<0.001
Abell 3528	12 ^h 54 ^m 18.36 ^s	-29°75.6"	146	16 219 ± 87	1047 ⁺⁵⁹ ₋₄₉	2.09	145	194	0.017	236	327	<0.001
Abell 3532	12 ^h 57 ^m 19.08 ^s	-30°13'32.7"	167	16 504 ± 60	780 ⁺⁴⁰ ₋₃₀	1.56	117	150	0.061	233	303	0.018
Abell 3535	12 ^h 57 ^m 48.60 ^s	-28°17'52.1"	55	20 190 ± 61	455 ⁺⁵¹ ₋₃₈	0.91	37	46	0.041	53	78	0.010
Abell 3570	13 ^h 46 ^m 50.88 ^s	-37°32'98.3"	31	11 261 ± 83	463 ⁺⁷⁴ ₋₅₀	0.93	19	17	0.589	37	34	0.627
Abell 3653	19 ^h 52 ^m 37.92 ^s	-52°74.1"	106	32 234 ± 81	836 ⁺⁶⁴ ₋₅₂	1.67	50	61	0.164	92	123	0.044
Abell 3716	20 ^h 51 ^m 16.56 ^s	-52°25'03.4"	124	13 517 ± 74	825 ⁺⁵⁸ ₋₄₈	1.65	110	113	0.226	196	365	<0.001
Abell 3744	21 ^h 7 ^m 13.80 ^s	-25°17'33.7"	69	11 483 ± 72	596 ⁺⁵⁹ ₋₄₅	1.19	64	88	0.066	88	129	0.007

^aRedshifts for this sample are sourced from: Fairall 1988; Teague, Carter & Gray 1990; da Costa et al. 1991; Dalton et al. 1994; Collins et al. 1995; Postman & Lauer 1995; Quintana et al. 1995; Loveday et al. 1996; Quintana & Ramirez 1996; Shectman et al. 1996; Caldwell & Rose 1997; da Costa et al. 1998; Katgert et al. 1998; Vettolani et al. 1998; Wegner et al. 1999; Schindler 2000; Bardelli et al. 2001; Colless et al. 2001; Donnelly et al. 2001; Christlein & Zabludoff 2003; Kaldare et al. 2003; Paturel et al. 2003; Jones et al. 2004; Smith et al. 2004; Pimbblet et al. 2006.

^bThis is the number of NED galaxies within 30 arcmin used to compute cz and σ from.




Figure 1 Spectroscopic observations of the components of the dumbbell BCM galaxy in Abell 3653 have been made by Postman & Lauer (1995) and Pimblet et al. (2006); the circles in the plot denote the location of where the redshift was taken. They give redshifts of $z = 0.1091 \pm 0.0003$ and $z = 0.1099 \pm 0.0002$ respectively. Since both components are within 2σ of each other, we consider this galaxy to be a confirmed dumbbell BCM galaxy for the purposes of this work.

We therefore proceed by applying the Dressler & Shectman (1988) approach to each of our clusters and we refer the reader to that publication for details of its algorithmic execution. For the purposes of this work, it is sufficient to note that if there is little or no substructure, then we may expect the resultant Δ statistic to be of approximately the same value as the number of galaxies sampled. The final parameter of merit is then $P(\Delta)$ which gives the probability of the observed value of Δ occurring randomly when the redshifts of cluster members are randomly assigned to other members in a Monte Carlo fashion (i.e. very low values of $P(\Delta)$ indicate the presence of substructure).

Values of Δ and $P(\Delta)$ are computed for each cluster within radial limits of both R_V and $2R_V$ (Tables 1 and 2). The reason for looking at both these radii is that it may be possible that the Δ statistic is insensitive to substructure at small (<2 Mpc) radii (Pinkney et al. 1996). We also plot the ‘bubble plots’ in Figure 2 for the results of the DS test for the dumbbell sample. In these plots, substructure can be interpreted as spatially close overlapping circles (Dressler & Shectman 1988). Within R_V , we find that the fraction of clusters with certain substructure (i.e. $P(\Delta) < 0.001$) is 2/13 (15%) for the dumbbell sample and 1/14 (7%) for the control sample. At $2R_V$, these fractions change to 5/13 (39%) and 4/14 (29%) respectively (in addition, a further two of control clusters are very close to being considered as having subclustering). We consider the difference in subclustering fractions between the two cluster samples to be statistically insignificant.

Table 2. The control-cluster sample

Cluster	RA (J2000)	Dec (J2000)	cz (km s ⁻¹)	σ (km s ⁻¹)	R_V (Mpc)	N_{R_V}	Δ_{R_V}	$P(\Delta)_{R_V}$	N_{2R_V}	Δ_{2R_V}	$P(\Delta)_{2R_V}$
Abell 2734	0 ^h 11 ^m 19.44 ^s	-28°30'99.6"	18 646 ± 94	1038 ⁺⁷³ ₋₈₄	2.08	120	153	0.127	214	427	<0.001
Abell 2751	0 ^h 16 ^m 19.92 ^s	-31°13'14.0"	31 863 ± 103	763 ⁺⁸⁰ ₋₉₅	1.47	43	67	0.014	99	128	0.027
Abell 2755	0 ^h 17 ^m 34.80 ^s	-35°66'9.6"	28 469 ± 135	829 ⁺¹⁰² ₋₁₂₉	1.66	29	21	0.844	64	87	0.180
Abell 2798	0 ^h 37 ^m 27.12 ^s	-28°18'86.4"	33 516 ± 108	739 ⁺⁸⁴ ₋₁₀₂	1.48	41	60	0.008	77	120	0.004
Abell 2811	0 ^h 42 ^m 8.88 ^s	-28°19'33.2"	32 557 ± 168	988 ⁺¹²⁶ ₋₁₆₂	1.98	83	111	0.193	175	264	<0.001
Abell 2829	0 ^h 51 ^m 19.20 ^s	-28°18'68.4"	33 565 ± 123	793 ⁺⁹⁴ ₋₁₁₇	1.59	39	50	0.043	69	118	0.010
Abell 3027	2 ^h 30 ^m 51.12 ^s	-33°34'9.2"	23 166 ± 97	907 ⁺⁸⁰ ₋₁₁₅	1.81	85	111	0.041	166	233	0.002
Abell 0389	2 ^h 51 ^m 28.80 ^s	-24°34'16.4"	34 085 ± 118	667 ⁺⁸⁰ ₋₁₁₅	1.33	26	24	0.632	45	55	0.173
Abell 3094	3 ^h 11 ^m 25.18 ^s	-26°32'40.0"	20 475 ± 77	774 ⁺⁶¹ ₋₇₀	1.55	89	110	0.084	140	233	<0.001
Abell 0957	10 ^h 13 ^m 49.80 ^s	-0°32'54.4"	13 623 ± 79	722 ⁺⁶³ ₋₇₃	1.44	77	95	0.084	115	149	0.065
Abell 1651	12 ^h 59 ^m 24.00 ^s	-4°68'0.4"	25 152 ± 107	817 ⁺⁸³ ₋₉₉	1.63	90	127	0.194	173	218	0.213
Abell 1750	13 ^h 30 ^m 49.68 ^s	-1°31'10.4"	25 647 ± 113	981 ⁺⁸⁸ ₋₁₀₄	1.96	101	168	<0.001	193	296	<0.001
Abell 2597	23 ^h 25 ^m 16.68 ^s	-12°44'6.4"	24 691 ± 117	597 ⁺¹¹⁷ ₋₉₀	1.19	21	16	0.529	36	37	0.276
Abell S1136	23 ^h 36 ^m 14.76 ^s	-31°21'56.4"	18 688 ± 92	617 ⁺⁸⁷ ₋₈₇	1.23	38	51	0.097	55	79	0.025




Figure 2 Results of applying the DS test to the dumbbell sample. In these bubble plots, each circle's radius is scaled proportional to the Dressler & Shectman (1988) δ statistic (i.e. proportional to the deviation of a localized group of galaxies mean velocity away from the whole cluster's mean velocity). Substructure is therefore indicated by spatially close and relatively big overlapping circles.

This is somewhat surprising as we may have naively expected there to have been recent cluster activity in these dumbbell cluster systems (see also Quintana et al. 1996) compared to the general cluster population. We re-emphasize the caveat of the DS test limitations: with some of our clusters (size ~ 30), the confidence level in the Δ statistic may drop to only 95% to be able to detect a 3:1 merger (Pinkney et al. 1996) — ~ 5 clusters at radii $< R_v$ and ~ 1 cluster at $< 2R_v$ in the dumbbell sample; and similar numbers in the control sample. Even with this confidence level and sample sizes, the data suggest that the incidence of 3:1 mass mergers is equal in both samples.

Given that Quintana et al. (1996) demonstrate that each component of the dumbbell in NGC 4782/3 system occupies different sub-groups, it is therefore a natural next step to ask whether the other galaxies in our samples also have large peculiar velocities?

4 Peculiar Velocities

Pimblet et al. (2006) have already noted that the dumbbell BCM galaxy from Abell 3653 has a very large peculiar velocity relative to the cluster frame (for both components of the dumbbell, Figure 1). Using the positions of




Figure 2 (Continued)

the individual dumbbell components given by Gregorini et al. (1994), we compare each of their redshifts with the mean cluster velocity from Table 1. The results of this are displayed in Table 3. The error listed on the velocity offset, Δcz , in this Table is simply the mean cluster velocity error (Table 1) added in quadrature to the galaxy velocity error given by NED (where more than 1 redshift measurement of a given dumbbell component is available, we choose the most recent since there is some uncertainty concerning which redshift may refer to what component in older measurements, e.g. Green, Godwin & Peach 1988). We perform a similar analysis on the control sample, identifying the BCM by eye and a magnitudinal comparison using NED (see Table 4).

For the control sample, we find that 3/14 (21%) of the clusters have BCMs with significant peculiar velocities (i.e. >3 standard errors away from $|\Delta cz| = 0$). Meanwhile in the dumbbell sample, 8/13 (62%) clusters have at least 1 dumbbell component with significant peculiar velocities (7 of which are large, $>300 \text{ km s}^{-1}$); but consistent to the control sample, only 3/13 (23%) have *both* dumbbell components with significant peculiar velocities. We note that for Abell 0533 the two dumbbell components have a very significant velocity offset ($>1600 \text{ km s}^{-1}$) which questions whether this system should be considered as a proper dumbbell system. Notwithstanding Abell 0533, we suggest that a cluster with a BCM dumbbell system is more likely to possess a significant peculiar velocity in at least

one of the dumbbell components than a non-dumbbell BCM cluster. Save for Abell 0533, no peculiar velocity is beyond 1.1 times the velocity dispersion measurement of the clusters, very much in line with the results of Valentijn & Casertano (1988).

5 Discussion and Conclusions

Even though the dumbbell sample is a collection of redshifts from many disparate sources, we believe that we have assembled a sufficient quantity of (bright) cluster members to adequately map out each of the clusters to a similar level found in 2dFGRS (see also Tables 1 and 2). Although the sample is likely not an optimal (homogeneously selected) one, we are nonetheless confident that the results presented would not change significantly with the addition of further observations. Indeed, subtracting a small percentage of galaxies from the better sampled clusters produces no significant change in the measured peculiar velocities, or substructuring.

Our results show that dumbbell BCM clusters are no more or less likely than other clusters to possess (3:1) subclustering (see also Oegerle & Hill 2001). However, they are more likely to have at least one BCM component with a significant peculiar velocity. We suggest the unifying interpretation of these observations is that our dumbbell BCM clusters may be in various states of post-merger activity. Those with both subclustering and peculiar velocities (e.g. Abell 3391) are probably in the early stages of

Table 3. Peculiar velocities of the dumbbells

Dumbbell component	RA (J2000)	Dec (J2000)	$ \Delta cz $ (km s ⁻¹)	Significance
Abell 2860 1	1 ^h 4 ^m 50.13 ^s	-39°2749.6''	539 ± 110	4.90
" " 2	1 ^h 4 ^m 50.11 ^s	-39°2749.8''	634 ± (97) ^a	(6.54) ^a
Abell 2911 1	1 ^h 26 ^m 5.42 ^s	-37°3472.9''	143 ± 120	1.19
" " 2	Uncertain	Uncertain	Uncertain ^b	-
Abell 3151 1	3 ^h 40 ^m 26.94 ^s	-28°2437.5''	376 ± 132	2.85
" " 2	3 ^h 40 ^m 25.14 ^s	-28°2439.0''	426 ± 128	3.33
Abell 3266 1	4 ^h 31 ^m 13.29 ^s	-61°1631.8''	247 ± 103	2.40
" " 2	4 ^h 31 ^m 12.18 ^s	-61°1635.1''	114 ± 101	1.13
Abell 0533 1	5 ^h 1 ^m 8.29 ^s	-22°2095.5''	255 ± 241	1.06
" " 2	5 ^h 1 ^m 6.63 ^s	-22°2096.2''	1906 ± 230	8.29
Abell 3391 1	6 ^h 26 ^m 20.22 ^s	-53°2494.0''	489 ± 133	3.68
" " 2	6 ^h 26 ^m 17.80 ^s	-53°2489.4''	68 ± 142	0.48
Abell 3528 1	12 ^h 54 ^m 23.40 ^s	-29°64.8''	100 ± 107	0.93
" " 2	12 ^h 54 ^m 22.32 ^s	-29°46.8''	0 ± 101	0.00
Abell 3532 1	12 ^h 57 ^m 21.96 ^s	-30°1308.9''	257 ± 64	4.02
" " 2	12 ^h 57 ^m 19.80 ^s	-30°1312.9''	36 ± 67	0.54
Abell 3535 1	12 ^h 57 ^m 55.44 ^s	-28°1720.0''	485 ± 92	5.27
" " 2	12 ^h 57 ^m 54.72 ^s	-28°1728.0''	462 ± 75	6.16
Abell 3570 1	13 ^h 46 ^m 47.28 ^s	-37°3268.4''	24 ± 111	0.22
" " 2	13 ^h 46 ^m 46.92 ^s	-37°3282.4''	49 ± 120	0.41
Abell 3653 1	19 ^h 53 ^m 3.48 ^s	-52°133.2''	736 ± 105	7.01
" " 2	19 ^h 53 ^m 2.76 ^s	-52°134.6''	495 ± 126	3.93
Abell 3716 1	20 ^h 52 ^m 0.48 ^s	-52°2718.0''	559 ± 92	6.08
" " 2	20 ^h 51 ^m 56.88 ^s	-52°2710.8''	255 ± 88	2.90
Abell 3744 1	21 ^h 7 ^m 25.68 ^s	-25°1543.3''	27 ± 94	0.29
" " 2	21 ^h 7 ^m 24.60 ^s	-25°1557.0''	211 ± 75	2.81

^aRedshift error not recorded for this component, the error quoted is simply the error on the cluster mean velocity and should therefore be taken as a lower bound.

^bUnable to unambiguously distinguish the second component and its redshift from the first component.

Table 4. BCM peculiar velocities in the control sample

Cluster	BCM RA (J2000)	BCM Dec (J2000)	$ \Delta cz $ (km s ⁻¹)	Significance
Abell 2734	0 ^h 11 ^m 21.67 ^s	-28°3075.8''	171 ± 101	1.69
Abell 2751	0 ^h 16 ^m 13.65 ^s	-31°1392.1''	245 ± 261	0.94
Abell 2755	0 ^h 17 ^m 40.99 ^s	-35°720.7''	284 ± 224	1.27
Abell 2798	0 ^h 37 ^m 32.20 ^s	-28°1915.9''	249 ± 116	2.15
Abell 2811	0 ^h 42 ^m 8.83 ^s	-28°1928.8''	23 ± 179	0.13
Abell 2829	0 ^h 51 ^m 22.53 ^s	-28°1897.5''	416 ± 126	3.30
Abell 3027	2 ^h 30 ^m 49.42 ^s	-33°371.8''	384 ± 132	2.91
Abell 0389	2 ^h 51 ^m 24.80 ^s	-24°3399.4''	331 ± 191	1.73
Abell 3094	3 ^h 11 ^m 25.00 ^s	-26°3351.9''	75 ± 92	0.82
Abell 0957	10 ^h 13 ^m 38.28 ^s	-0°3331.8''	253 ± 95	2.66
Abell 1651	12 ^h 59 ^m 22.56 ^s	-4°706.3''	419 ± 111	3.97
Abell 1750	13 ^h 30 ^m 50.76 ^s	-1°3097.0''	441 ± 116	3.80
Abell 2597	23 ^h 25 ^m 19.92 ^s	-12°446.0''	189 ± 118	1.60
Abell S1136	23 ^h 36 ^m 16.56 ^s	-31°2169.3''	19 ± 112	0.17

mixing: the bulk of the galaxies are separable in to sub-groups with a dumbbell component belonging to one (or even each) sub-grouping (see also Quintana et al. 1996). As the merger progresses, we may expect the less massive galaxies to homogenize first eventually leaving only

a large peculiar velocity in one or both dumbbell components as the singular signpost to recent cluster activity, before the individual components of the dumbbell BCM itself relax with respect to each other and the rest of the merged cluster (Abell 2911). Alternatively, we note that

intermediate clusters such as Abell 3570 may simply be the result of ellipticals preferentially being on radial orbits (e.g. Ramirez & de Souza 1998; see also Hwang & Lee 2008) and hence observationally indistinguishable. The dumbbell itself may be a final transient phase before ultimately merging into a more massive cD-like galaxy (e.g. Tremaine 1990; West 1994). These results and the emerging picture of the evolution of dumbbells with respect to cluster substructure are also consistent with other observational evidence that the timescale for clusters to accrete new galaxies is much shorter than the timescale required for the central galaxy to merge with the accreted galaxies (e.g. Cooray & Milosavljević 2005; Brough et al. 2008).

Acknowledgments

We thank the efforts of the 2dFGRS team and Anglo-Australian Observatory in creating the 2dFGRS survey. We also thank H. Andernach and I. G. Roseboom for fruitful discussion during the course of this work and the diligent work of the referee whose comments have improved this work. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

References

- Abell, G. O., 1958, *ApJS*, 3, 211
- Abell, G. O., Corwin, H. G., Jr. & Olowin, R. P., 1989, *ApJS*, 70, 1
- Bardelli, S., Zucca, E. & Baldi, A., 2001, *MNRAS*, 320, 387
- Brough, S., Couch, W. J., Collins, C. A., Jarrett, T., Burke, D. J. & Mann, R. G., 2008, *MNRAS*, 385, L103
- Caldwell, N. & Rose, J. A., 1997, *AJ*, 113, 492
- Christlein, D. & Zabludoff, A. I., 2003, *ApJ*, 591, 764
- Colless, M. et al., 2001, *MNRAS*, 328, 1039
- Collins, C. A., Guzzo, L., Nichol, R. C. & Lumsden, S. L., 1995, *MNRAS*, 274, 1071
- Cooray, A. & Milosavljević, M., 2005, *ApJ*, 627, L85
- da Costa, L. N., Pellegrini, P. S., Davis, M., Meiksin, A., Sargent, W. L. W. & Tonry, J. L., 1991, *ApJS*, 75, 935
- da Costa, L. N. et al., 1998, *AJ*, 116, 1
- Craddace, R. et al., 2002, *ApJS*, 140, 239
- Dalton, G. B., Efstathiou, G., Maddox, S. J. & Sutherland, W. J., 1994, *MNRAS*, 269, 151
- De Propris, R. et al., 2002, *MNRAS*, 329, 87
- Donnelly, R. H., Forman, W., Jones, C., Quintana, H., Ramirez, A., Churazov, E. & Gilfanov, M., 2001, *ApJ*, 562, 254
- Dressler, A. & Shectman, S. A., 1988, *AJ*, 95, 985
- Dubinski, J., 1998, *ApJ*, 502, 141
- Ebeling, H., Voges, W., Bohringer, H., Edge, A. C., Huchra, J. P. & Briel, U. G., 1996, *MNRAS*, 281, 799
- Ebeling, H., Barrett, E., Donovan, D., Ma, C.-J., Edge, A. C. & van Speybroeck, L., 2007, *ApJ*, 661, L33
- Fairall, A. P., 1988, *MNRAS*, 233, 691
- Fitchett, M. & Webster, R., 1987, *ApJ*, 317, 653
- Girardi, M., Biviano, A., Giuricin, G., Mardirossian, F. & Mezzetti, M., 1993, *ApJ*, 404, 38
- Girardi, M., Giuricin, G., Mardirossian, F., Mezzetti, M. & Boschin, W., 1998, *ApJ*, 505, 74
- Gregorini, L., Vettolani, G., de Ruiter, H. R. & Parma, P., 1992, *A&AS*, 95, 1
- Gregorini, L., de Ruiter, H. R., Parma, P., Sadler, E. M., Vettolani, G. & Ekers, R. D., 1994, *A&AS*, 106, 1
- Green, M. R., Godwin, J. G. & Peach, J. V., 1988, *MNRAS*, 234, 1051
- Hoessel, J. G. & Schneider, D. P., 1985, *AJ*, 90, 1648
- Hwang, H. S. & Lee, M. G., 2008, *ApJ*, 676, 218
- Jones, D. H. et al., 2004, *MNRAS*, 355, 747
- Kaldare, R., Colless, M., Raychaudhury, S. & Peterson, B. A., 2003, *MNRAS*, 339, 652
- Katgert, P., Mazure, A., den Hartog, R., Adami, C., Biviano, A. & Perea, J., 1998, *A&AS*, 129, 399
- Lacey, C. & Cole, S., 1993, *MNRAS*, 262, 627
- Laine, S., van der Marel, R. P., Lauer, T. R., Postman, M., O'Dea, C. P. & Owen, F. N., 2003, *AJ*, 125, 478
- Lauer, T. R., 1988, *ApJ*, 325, 49
- Loveday, J., Peterson, B. A., Maddox, S. J. & Efstathiou, G., 1996, *ApJS*, 107, 201
- Merritt, D., 1985, *ApJ*, 289, 18
- Oegerle, W. R. & Hill, J. M., 2001, *AJ*, 122, 2858
- Ostriker, J. P. & Tremaine, S. D., 1975, *ApJ*, 202, L113
- Paturel, G., Theureau, G., Bottinelli, L., Gouguenheim, L., Coudreau-Durand, N., Hallet, N. & Petit, C., 2003, *A&A*, 412, 57
- Pimblet, K. A., Roseboom, I. G. & Doyle, M. T., 2006, *MNRAS*, 368, 651
- Pinkney, J., Roettiger, K., Burns, J. O. & Bird, C. M., 1996, *ApJS*, 104, 1
- Postman, M. & Lauer, T. R., 1995, *ApJ*, 440, 28
- Quintana, H., Ramirez, A., Melnick, J., Raychaudhury, S. & Slezak, E., 1995, *AJ*, 110, 463
- Quintana, H., Ramirez, A. & Way, M. J., 1996, *AJ*, 112, 36
- Ramirez, A. C. & de Souza, R. E., 1998, *ApJ*, 496, 693
- Scaramella, R., Zamorani, G., Vettolani, G. & Chincarini, G., 1991, *AJ*, 101, 342
- Shectman, S. A., Landy, S. D., Oemler, A., Tucker, D. L., Lin, H., Kirshner, R. P. & Schechter, P. L., 1996, *ApJ*, 470, 172
- Schindler, S., 2000, *A&AS*, 142, 433
- Smith, R. J. et al., 2004, *AJ*, 128, 1558
- de Souza, R. & Quintana, H., 1990, *AJ*, 99, 1065
- Teague, P. F., Carter, D. & Gray, P. M., 1990, *ApJS*, 72, 715
- Tremaine, S., 1990, *Dynamics and Interactions of Galaxies*, 394
- Valentijn, E. A. & Casertano, S., 1988, *A&A*, 206, 27
- Vettolani, G. et al., 1998, *A&AS*, 130, 323
- Wegner, G., Colless, M., Saglia, R. P., McMahan, R. K., Davies, R. L., Burstein, D. & Baggle, G., 1999, *MNRAS*, 305, 259
- West, M. J., Oemler, A. J. & Dekel, A., 1988, *ApJ*, 327, 1
- West, M. J., 1994, *MNRAS*, 268, 79
- Woudt, P. A., Kraan-Korteweg, R. C., Lucey, J., Fairall, A. P. & Moore, S. A. W., 2008, *MNRAS*, 383, 445
- Yamada, T., Koyama, Y., Nakata, F., Kajisawa, M., Tanaka, I., Kodama, T., Okamura, S. & De Propris, R., 2002, *ApJ*, 577, L89
- Zabludoff, A. I., Huchra, J. P. & Geller, M. J., 1990, *ApJS*, 74, 1
- Zabludoff, A. I., Geller, M. J., Huchra, J. P. & Ramella, M., 1993, *AJ*, 106, 1301