

COMMONWEALTH AGRICULTURAL BUREAUX

Forthcoming Publication

Review of Biological Control of Pests in the Commonwealth Caribbean and Bermuda up to 1982

edited by M J W COCK

Contributors FD BENNETT, MJW COCK, IW HUGHES, FJ SIMMONDS and M YASEEN

Commonwealth Institute of Biological Control Technical Communication No. 9

Though biological control has been used in the Caribbean since the nineteenth century, it has been significantly intensified since the establishment of the West Indian Station in Trinidad, by the Commonwealth Institute of Biological Control, shortly after World War Two.

This review covers the practice of classical biological control of pests and weeds of both agricultural and human importance by the introduction of exotic natural enemies, and will form essential reading to plant protection workers as well as entomologists and weed researchers.

244 x 187mm, hardback, 244pp. (June 1985)

ISBN 0 85198 550 5

Obtainable from:

Commonwealth Agricultural Bureaux, (PRO. DPT. 504) Farnham House, Farnham Royal, Slough SL2 3BN, UK. Tel: Farnham Common (02814) 2662 Telex: 847964 Cables: Comag, Slough



Forthcoming Publication

COMMONWEALTH AGRICULTURAL BUREAUX

Biological Control Programmes against Insects and Weeds in Canada 1969-1980

edited by J S KELLEHER and M A HULME

Commonwealth Institute of Biological Control Technical Communication No. 8

This is the third book reviewing progress in biological control programmes against insect pests and weeds affecting agriculture and forestry in Canada. It is compiled from authoritative summaries written by research staff of the Canada Department of Agriculture and the Canadian Forestry Service engaged in the implementation of these programmes. Strict editorial control has ensured a uniformity of style and treatment of each programme so the book provides an invaluable record of what has been achieved which will be essential reading for all engaged in the control of pests and weeds in temperate regions, and an important source of information for all interested in the development of biological control as a major component of pest management.

Comparison with the two previous reviews demonstrates how the scope of biological control is widening with increased use of pathogens and a broadened ecological approach to the solution of major pest and weed problems.

Obtainable from:

Commonwealth Agricultural Bureaux, (PRO. DPT. 372) Farnham House, Farnham Royal, Slough SL2 3BN, UK, Tel: Farnham Common (02814) 2662 Telex: 847964 Cables: Comag, Slough

NOTES FOR AUTHORS

Papers submitted for publication in the Bulletin of Entomological Research should be sent, in duplicate, to the Director, Commonwealth Institute of Entomology, 56 Queen's Gate, London SW7 5JR. They must deal with original research concerning insects, mites or ticks of economic importance in the agricultural, medical or veterinary fields; they must be of more than purely local interest, and taxonomic papers will not be accepted unless the organisms dealt with are of definite importance in the field of applied entomology. Review articles may also be submitted, but will normally be by invitation. Papers must not have been published or accepted for publication elsewhere. They will normally be published in order of their receipt in definitive form.

For general guidance on presentation and style, including the preparation of illustrations, contributors are urged to consult *Writing Scientific Papers in English. An ELSE-Ciba Foundation Guide for Authors* (by Maeve O'Connor & E. Peter Woodford), 116 pp., Amsterdam, Associated Scientific Publishers, 1975, ISBN 90-219-4035-3.

Text

Papers should be typewritten (double spaced, with ample margins and on one side of the paper only) and a current number of the *Bulletin* should be consulted for details of style and lay-out. Titles should be as brief as is consistent with clarity. Abbreviations should conform to *British Standard 1991*, Part I, 1967, and are alike in singular and plural; stops will be used only where ambiguity might occur, e.g. 'in.' not 'in' for inch, but 'in²' for square inch. Metric units are preferred; if non-metric units are used, metric equivalents should be given at least once in each article.

Tables

Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must on no account be of a size or form that will not conveniently fit into the page of the *Bulletin*.

Illustrations

Line drawings and graphs must be in jet-black indian ink, preferably on bristol board or tracing cloth. Graphs may be drawn on paper ruled in pale blue or light grey; paper having green, red, yellow or brown lines should not be used. Any stippling or hatching, as well as any numerals or lettering, on the text-figures must be completed, by the author, in a form suitable for reproduction; this work is *not* undertaken by the printer. Text-figures, when reproduced, cannot exceed 12.7 cm in width; if, therefore, the original drawings are larger, the lettering on them must be in proportion, so that it is still legible when reduced; special attention should be paid to the relative scales of lettering and line thicknesses in relation to the figure as a whole. Photographs for reproduction as figures are admissible when they are a real contribution to the text. Captions to figures should be typed on a separate sheet.

Abstract

Each paper must contain a carefully prepared and accurate informative abstract that shall be complete in itself and intelligible without reference to the text or figures. It should not exceed 250 words, and should be included at the beginning immediately after the title, author and author's affiliation.

References

References must be based on the name and year system and conform to the present usages of the *Bulletin*. Titles of journals should be abbreviated as in the 4th edition of *The World List of Scientific Periodicals*. References to unpublished information should normally be included only in the text—(A. B. Jones, pers. comm.) or (A. B. Jones. Control of sandflies by insecticides. Unpublished report. Ross Institute, London, 1950).

Offprints

Free copies of offprints are allowed as follows: One author-20 copies; two or more authors-30 copies. Further copies may be obtained on payment, and the number required should be specified when the paper is submitted.

BULLETIN OF ENTOMOLOGICAL RESEARCH

(Word List Abbreviation: Bull. ent. Res.) ISSN 0007-4853

| HOLLOWAY, G. J. & SMITH, R. H. Inheritance of the ability of <i>Sitophilus oryzae</i> (L.) (Coleoptera: Curculionidae) to feed and breed on yellow split-pea (<i>Pisum sativum</i>) | PAGE 367 |
|---|-------------|
| BROOKES, C. P. & LOXDALE, H. D. A device for simultaneously homogenizing numbers of individual small insects for electrophoresis | 377 |
| SAMWAYS, M. J. Relationship between red scale, <i>Aonidiella aurantii</i> (Maskell) (Hemiptera: Diaspididae), and its natural enemies in the upper and lower parts of citrus trees in South Africa | 379 |
| COLLIER, R. H. & FINCH, S. Accumulated temperatures for predicting the time of emergence in the spring of the cabbage root fly, <i>Delia radicum</i> (L.) (Diptera: Anthomyiidae) | 395 |
| NYIRENDA, G. K. C. Persistent populations of males of the African armyworm, Spodoptera exempta (Walker) (Lepidoptera: Noctuidae), in Malawi | 405 |
| KING, A. B. S. Factors affecting infestation by larvae of <i>Phyllophaga</i> spp. (Coleoptera: Scarabaeidae) in Costa Rica | 417 |
| GOLDSON, S. L., DYSON, C. B., PROFFITT, J. R., FRAMPTON, E. R. & LOGAN, J. A. The effect of <i>Sitona discoideus</i> Gyllenhal (Coleoptera: Curculionidae) on lucerne yields in New Zealand | 429 |
| KIRK-SPRIGGS, A. H. <i>Meligethes heteropus</i> Gerstaecker (Coleoptera: Nitidulidae), a new pest of bulrush millet in West Africa | 443 |
| TORR, S. J. The susceptibility of <i>Glossina pallidipes</i> Austen (Diptera: Glossinidae) to insecticide deposits on targets | 451 |
| MARTIN, J. H. The whitefly genus <i>Dialeurolonga</i> (Hemiptera: Aleyrodidae) on citrus, with the redescription of a Malagasian species | 459 |
| CHARLWOOD, J. D., DAGORO, H. & PARU, R. Blood-feeding and resting behaviour in the Anopheles punctulatus Dönitz complex (Diptera: Culicidae) from coastal Papua New Guinea | 463 |
| BOWDEN, J., SHERLOCK, P. L. & DIGBY, P. G. N. Studies of elemental composition as a biological marker insects. III. Comparison of apterous and alate cereal aphids, especially <i>Rhopalosiphum padi</i> (L.) (Hemiptera: Aphididae), from oats and wheat, and from oats infected with or free from barley yellow dwarf virus | 477 |
| STEELE, G. M. & RANDOLPH, S. E. An experimental evaluation of conventional control measures against the sheep tick, <i>Ixodes ricinus</i> (L.) (Acari: Ixodidae). I. A unimodal seasonal activity pattern | 477 |
| RANDOLPH, S. E. & STEELE, G. M. An experimental evaluation of conventional control measures against the sheep tick, <i>Ixodes ricinus</i> (L.) (Acari: Ixodidae). II. The dynamics of the tick-host interaction | 501 |
| WARNES, M. L. & FINLAYSON, L. H. Responses of the stable fly, <i>Stomoxys calcitrans</i> (L.) (Diptera: Muscidae), to carbon dioxide and host odours. I. Activation | 519 |
| FLINT, S. A comparison of various traps for <i>Glossina</i> spp. (Glossinidae) and other Diptera | 529 |
| HUGHES, P. B. & RAFTOS, D. A. Genetics of an esterase associated with resistance to organophosphorus insecticides in the sheep blowfly, <i>Lucilia cuprina</i> (Wiedemann) (Diptera: Calliphoridae) | 535 |
| WILLIAMS, D. J. <i>Pseudococcus mandio</i> sp. n. (Hemiptera: Pseudococcidae) on cassava roots in Paraguay, Bolivia and Brazil | 545 |

© Commonwealth Agricultural Bureaux, 1985

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.