

Minimizers and Quasiminimizers

14.1 Quasiminimizers

In addition to currents and varifolds, there are several other ways to model minimal surfaces and related objects, see [139, 161]. Quasiminimizers provide a very natural and general setting for many variational problems. Let $E \subset \mathbb{R}^n$ be closed and unbounded such that for a fixed positive integer m , $0 < \mathcal{H}^m(E \cap B(x, r)) < \infty$ for $x \in E, r > 0$. We say that E is an m -quasiminimizer if for some $M < \infty$,

$$\mathcal{H}^m(E \cap W) \leq M \mathcal{H}^m(f(E \cap W))$$

for all Lipschitz mappings $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $W = \{x: f(x) \neq x\}$ is bounded. If this holds with $M = 1$, then E minimizes m -dimensional Hausdorff measure. The setting in the papers quoted below is more general. In particular, there is also a local, often very useful, version, but we skip it here. The quasiminimizers were introduced by Almgren in [9] under the name restricted sets. He proved that they are AD- m -regular and m -rectifiable. David and Semmes investigated them in [150]. They re-proved Almgren's results and went further. The following is a special case of their results:

Theorem 14.1 *If $E \subset \mathbb{R}^n$ is a closed m -quasiminimizer, then E is AD- m -regular, uniformly m -rectifiable and it contains big pieces of Lipschitz graphs (recall Section 5.2).*

Both Almgren's and David–Semmes's proofs use Lipschitz projections into k -dimensional cubical skeleta like in the Federer–Fleming proof of the deformation theorem of currents. First this gives AD-regularity. Then, by David and Semmes, via many complicated constructions, the big pieces of the Lipschitz graphs condition are verified.

The codimension 1 case was studied by different methods in [149] and [264].

All these papers contain many interesting results on and connections with various geometric variational problems.

There is much later work along these lines, see David's long paper [139] for a very general setting, for discussion and references. It seems to give the most general rectifiability results. In particular, there he used sliding conditions; the deformations were required to preserve given boundary pieces but were allowed to slide along them.

When minimizing Hausdorff measure the existence of minimizers is often a difficult question, both for the lack of lower semicontinuity and compactness. De Lellis, Ghiraldin and Maggi [162] established a general result to deal with this. For this they used Preiss's Theorem 4.11.

14.2 Mumford–Shah Functional

Let $\Omega \subset \mathbb{R}^n$ be a domain and g a bounded measurable function in Ω . The *Mumford–Shah functional* J is then defined by

$$J(u, K) = \int_{\Omega \setminus K} (u - g)^2 + \mathcal{H}^{n-1}(K) + \int_{\Omega \setminus K} |\nabla u|^2$$

for

$$(u, K) \in \mathcal{A}(\Omega) := \{(u, K) : K \subset \Omega \text{ relatively closed and } u \in W_{loc}^{1,2}(\Omega \setminus K)\}.$$

We assume that there are $(u, K) \in \mathcal{A}(\Omega)$ with $J(u, K) < \infty$, which is always true if $\Omega \subset \mathbb{R}^n$ is bounded. For many aspects of the Mumford–Shah functional, including applications to image segmentation and conjectures and results on minimizers, see the books [15] and [138]. Here we restrict the discussion to things related to rectifiability.

A minimizer for J is a pair $(u, K) \in \mathcal{A}(\Omega)$ which gives the smallest value for J . Minimizers always exist, although it is far from obvious since Hausdorff measure is not lower semicontinuous. One way to prove the existence is to first minimize

$$\int_{\Omega} (u - g)^2 + \mathcal{H}^{n-1}(S_u) + \int_{\Omega} |\nabla u|^2$$

for $u \in SBV(\Omega)$, recall Section 12.3. Minimizers for this exist by the compactness properties of SBV. To get from this a minimizer for J , the problem that S_u need not be closed has to be dealt with. Here one cannot use the full $BV(\Omega)$, since it would give 0 for the infimum. Anyway, now S_u is $(n - 1)$ -rectifiable by Theorem 12.13. This approach is discussed in [15]. In [138], a different approach without SBV is explained.

For a minimizer (u, K) , u is in $C^1(\Omega \setminus K)$, which follows from the fact that it solves the PDE $\Delta u = u - g$. For K there are conjectures which are only partially solved. David and Semmes proved the following in [148], see also [138]:

Theorem 14.2 *If (u, K) is a minimizer for J and $B(x, 2r) \subset \Omega$, then $K \cap B(x, r)$ is contained in an AD- $(n-1)$ -regular uniformly $(n-1)$ -rectifiable set.*

The key to the proof is that the failure of the Poincaré inequality in the complement of an AD- $(n-1)$ -regular set E at most scales implies uniform rectifiability of E . This is understandable because the validity of the Poincaré inequality requires that E does not separate the space too much. More precisely: E is uniformly $(n-1)$ -rectifiable if there exists a positive number c such that for all $M \geq 1$ the set $F(E, c, M)$ of pairs (x, r) , $x \in E$, $0 < r < d(E)$, satisfying the following condition, is a Carleson set: for all balls $B(x_i, r_i) \subset B(x, r) \setminus E$, $i = 1, 2$, with $r_i > cr$ and for all $f \in W^{1,1}(B(x, Mr) \setminus E)$,

$$\left| r_1^{-n} \int_{B(x_1, r_1)} f - r_2^{-n} \int_{B(x_2, r_2)} f \right| \leq Mr^{1-n} \int_{B(x, r) \setminus E} |\nabla f|. \quad (14.1)$$

David and Semmes proved this by showing that this condition implies the local symmetry of Theorem 5.9. Another proof is described in [138]. The converse is false; an example is a coordinate hyperplane with the balls of radius $1/10$ centred in the integer lattice removed.

For slight simplicity, assume $\Omega = \mathbb{R}^n$. To prove that for a minimizer (u, K) the set $F(K, c, M)$ is a Carleson set, one applies (14.1) with $u = f$ and constructs a competitor to get for some $p < 2$,

$$\omega_p(x, Mr) = r^{p/2-n} \int_{B(x, Mr) \setminus K} |\nabla u|^p > \varepsilon(M) > 0.$$

As $r^{1-n} \int_{B(x, r) \setminus K} |\nabla u|^2$ is bounded, it is not very difficult to prove that the set of (x, r) such that $\omega_p(x, r) > \varepsilon$ satisfies a Carleson condition, from which it follows that $F(K, c, M)$ is a Carleson set.

Theorem 14.2 holds for a much larger class of quasiminimizers.

14.3 Some Free Boundary Problems

In [141], David, Engelstein and Toro studied the following two-phase free boundary problem. Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and q_+ and q_- bounded continuous functions on Ω . Let

$$J(u) = \int_{\Omega} (|\nabla u(x)|^2 + q_+(x)^2 \chi_{\{u>0\}}(x) + q_-(x)^2 \chi_{\{u<0\}}(x)) \, dx.$$

Among other things they proved that if u is an almost minimizer (we omit the definition) for J , then, under slight extra conditions, the sets $\Omega \cap \partial\{x \in \Omega: u(x) > 0\}$ and $\Omega \cap \partial\{x \in \Omega: u(x) < 0\}$ are locally AD- $(n-1)$ -regular and uniformly $(n-1)$ -rectifiable. The proof is a complicated mixture of potential theory and geometric measure theory. In particular, proving the AD-regularity is quite demanding and achieved with estimates for the harmonic measure.

We shall return to the corresponding one-phase problem in Section 15.6.

Rigot [390] proved the uniform rectifiability of sets almost minimizing perimeter, recall Section 12.1. Let $g(0, \infty) \rightarrow (0, \infty)$ with $g(x) = o(x^{(n-1)/n})$.

Theorem 14.3 *Let $E \subset \mathbb{R}^n$ be Lebesgue measurable. If*

$$P(E) \leq P(F) + g(\mathcal{L}^n((E \setminus F) \cup (F \setminus E)))$$

whenever $F \subset \mathbb{R}^n$ is Lebesgue measurable and $F = E$ outside some compact set, then E is equivalent to E' for which $\partial E'$ is AD- $(n-1)$ -regular and uniformly $(n-1)$ -rectifiable.

She proved this by showing that $\partial E'$ is a Semmes surface, recall Section 8.7.