British Journal of Nutrition (2011), **106**, 1763 © The Author 2011

Letter to the Editor

Comment on Jia et al.

(First published online 21 September 2011)

Jia et al. (1) report that dietary fish oil or curcumin, given alone, promotes mucosal inflammation and damage, whereas, given together, resolution of inflammation was promoted and major inflammatory factors, especially NF-kB, were suppressed. These findings are suggested to reflect an interaction of dietary lipids and curcumin in mucosal homeostasis and in the control of colonic inflammation through the suppression of PG release in colonic mucosa by both compounds through the Toll-like receptor 4/cyclo-oxygenase-2/PGE signalling axis. However, these effects might also arise through the actions of these two compounds on vitamin D signalling through vitamin D receptors (VDR), since both bind to the VDR⁽²⁾. DHA competes with calcitriol for binding to the VDR, as do other PUFA and biliary lithocholic acid, which may tend to reduce VDR signalling activity, while curcumin can have calcitriollike effects since its binding to the VDR induces some genomic effects of the activated VDR and it can also induce rapid nongenomic VDR-mediated effects such as activation of chloride channels⁽²⁾. Vitamin D reduces inflammation generally, including in the colonic mucosa, and is activated in the gut as in most target tissues (3,4). Furthermore, pharmacological preparations of fish oil contain small amounts of vitamin D but large amounts of vitamin A, which antagonises many actions of vitamin D. For example, vitamin A reduces vitamin Dinduced absorption of dietary Ca⁽⁵⁾. It would be of interest to know whether the reported effects and interactions of fish oil and curcumin on colonic inflammation are related to variations in vitamin D intake (or to vitamin D repletion as judged by serum 25-hydroxyvitamin D concentration) or to variations in vitamin A intake, or both.

Barbara J. Boucher

Blizard Institute
Centre for Diabetes
Barts and The London School of Medicine
and Dentistry
Queen Mary University of London
2 Newark Street
London E1 2AT
UK
email bboucher@doctors.org.uk

doi:10.1017/S0007114511005368

References

- 1. Jia Q, Ivanov I, Zlatev ZZ, *et al.* (2011) Dietary fish oil and curcumin combine to modulate cytogenetics and gene expression in dextran sodium sulphate treated rats. *Br J Nutr* **106**, 519–529.
- Haussler MR, Haussler CA, Bartik L, et al. (2009) Vitamin D receptor: molecular signalling and actions of nutritional ligands in disease prevention. Nutr Rev 66, S98–S112.
- Adams JS & Hewison M (2008) Unexpected actions of vitamin D: new perspectives on the regulation of innate and adaptive immunity. Nat Clin Pract Endocrinol Metab 4, 80–90.
- Hopkins MH, Owen J, Ahearn TU, et al. (2011) Effects of supplemental vitamin D and calcium on biomarkers of inflammation in colorectal adenoma patients: a randomized controlled clinical trial. Cancer Prev Res (Phila) (epublication ahead of print version 30 June 2011).
- Johansson S & Melhus H (2001) Vitamin A antagonizes calcium responses to vitamin D in man. J Bone Miner Res 16, 1899–1905.