NOTES FOR AUTHORS

The Journal of Helminthology publishes papers on all aspects of helminths, particularly those of medical or veterinary importance. Taxonomic contributions will be acceptable if they contribute to the systematics of a group and particularly if they employ biochemical or molecular biological techniques. Short reviews will also be welcome.

Page Format. The Journal is printed in a two-column format (column width of 80 mm) with a text area of 170×225 mm.

Text. Papers should be typed, on one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used.

When the paper has been accepted word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with fewer type-setting errors.

Abstract. Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words. A short title should be provided as a running head.

Tables. Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or landscape (if absolutely necessary) format. Table captions should be typed on a separate sheet.

Illustrations. Copies only of artwork should be submitted. The original illustrations should accompany the paper after acceptance and Text revision. figures, line drawings, computer-generated figures and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. They should be glossy prints of the same size as they are to appear in the Journal. All figures and letters on

photographs must be inserted by the author. Figure and captions should be typed on a separate sheet.

Voucher specimens. The deposition of voucher specimens should be considered where appropriate.

References. References must be based on the name and year system, give full journal titles and conform to the following styles:

- Grønvold, J., Wolstrup, J., Larsen, M., Henriksen, S.A. & Nansen, P. (1993) Biological control of Ostertagia ostertagi by feeding selected nematode-trapping fungi to calves. Journal of Helminthology 67, 31–36. Grove, D.I. (1990) <u>A history of human helminthol-</u>
- ogy. 850 pp. Wallingford, CAB International.
- Southgate, V.R. & Rollinson, D. (1987) Natural history of transmission and schistosome interactions. pp. 347-378 in Rollinson, D. & Simpson, A.J.G. (Eds) The biology of schistosomes: from genes to latrines. London, Academic Press.

Citation of authors in the text should appear in the form: Polaszek (1990) or (Polaszek, 1990). More than one author should be cited in chronological order as: (Holloway et al., 1987; Walker & Huddleston, 1988).

Offprints. 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. Three copies of the manuscript, which must be in English or French (with an English summary) should be accompanied by a letter signed by all the authors and together with artwork submitted to:

The Editor Journal of Helminthology International Institute of Parasitology 395A Hatfield Road St Albans, Herts AL4 0XU, UK.

SEPTEMBER 1994

Journal of Helminthology

Research Papers

| Agatsuma, T., Terasaki, K., Yang L. & Blair, D. Genetic variation in the triploids of Japan Fasciola species, and relationships with other species in the genus | |
|--|-----|
| | 181 |
| Behnke, J.M., Pritchard, D.I., Wakelin, D., Park, J.R., McNicholas, A.M. & Gilbert, F.S. | |
| Effect of ivermectin on infection with gastro-intestinal nematodes in Sierra Leone | 187 |
| Díez-Baños, P., Morrondo-Pelayo, P., Feijoo-Penela, A., Carrillo-González, B. & | |
| López-Sández, C. Relationship between the excretion of protostrongylid larvae in sh North-west Spain and climatic conditions | 197 |
| Kaufman, A.R. & Fried, B. Infectivity, growth, distribution and fecundity of a six versus | |
| twenty-five metacercarial cyst inoculum of Echinostoma caproni in ICR mice | 203 |
| Manga-González, M.Y. & Morrondo-Pelayo, M.P. Larval development of ovine Neostrongylus | |
| linearis in four experimentally infected mollusc species | 207 |
| Mattison, R.G., Hanna, R.E.B. & Nizami, W.A. Ultrastructure and histochemistry of the | |
| tegument of juvenile paramphistomes during migration in Indian ruminants | 211 |
| Mendoza-de Gives, P., Zavaleta-Mejia, E., Herrera-Rodriguez, D. & Quiróz-Romero, H. In vitro trapping capability of Arthrobotrys spp. on infective larvae of Haemonchus contorts and Nacobbus aberrans | 223 |
| Olmeda-García, A.S. & Rodríguez-Rodríguez, J.A. Stage-specific development of a filarial nematode (Dipetalonema dracunculoides) in vector ticks | 231 |
| O'Lorcain, P. Prevalence of Toxocara canis ova in public playgrounds in the Dublin area of Ireland | 237 |
| Papazahariadou, M.G., Koutinas, A.F., Rallis, T.S. & Haralabidis, S.T. Prevalence of | |
| microfilaraemia in episodic weakness and clinically normal dogs belonging to hunting breeds | 243 |
| Roots, C.D., Lewis, J.W. & Churchfield, J.S. The morphology of hymenolepidid and dilepidid cestodes from common and pygmy shrews (Soricidae) in Southeast England | 247 |
| a rumen amphistome Cotylophoron cotylophorum | n |
| Sun, S. & Sugane, K. Complete structure of the gene encoding an immunodominant antigen of Dirofilaria immitis and larva-specific synthesis of primary transcript | 259 |
| Research Notes | |
| Fan, P.C., Chung, W.C. & Wu, J.C. Experimental infection of an isolate of <i>Taenia solium</i> from Hainan in domestic animals | 265 |
| Ito, A., Nakao, M., Ito, M., Matsuzaki, T., Kamiya, M. & Kutsumi, H. Antibody responses in the wild vole, <i>Clethrionomys rufocanus bedfordiae</i> , naturally infected with <i>Echinococcus multilocularis</i> by Western blotting | 267 |
| Negita, T. & Ito, A. In vitro hatching of oncospheres of Taenia taeniaeformis using eggs isolated | |
| from fresh, frozen, formalin-fixed and ethanol-fixed segments | 271 |

C CAB INTERNATIONAL, 1994

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.