

Crops and Soils Research Paper

Cite this article: Alan O, Budak B, Sen F, Ongun AR, Tepecik M, Ata S (2025). Solid and liquid digestate generated from biogas production as a fertilizer source in processing tomato yield, quality and some health-related compounds. *The Journal of Agricultural Science* **163**, 55–70. <https://doi.org/10.1017/S0021859624000741>

Received: 25 June 2024
Revised: 9 November 2024
Accepted: 6 December 2024
First published online: 20 January 2025


Keywords:

blossom-end rot; Brix yield; digestate application; lycopen; *Lycopersicon esculentum* Mill

Corresponding author:

Ozlem Alan;
Email: ozlem.alan@ege.edu.tr

Solid and liquid digestate generated from biogas production as a fertilizer source in processing tomato yield, quality and some health-related compounds

Ozlem Alan¹ , Bulent Budak¹, Fatih Sen², Ali Riza Ongun³, Mahmut Tepecik³ and Samet Ata⁴

¹Odemis Vocational Training School, Ege University, Izmir, Turkey; ²Department of Horticulture, Faculty of Agriculture, Ege University, Izmir, Turkey; ³Department of Soil Science and Plant Nutrition, Faculty of Agriculture, Ege University, Izmir, Turkey and ⁴SÜTAŞ Dairy Products Inc, Izmir, Turkey

Abstract

To manage anaerobic digestion residues (digestate) sustainably, it is important to determine their agricultural properties. In the present study, the effects of two digestate fractions (solid and liquid) on processing tomato yield parameters, quality traits, health-related compounds and some fruit physiological disorders were evaluated. The solid and liquid digestate fractions were compared with chemically fertilized and unfertilized control to evaluate the potential of the digestate as a fertilizer. A 2-year experiment was conducted in a randomized complete block design, with three replications and using two tomato varieties: cv. 'Arte F₁' and cv. 'Zeplin F₁'. The results indicated that (1) compared with chemical fertilizer, the solid digestate produced equal or even better results in terms of fruit size, yield parameters (solid digestate treatment increased the total fruit weight per plant by an average of 30.7, 8.2 and 22.4% in 2019 and 25.3, 14.2 and 17.9% in 2022 compared with control, chemical fertilizer and liquid treatments, respectively) and percentage of fruit affected by sunscald and blossom-end rot in both years; (2) use of liquid digestate led to similar or significantly higher fruit size, yield parameters and percentage of fruit affected by sunscald and blossom-end rot than control in both years and (3) use of both solid and liquid digestate fractions significantly maintained or improved fruit quality in terms of colour traits, pericarp thickness, dry matter content, total soluble solid content, titratable acidity, pH, vitamin C and antioxidant activity. However, the effects of solid and liquid digestate fractions varied with year and variety.

Introduction

Global dependence on fossil fuels; depletion of energy-producing raw materials, such as oil, coal and natural gas; and growing energy demands have forced countries to actively search for alternative energy sources, such as renewable energy, which are environmentally friendly (Panwar *et al.*, 2011). One of the most important alternative energy sources is anaerobic biomass digestion. Many studies have highlighted that the production of biogas through anaerobic digestion (AD) offers additional advantages (socio-economic, technological and environmental opportunities) over other forms of renewable energy production (Rocha-Meneses *et al.*, 2023).

AD effectively converts biowaste into two valuable economic by-products: (1) biogas, a sustainable energy source and (2) anaerobic digestate, sometimes known as 'digestate', which can be used to fertilize and amend soil (Nkoa, 2014). AD produces clean energy from waste materials (e.g. food waste, animal manure and agricultural waste and residues) while reducing environmental risk from untreated waste (Bhatt and Tao, 2020; Lee *et al.*, 2021). However, AD faces challenges in sustainably managing digestive residues. According to Czekala (2022), excessive digestate production may occur as a result of intensive biogas plants, leading to additional environmental problems if this extra digestate is not well managed (Antoniou *et al.*, 2019).

AD preserves large quantities of nutrients from the feedstock, resulting in a nutrient-rich digestate that serves as a fertilizer for plants. Digestates are a complex mixture of organic compounds, minerals (macroelements and microelements) and microorganisms and can affect crop growth, yield and quality (Bolzonella *et al.*, 2018). Albuquerque *et al.* (2012) indicated that agrochemical characteristics of digestates vary depending on substrate composition (feedstock) and digestion type. AD can be used as a fertilizer for various species. Doyeni *et al.* (2021) reported that the digestate increased the grain yield and quality of *Triticum aestivum* L. compared with a synthetic N fertilizer. Moreover, Koszel *et al.* (2020) reported that digestate

treatment in winter rapeseed increased the yield as well as protein, macronutrient and saturated and unsaturated fatty acid contents compared with the control. Panuccio *et al.* (2019) emphasized that digestates increased phenol and flavonoid contents as well as antioxidant activity (AA) in cucumber compared with the unfertilized control. Finally, Barzee *et al.* (2019) reported that tomatoes fertilized using digestate had a higher soluble solid content compared with those fertilized using a synthetic fertilizer.

Digestate can be separated into solid and liquid fractions, which have different physical and chemical properties (Yu *et al.*, 2021). Thus, the solid and liquid fractions of digestates vary from each other. This variation can be attributed to the origin of substrates, operating parameters used in the digester and the type of solid–liquid separation used (Akhiar *et al.*, 2017). Defined attributes, such as dry or organic matter content, NH_4 content, C/N ratio and nutrient content of solid or liquid fractions that form the digestates, may have different efficiencies in plants and soils.

Conversely, chemical fertilization is one of the most widely used regimes in intensified agriculture (Hernández *et al.*, 2014). Although chemical fertilization is an effective regime for obtaining higher crop yields, its excessive and long-term use is expensive and causes several problems, such as degradation of soil fertility and soil microbial population, erosion, soil acidity and ground-water pollution (Shen *et al.*, 2021; Carricondo-Martínez *et al.*, 2022). The use of AD may result in additional benefits. Many studies have reported that the application of digestate has positive effects on soil physical, chemical and biological properties, which are key factors for soil functioning. Mayerová *et al.* (2023) reported that the long-term application of digestate improves bulk density, soil aggregate stability and water infiltration, and Makádi *et al.* (2016) indicated that the consecutive application of liquid digestate decreases the soil pH, despite the alkaline property of the applied digestate, especially in soils with low buffering capacity. Moreover, the addition of digestate into soil increases microbial biomass and activity of several enzymes involved in C and N cycles, which depend on digestate carbon and nutrient contents (Barra Caracciolo *et al.*, 2015; Van Midden *et al.*, 2023). Pan *et al.* (2018) reported that the application of digestate may suppress soil-borne plant diseases caused by antagonistic bacteria.

Tomato (*Lycopersicon esculentum* Mill.) is one of the most important industrial crops and accounts for a global production of 189.1 million tonnes harvested from approximately 5.2 million ha (FAOSTAT, 2022). More than 80% of the tomatoes cultivated globally are processed into sauce, juice, ketchup, canned tomato, stew, soup, etc., which have high nutritional qualities (Petropoulos *et al.*, 2020). Plants, particularly processing tomatoes, must have unique traits to improve production and fruit quality with respect to the processing sector while meeting the quality criteria of the market (Moura and Golynski, 2018).

Studies have investigated the use of digestate as a fertilizer for the tomato plant in greenhouses (under controlled environment) (Zheng *et al.*, 2019; Bergstrand *et al.*, 2020; Cristina *et al.*, 2020; Li *et al.*, 2023) and under field conditions (Yu *et al.*, 2010; Ferdous *et al.*, 2018; Morra *et al.*, 2021; Li *et al.*, 2023). In addition, studies have referred to different forms of digestates, such as pelleted, concentrated biogas slurry, liquid or solid fractions and biochar. The only data available on the two digestate fractions (solid and liquid) of tomato, which was grown in a greenhouse and used plastic pots (Panuccio *et al.*, 2021), are different from those obtained under open field conditions. Although these experiments provide a basis for the use of digestate as a fertilizer for tomato, more research is needed to determine the use of digestate

as a fertilizer under different treatment and environmental conditions. Furthermore, several studies have emphasized on tomato fruit quality and health-related traits, which can vary on the basis of genotype (Chea *et al.*, 2021) or genotype in interaction with the environment (Panthee *et al.*, 2013). Considering these findings, our study aimed to (1) compare the effects of the solid and liquid digestate fractions with those of chemical fertilizer (CF) on yield-quality traits and health-related compounds in processing tomato in a 2-year experiment conducted in an open field setting, (2) assess the fertilizer value of the two digestate fractions by accounting for the two genotypes of processing tomatoes and (3) evaluate new fertilization regimes in which AD can be used as the main source of fertilizer and sustainably produce similar yields and quality characteristics as CFs.

Materials and methods

Plant material and crop management

During 2019 and 2022, field experiments were conducted in the experimental fields of the Odemis Vocational School at Ege University, Izmir, Turkey (latitude 38°12'N, longitude 27°52'E and altitude 111 m a.s.l.). Field trials of the project were planned for two consecutive years. However, due to the COVID-19 pandemic in 2020 and 2021, the second year of field trials was performed in 2022. Two varieties of processing tomatoes were used as plant material: cv. 'Arte F₁' (May Seed Company, Turkey) and cv. 'Zeplin F₁' (Vilmorin Seed Company, France). These varieties are widely grown for the commercial production of tomato in Turkey. Tomato seedlings were obtained from a commercial nursery.

Soil samples were obtained from a depth of 0–30 cm before transplantation. The physical and chemical properties of the soil are listed in Table 1. The air temperature and mean total rainfall recorded during the cropping cycles (April–August) were 42–0.8°C and 24.4 mm in 2019 and 37.9–4.3°C and 27.1 mm in 2022, respectively (Fig. 1).

Solid digestate (SD) and liquid digestate (LD) fertilizers were obtained from the biogas plants of ENFAŞ I.C., a subsidiary of SÜTAŞ I.C., operating in the Izmir-Tire district of Turkey. The raw materials used in AD were manure (65%) (50% cow manure + 15% poultry/chicken manure), agricultural waste (30%) and dairy waste (5%). The heavy metal contents of SD and LD were below the values permitted by Turkish directives (RG, 2010). The chemical characteristics of SD and LD in 2019 and 2022 are shown in Table 2. The nutrient content of SD in 2019 and 2022 was as follows: total nitrogen (3.3 v. 1.1%), phosphorous (P_2O_5 , 1.97 v. 0.54%), potassium (K_2O , 2.1 v. 0.91%) and organic matter (81.3 v. 86.6%). The nutrient content of LD in 2019 and 2022 was as follows: total nitrogen (0.8 v. 0.6%), phosphorous (P_2O_5 , 0.48% v. not detected), potassium (K_2O , 1.15 v. 0.5%) and organic matter (6.6 v. 7.6%).

Field experiments were conducted under four treatment regimes: (1) control (C), unfertilized plants (null control); (2) CF, fertilization with 150 kg N/ha of NPK fertilizer; (3) SD and (4) LD. Composite fertilizer for CF; and experimental materials for SD and LD were homogeneously spread on the soil surface. Then, all experiment plots were tilled at a depth of 0–15 cm with a rototiller 1 week before tomato seedlings were transplanted into the open field (within the last week of April in both years). The seedlings were transplanted at a density of 2.75 plants/m² in a single row with a spacing of 1.40 m between each row and 0.26 m between plants in the row. A randomized complete

Table 1. Physical and chemical properties of the soil in the 2-year experiment

Properties	2019	2022	Properties	2019	2022
pH ^a (1:2.5)	7.31	7.23	Available P ^g (mg/kg)	7.04	11.0
Total salt ^b (%)	0.05	0.03	Available K ^h (mg/kg)	452	350
CaCO ₃ ^c (%)	2.80	1.11	Available Ca ^h (mg/kg)	540	480
Sand (%)	70.1	76.9	Available Mg ^h (mg/kg)	145	155
Clay (%)	8.60	6.78	Available Fe ⁱ (mg/kg)	4.72	4.05
Silt (%)	21.4	16.4	Available Zn ⁱ (mg/kg)	1.18	1.11
Texture ^d	Sandy loam	Sandy loam	Available Mn ⁱ (mg/kg)	10.2	10.6
Organic matter ^e (%)	1.37	1.24	Available Cu ⁱ (mg/kg)	0.42	0.85
Total N ^f (%)	0.084	0.072			

a, 1:2.5 water extract; b, 1:2.5 soil: conductimetric in water extract; c, calcimetric; d, hydrometric; e, Walkley-Black method; f, Kjeldahl method; g, available Olsen; h, available 1 N NH₄OAc extract; i, available DTPA extract.

block design with three replications was used. The experimental plots were 28 m² in area (5.6 m × 5.0 m).

For one tomato crop cycle, 150 kg/ha N was used, considering the nitrogen requirement of tomato plants. The amounts of SD and LD used were calculated considering N dosage (150 kg/ha). The moisture and nutrient contents of SD and LD differed in the 2 years of the experiment (Table 2). Therefore, the application rates of SD and LD in the second year were established according to the amount of N applied in the first year (SD, 16.66 tonne/ha in 2019 and 49.8 tonne/ha in 2022; LD, 21.16 L/ha in 2019 and 28.14 L/ha in 2022). For basic chemical fertilization, zinc-enriched composite fertilizer (15% N, 15% P₂O₅ and 15% K₂O + Zn) was used. In addition to basic fertilization, first dressing (side dressing) was applied in May and urea (46% N) was used as N source. The second dressing was applied in June and calcium ammonium nitrate (26% N) and potassium nitrate (13-0-46) were used as N sources. Drip irrigation and plant protection from pests were managed according to the farm standard.

Yield assessment

A single harvest was performed at the end of the growing seasons in each year – within the first weeks of August 2019 and 2022.

When the tomato fruits reached harvest maturity (ripe fruits accounting for approximately 85% of the total yield), eight plants from the centre of each replication were harvested randomly by hand in the morning. Then, the plants were taken to the processing lab and fruits were graded as red (ripe), green (unripe), unmarketable and affected by sunscald (SS) and blossom-end rot (BER). The mean fruit weight (MFW) was calculated using marketable red fruits, number of fruits per plant was calculated using red and green fruits, and TFW was calculated using marketable red fruits. Brix yield (t/ha) was calculated by dividing the marketable yield (t/ha) by total soluble solid (TSS) content, as described by Duman *et al.* (2003). Morphology, chemical composition and health-related compounds were determined using ripe fruits (20 fruits for each parameter).

Physical analysis of fruits

In total, 20 tomato fruits for each variety in each replication were used to determine MFW using a precision scale (XB 12100; Presica Instruments Ltd., Switzerland) as well as fruit diameter (FD), fruit length (FL) and fruit pericarp thickness (PT) using a digital compass (SC-6; Mitutoyo, Japan). The results were expressed in mm.




Figure 1. Air temperature and total rainfall recorded during the cropping cycles (April–August) in the two growing seasons (2019 and 2022) (T-Min, minimum temperature; T-Max, maximum temperature; T-Mean, monthly mean temperature).

Table 2. Main chemical characteristics of SD and LD used in the 2-year experiment

Parameters	SD		LD	
	2019	2022	2019	2022
pH	7.4	8.7	7.9	8.5
EC (dS/m)	6.03	1.27	4.7	3.3
Organic matter (%)	81.3	86.6	6.6	7.6
Total (humic + fulvic) acid (%)	42.9	39.1	5.8	5.72
Total nitrogen (%)	3.3	1.1	0.8	0.6
P ₂ O ₅ (%)	1.97	0.54	0.48	Not detected
K ₂ O (%)	2.1	0.91	1.15	0.5
Zn	286 mg/kg	192 ppm	128.1 mg/kg	0.002 (%)
Cu	35.01 mg/kg	19.5 ppm	9.82 mg/kg	Not detected
Cd	1.61 mg/kg	0.13 ppm	0.35 mg/kg	<0.01 (RL)
Pb	5.52 mg/kg	<0.01 (RL)	1.29 mg/kg	<0.01 (RL)
Cr	19.9 mg/kg	10.74	2.98 mg/kg	1.6 ppm
Ni	11.8 mg/kg	7.06	2.21 mg/kg	1.6 ppm
Hg	<0.01 (RL)	<0.01 (RL)	<0.01 (RL)	<0.01 (RL)
Sn	<0.01 (RL)	<0.01 (RL)	<0.01 (RL)	<0.01 (RL)

RL, reporting limit; SD, solid digestate; LD, liquid digestate.

The peel colour was measured at the equatorial level on both sides of 20 plums with a colorimeter (CR-400; Minolta Co., Osaka, Japan), and the average scores were recorded in terms of CIE $L^* a^* b^*$ values. Chroma (C^*), which indicates intensity or colour saturation, and hue angle (h°), which was expressed as follows: 0° (red-purple), 90° (yellow), 180° (bluish-green) and 270° (blue) (McGuire, 1992), were calculated by using the following equation:

$$C^* = [a^{*2} + b^{*2}]^{1/2} \quad (1)$$

$$h^\circ = \tan^{-1} [b^*/a^*] \quad (2)$$

The a^*/b^* ratio can be used to report the brightness of the red colour of tomato fruit and its products (Akdeniz *et al.*, 2012). Colour measurement in fruit pulp samples was conducted using the same method.

Fruit firmness (FF) was determined for 20 tomato fruits per replicate using a texture analyser (GS-15, GÜSS Manufacturing Ltd., South Africa) by plunging the 7.9 mm-diameter tip to 10 mm depth with 10 cm/min speed after removing the skin from the equatorial section of the fruit. The results were expressed in Newton (N).

Chemical analysis of fruits

The TSS content of tomato juice was determined using a digital refractometer (PR-1; Atago, Tokyo, Japan) and expressed in percentage. Dry matter (DM) content was determined by drying the samples in an oven (Memmert, Germany) at 65°C until a constant weight was obtained and calculated on the basis of the percentage of weight loss (AOAC, 1990). Titratable acidity (TA) was determined by titrating 10 ml of the juice with 0.1 N NaOH up to

pH 8.1. The results were expressed as gram citric acid per 100 ml of fruit juice. pH value was determined using a pH metre (MP220, Mettler Toledo, Germany) (Karaçalı, 2012).

Lycopene content, vitamin C content, TPC and AA

The amount of lycopene, i.e. the colour obtained from an extract of tomato sample treated with acetone solvent and homogenized, was measured using a spectrophotometer at 503 nm. The results were expressed as mg/kg and calculated using the following formula: lycopene (mg/kg) = $62.43 \times \text{OD}_{503}/\text{sample weight}$ (Davis *et al.*, 2003).

A sample of 50 g tomatoes was mixed with 50 ml of oxalic acid (0.4%), blended in a Waring blender (Blender 8011ES, USA) and filtered. Vitamin C (L-ascorbic acid) content in the filtrate was measured using 2,6-dichloroindophenol according to a titrimetric method (AOAC, 1995). The results were expressed as mg L-ascorbic acid per 100 g fresh FW.

Fruit extracts (in methanol) were prepared from tomato fruits according to the method described by Thaipong *et al.* (2006), with some modifications for TPC and AA analysis. TPC analysis was conducted using the Folin–Ciocalteu method according to Zheng and Wang (2001), with an incubation time of 120 min for colour development. Absorbance was measured at 725 nm using a spectrophotometer (Carry 100 Bio; Varian, Australia) and the results were expressed as milligram gallic acid equivalent (GAE) per 100 g of FW with reference to a standard curve of gallic acid (0–0.1 mg/ml).

The ferric reducing ability of plasma was assayed according to Benzie and Strain (1996), where reductants ('antioxidants') in the sample reduce Fe (III)/tripyridyltriazine complex to a blue ferrous form, with an increase in absorbance at 593 nm. The final results were expressed in μmol trolox equivalents (TE)/g FW with reference to a standard curve of trolox (25–500 μmol).

Statistical analysis

The experimental design was randomized for cultivated plants, with three replicates. For quality attributes and health-related assays, 20 fruits of both varieties were analysed and all assays were performed in triplicates. Statistical analysis of variance was performed using SPSS 19.0 for Windows (SPSS Inc., Chicago, IL, USA). Data from 2019 and 2022 were analysed separately. A split-plot model with three replicates was used for variance analysis of all parameters, where variety was attributed to the main plots and fertilizer treatments to the subplots. Significant differences among groups were compared using least significant difference (LSD) test at $P < 0.05$. In addition, the technique for order preference by similarity to ideal solutions (TOPSIS) method, one of the multi-criteria decision-making methods, was used.

Results

Yield parameters

The fruit diameter, fruit length, mean fruit weight, the number of fruits per plant, total fruit weight and Brix yield of the treatments are presented in Table 3. Regarding FD, no significant differences were found in 2019; however, in 2022, treatment and the interaction between experimental factors had significant effects ($P < 0.01$). On the basis of average results, CF and SD treatments increased FD by approximately 11.4 and 10.3% compared with C treatment and by 4.4 and 3.4% compared with LD treatment, respectively.

Regarding FL, only variety had a significant effect in 2019 ($P < 0.01$). In the second year of the experiment, variety, treatment and the interaction between experimental factors had significant effects ($P < 0.01$, $P < 0.01$ and $P < 0.05$, respectively). CF and SD treatments of cv. 'Zeplin' had similar and highest FL values. The lowest FL was found with C treatment of cv. 'Arte'. On the basis of average results, SD and CF treatments increased FL by approximately 11.8 and 9.8% compared with C treatment and by 4.0 and 2.1% compared with LD treatment, respectively. MFW was significantly affected by variety, treatment and interaction ($P < 0.01$) in 2019; however, in 2022, treatment and the interaction between experimental factors were found to be significant ($P < 0.01$) (Table 3). The MFW changed between 104.3 g (SD treatment of cv. 'Zeplin') and 82.3 g (LD treatment of cv. 'Arte'). On the basis of average results, SD treatment increased MFW by 11.3, 4.8 and 12.3% compared with C, CF and LD treatments, respectively. In 2022, MFW changed between 94.0 g (CF treatment of cv. 'Zeplin') and 71.7 g (C treatment of cv. 'Zeplin'). According to average results, CF and SD treatments increased MFW by approximately 19.3 and 17.5% compared with C treatment and by 5.8 and 4.2% compared with LD treatment, respectively.

In 2019, the number of fruits per plant was significantly influenced by treatment ($P < 0.01$) (Table 3). CF, SD and LD treatments increased fruit number per plant by an average of 59.1, 62.4 and 54.0%, respectively, compared with control. In 2022, the differences between treatments and interactions were statistically significant ($P < 0.01$). The highest fruit number was obtained from the CF treatment of cv. 'Arte', whereas C treatment of cv. 'Arte' yielded the lowest. On the basis of average results, CF treatment resulted in the highest fruit number that significantly differed by unfertilized and fertilized thesis and this value was 56.9, 14.5 and 42.9% greater than those obtained from C, SD and LD treatments, respectively (Table 3).

For total fruit weight, statistical analysis revealed differences ($P < 0.01$) only between treatments in both years (Table 3). In 2019, the TFW value of SD treatment was the highest, with 2.51 kg per plant, followed by that of CF treatment (2.32 kg per plant). In the second year of the experiment, SD treatment resulted in the highest TFW (2.97 kg per plant) that significantly differed by unfertilized and fertilized thesis and this value was 25.3, 14.2 and 17.9% greater than those obtained from C, CF and LD treatments, respectively.

Regarding Brix yield, only treatment had a significant effect ($P < 0.01$) in both years (Table 3). The highest values were observed in CF and SD treatments in both years. CF and SD treatments increased Brix yield by approximately 33.0 and 40.5% in 2019 and 25.0 and 21.2% in 2022 compared with C treatment and by 10.0 and 16.3% in 2019 and 17.2 and 13.7% in 2022 compared with LD treatment, respectively.

As a result of ANOVA including the interactions between the experimental year (Y), treatment (T) and variety (V) of the parameters in Table 3, it was found that the mean of the years is different at 0.05 significance level.

Quality parameters

The colour characteristics of fruits are presented in Table 4. Fruit skin C^* was significantly affected by variety ($P < 0.01$) and treatment ($P < 0.05$) in 2019; however, in 2022, only treatment affected fruit skin C^* ($P < 0.05$). LD and CF treatments yielded the highest skin C^* values, followed by C treatment. In the second year of the experiment, CF, LD and C treatments showed similar and higher skin C^* values than SD treatment. For the h° value of fruit, variety and interaction showed significant differences in 2019 ($P < 0.01$ and $P < 0.05$, respectively). CF and SD treatments of cv. 'Zeplin' showed similar and lowest h° values. In the second year of the experiment, fruit h° values were significantly affected by variety and treatment ($P < 0.01$). Considering average results, CF treatment yielded the lowest h° value (39.5), whereas SD, C and LD treatments gave the highest h° values (Table 4). The a^*/b^* ratio of fruit skin was significantly affected by variety, treatment ($P < 0.01$ and $P < 0.05$, respectively) and the interaction between experimental factors ($P < 0.05$) in 2019 (Table 4). The highest skin a^*/b^* ratio was found with CF treatment of cv. 'Zeplin', followed by SD treatment of cv. 'Zeplin'. C and CF treatments of cv. 'Arte' gave the lowest a^*/b^* ratio. In 2022, the skin a^*/b^* ratio was significantly affected by variety and treatment ($P < 0.01$). The highest a^*/b^* ratio was attained from CF treatment, with a mean value of 1.22.

Pulp C^* was significantly influenced by variety and treatment in 2019 ($P < 0.01$) (Table 4). The highest and lowest values of pulp C^* were obtained from SD and CF treatments, respectively. In 2022, the effects of treatment and interaction were significant ($P < 0.01$). Pulp C^* values changed between 34.1 (C treatment of cv. 'Arte') and 40.3 (SD treatment of cv. 'Zeplin'). Pulp h° was significantly affected by treatment in 2019 ($P < 0.05$). The lowest pulp h° value was obtained from CF treatment, followed by SD treatment, whereas control yielded the highest pulp h° value. In 2022, pulp h° values were significantly affected by treatment and interaction ($P < 0.01$ and $P < 0.05$, respectively). CF and SD treatments of cv. 'Zeplin' and cv. 'Arte' had similar and lowest h° values. For the a^*/b^* ratio of pulp, non-significant differences were found in the first year of the experiment; however, the a^*/b^* ratio of pulp was significantly affected by treatment ($P < 0.01$) and interaction ($P < 0.05$) in 2022 (Table 4). CF and SD

Table 3. Effect of digestate fractions on fruit size and yield parameters of processing tomato cultivars

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
FD (mm)	C	48.7	48.0	48.3	44.0	45.2	44.6 c	49.0 a	47.8 b
	CF	49.0	49.3	49.2	47.8	51.6	49.7 a		
	SD	49.3	51.0	50.2	51.3	47.1	49.2 ab		
	LD	50.7	47.7	49.2	46.5	48.7	47.6 b		
	Mean	49.0	49.0		47.4	48.1			
LSD 0.05	Variety	NS			NS				
	Treatment	NS			1.838**				
	Interaction	NS			2.599**				
FL (mm)	C	62.7	65.3	64.0	57.9	64.3	61.1 c	64.9 a	65.6 b
	CF	62.0	68.0	65.0	63.2	71.0	67.1 ab		
	SD	59.7	70.0	64.8	67.7	68.8	68.3 a		
	LD	62.3	69.0	65.7	64.1	67.3	65.7 b		
	Mean	61.7 b	68.1 a		63.2 b	67.9 a			
LSD 0.05	Variety	1.871**			1.596**				
	Treatment	NS			2.257**				
	Interaction	NS			3.192*				
MFW (g)	C	83.7	85.0	84.3 c	74.7	71.7	73.2 c	87.8 a	82.3 b
	CF	84.0	95.0	89.5 b	80.7	94.0	87.3 a		
	SD	83.3	104.3	93.8 a	91.0	81.0	86.0 a		
	LD	82.3	84.7	83.5 c	81.7	83.3	82.5 b		
	Mean	83.3 b	92.3 a		82.0	82.5			
LSD 0.05	Variety	2.242**			NS				
	Treatment	4.484**			3.461**				
	Interaction	4.970**			4.894**				
Fruit number (no/plant)	C	24.7	22.7	23.7 b	36.0	39.7	37.8 d	34.1 a	47.6 b
	CF	37.3	38.0	37.7 a	61.7	57.0	59.3 a		
	SD	42.0	35.0	38.5 a	51.7	52.0	51.8 b		
	LD	35.7	37.3	36.5 a	38.0	45.0	41.5 c		
	Mean	33.3	34.9		46.8	48.4			
LSD 0.05	Variety	NS			NS				
	Treatment	4.124**			2.813**				
	Interaction	NS			3.979**				
TFW (kg/plant)	C	1.99	1.84	1.92 c	2.24	2.50	2.37 b	2.20 a	2.62 b
	CF	2.18	2.46	2.32 ab	2.56	2.63	2.60 b		
	SD	2.39	2.64	2.51 a	3.13	2.80	2.97 a		
	LD	1.96	2.13	2.05 bc	2.35	2.69	2.52 b		
	Mean	2.13	2.27		2.57	2.66			
LSD 0.05	Variety	NS			NS				
	Treatment	0.337**			0.291**				
	Interaction	NS			NS				
BY (tonne/ha)	C	9.45	9.02	9.23 c	10.25	10.98	10.62 b	11.41 a	12.02 b

(Continued)

Table 3. (Continued.)

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
	CF	12.27	12.29	12.28 a	13.02	13.53	13.27 a		
	SD	12.71	13.24	12.97 a	13.33	12.40	12.87 a		
	LD	10.63	11.66	11.15 b	11.27	11.37	11.32 b		
	Mean	11.26	11.55		11.97	12.07			
LSD 0.05	Variety	NS			NS				
	Treatment	0.989**			0.922**				
	Interaction	NS			NS				

C, control; CF, chemical fertilizer; SD, solid digestate; LD, liquid digestate; FD, fruit diameter; FL, fruit length; MFW, mean fruit weight; TFW, total fruit weight; BY, Brix yield.

* $P < 0.05$; ** $P < 0.01$; NS, not significant.

treatments of cv. 'Zeplin' and cv. 'Arte' showed similar and highest a*/b*.

At the 0.05 significance level, the mean of the years differed, according to the results of an ANOVA that included the interactions between the experimental year (Y), treatment (T) and variety (V) of the factors in Table 4.

The pericarp thickness, fruit firmness, dry matter, total soluble solid, titratable acidity and pH of the treatments are presented in Table 5. PT was significantly affected by variety ($P < 0.01$) and treatment ($P < 0.05$) in the first year of the experiment. However, in 2022, statistical analysis revealed differences between the treatment and the interaction between experimental factors ($P < 0.01$). On the basis of average results, in 2019, C treatment yielded the highest PT, followed by SD treatment. C and SD treatments increased PT by approximately 13.0 and 7.1% compared with CF treatment and by 9.9 and 4.5% compared with LD treatment, respectively. In 2022, PT values changed between 6.99 and 5.83 mm. According to the average results, SD treatment resulted in the highest value of PT that significantly differed by unfertilized and fertilized thesis and SD treatment increased 7.3, 12.8 and 6.2% compared with C, CF and LD treatments, respectively.

Regarding fruit firmness, no significant differences were found in 2019; however, in 2022, FF was significantly affected by variety, treatment and their interactions ($P < 0.01$, $P < 0.05$ and $P < 0.01$, respectively) (Table 5). FF changed between 28.3 and 36.8 N. On the basis of average results, CF treatment increased FF by 11.0, 7.7 and 7.7% compared with C, SD and LD treatments, respectively.

For dry matter content was significantly influenced by variety and treatment ($P < 0.01$) in 2019; however, in 2022, the differences were statistically significant between the treatment ($P < 0.01$) and interaction ($P < 0.05$) (Table 5). In the first year of the experiment, the DM content decreased among the treatments in the following order: LD (6.11%) > SD (5.90%) > CF (5.82%) > C (5.55%). In 2022, the highest DM content was obtained from the CF treatment of cv. 'Arte', whereas the lowest DM content was obtained from the control treatment of cv. 'Arte'. According to the average results, the CF treatment resulted in the highest value of DM that significantly differed by unfertilized and fertilized thesis and this treatment increased 9.2, 9.0 and 6.9% compared with C, SD and LD treatments, respectively.

Total soluble solid content (Brix) was significantly affected by treatment ($P < 0.01$) in 2019. CF, SD and LD treatments showed

similar and higher TSS contents than C treatment (Table 5). In 2022, the effects of treatment and interaction on TSS content were significant ($P < 0.01$). TSS content changed between 5.23° Brix (CF treatment of cv. 'Zeplin') and 4.30° Brix (LD treatment of cv. 'Zeplin'). On the basis of average results, TSS content of fertilizer treatments was in the order of CF (5.20° Brix) > LD (4.58° Brix) > C (4.57° Brix) > SD (4.45° Brix).

The effects of variety, treatment and their interactions on TA were statistically significant ($P < 0.01$) in both years (Table 5). TA changed between 0.41 g/100 ml and 0.56 g/100 ml in 2019. According to the average results, SD and LD treatments increased TA by 11.1 and 8.9% compared with CF treatment and by 19 and 16.7% compared with C treatment, respectively. In 2022, TA changed between 0.29 g/100 ml and 0.56 g/100 ml. Considering average results, the CF treatment resulted in the highest value of TA that significantly differed by unfertilized and fertilized thesis and this treatment increased 18.4, 18.4 and 25% compared with C, SD and LD treatments, respectively.

Fruit pH was influenced by variety and treatment ($P < 0.05$ and $P < 0.01$, respectively) in 2019; however, in 2022, treatment had no effect on fruit pH, whereas variety and interaction had significant effects ($P < 0.01$ and $P < 0.05$, respectively) (Table 5). In 2019, SD treatment yielded the lowest fruit pH, followed by LD treatment. In the second year of the experiment, fruit pH was the lowest with the C and SD treatments of cv. 'Arte'.

The application of ANOVA to the parameters presented in Table 5, which includes the interactions between the experimental year (Y), treatment (T) and variety (V), revealed that the mean values for the years are statistically different at the 0.05 level of significance.

Health-related compounds

Table 6 presents the contents of lycopene, vitamin C and total phenolics as well as AA of tomato fruits. Lycopene content was significantly influenced by variety, treatment and their interactions ($P < 0.01$) in both years. CF and LD treatments of cv. 'Zeplin' showed similar and higher lycopene contents than all other treatments in both years. The lowest lycopene content was found with the C treatment of cv. 'Arte' in both years.

Variety, treatment and interaction had a significant effect on vitamin C content ($P < 0.01$, $P < 0.01$ and $P < 0.05$, respectively) in 2019; however, in 2022, the effects of variety and treatment were significant ($P < 0.01$) (Table 6). LD and SD treatments of

Table 4. Effect of digestate fractions on the colour characteristics of processing tomato cultivars

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
C* (fruit skin)	C	43.6	46.2	44.9 ab	40.7	41.1	40.9 a	45.0 a	40.7 b
	CF	45.5	45.6	45.5 a	41.6	41.9	41.7 a		
	SD	42.6	44.5	43.6 b	39.3	39.0	39.2 b		
	LD	44.3	47.2	45.8 a	40.5	41.8	41.2 a		
	Mean	44.0 b	45.9 a		40.5	40.9			
LSD 0.05	Variety	1.062**			NS				
	Treatment	1.503*			1.441*				
	Interaction	NS			NS				
h° (fruit skin)	C	47.2	46.2	46.7	43.2	41.9	42.5 a	45.9 a	41.6 b
	CF	47.2	43.8	45.5	40.8	38.1	39.5 b		
	SD	46.1	45.0	45.6	44.0	41.3	42.7 a		
	LD	45.7	45.4	45.6	42.0	41.3	41.7 a		
	Mean	46.6 a	45.1 b		42.5 a	40.6 b			
LSD 0.05	Variety	0.717**			0.771**				
	Treatment	NS			1.090**				
	Interaction	1.434*			NS				
a*/b* (fruit skin)	C	0.93	0.96	0.94 b	1.05	1.12	1.09 b	0.97 a	1.13 b
	CF	0.93	1.04	0.99 a	1.16	1.28	1.22 a		
	SD	0.96	1.00	0.98 a	1.04	1.12	1.08 b		
	LD	0.95	0.98	0.97 ab	1.11	1.14	1.12 b		
	Mean	0.94 b	1.00 a		1.09 b	1.16 a			
LSD 0.05	Variety	0.021**			0.032**				
	Treatment	0.030*			0.046**				
	Interaction	0.043*			NS				
C* (pulp)	C	27.8	29.8	28.8 ab	34.1	39.4	36.7 a	28.5 a	36.6 b
	CF	24.6	28.9	26.7 c	39.4	35.5	37.5 a		
	SD	28.9	31.1	30.0 a	37.4	40.3	38.9 a		
	LD	27.6	29.4	28.5 b	34.2	32.7	33.5 b		
	Mean	27.3 b	29.8 a		36.3	36.9			
LSD 0.05	Variety	1.056**			NS				
	Treatment	1.494**			2.175**				
	Interaction	NS			3.076**				
h° (pulp)	C	35.0	33.8	34.4 a	29.9	31.5	30.7 a	33.4 a	29.7 b
	CF	33.8	30.9	32.3 c	28.6	28.0	28.3 b		
	SD	33.1	32.2	32.7 bc	28.9	28.5	28.7 b		
	LD	33.3	34.8	34.0 ab	30.4	31.8	31.1 a		
	Mean	33.8	32.9		29.4	29.9			
LSD 0.05	Variety	NS			NS				
	Treatment	1.540*			0.825**				
	Interaction	NS			1.167*				
a*/b* (pulp)	C	1.45	1.50	1.48	1.74	1.63	1.69 b	1.53 a	1.76 b

(Continued)

Table 4. (Continued.)

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
	CF	1.50	1.68	1.59	1.84	1.88	1.86 a		
	SD	1.54	1.59	1.56	1.81	1.84	1.83 a		
	LD	1.53	1.45	1.49	1.71	1.62	1.66 b		
	Mean	1.50	1.55		1.77	1.74			
LSD 0.05	Variety	NS			NS				
	Treatment	NS			0.058**				
	Interaction	NS			0.081*				

C, control; CF, chemical fertilizer; SD, solid digestate; LD, liquid digestate; C*, chroma; h°, hue angle; a*/b*, ratio.

* $P < 0.05$; ** $P < 0.01$; NS, not significant.

cv. 'Zeplin' showed similar and highest vitamin C values. The lowest content of vitamin C was found with the C treatment of cv. 'Arte' in both years.

Concerning another health-related compound, total phenolic content was significantly affected by variety, treatment and their interactions ($P < 0.01$) in both years (Table 6). In 2019, TPC changed between 54.8 mg GAE/100 g FW and 42.3 mg GAE/100 g FW. In 2022, TPC was the highest with the CF treatment of cv. 'Arte' and CF treatment of cv. 'Zeplin', whereas the C treatment of cv. 'Zeplin' yielded the lowest amount of TPC. On the basis of average results, TPC was the highest with C and CF treatments in 2019, whereas in the second year of the experiment, TPC was highest only with CF treatment.

AA was significantly affected ($P < 0.01$) by variety ($P < 0.05$) in 2019; however, in 2022, variety had no effect on AA, whereas treatment had a significant effect ($P < 0.05$) (Table 6). The AA of tomato fruits reduced in the following order: CF (2.91 $\mu\text{mol TE/g FW}$) > LD (2.79 $\mu\text{mol TE/g FW}$) > C (2.68 $\mu\text{mol TE/g FW}$) > SD (2.30 $\mu\text{mol TE/g FW}$).

The mean of the years differs at the 0.05 significant level, according to the results of an ANOVA that takes into account the interactions between the experimental year (Y), treatment (T) and variety (V) of the factors in Table 6.

Physiological disorders of tomato fruit

The ratio of SS and BER in tomato fruits is presented in Table 7. SS was significantly affected by variety, treatment and interaction ($P < 0.01$) in both years. In 2019, SD treatment decreased fruit SS by approximately 61.7, 37.9 and 52.6% compared with C, CF and LD treatments, respectively. In the second year of the experiment, CF and SD treatments decreased fruit SS by approximately 60.2 and 65.9% compared with C treatment and by 62.0 and 67.4% compared with LD treatment, respectively.

BER was significantly influenced by variety ($P < 0.05$), treatment and interaction ($P < 0.01$) in 2019; however, in 2022, BER was significantly influenced only by treatment ($P < 0.01$) (Table 7). In 2019, according to the average results, CF and SD treatments decreased BER by approximately 58.5 and 63.4% compared with C treatment and by 54.1 and 59.5% compared with LD treatment (2019), respectively. In the second year of the experiment, the lowest value of fruit BER was achieved by SD treatment. This value was 83.3, 71.4 and 83.3% lower than those obtained from C, CF and LD treatments, respectively.

As a result of ANOVA including the interactions between the experimental year (Y), treatment (T) and variety (V) of the parameters in Table 7, it was found that the mean of the years is different at 0.05 significance level.

Discussion

Effect of fertilizer treatments on yield parameters

The present study showed that tomato plants fertilized with SD had equal or even higher fruit size and yield production parameters than tomato plants fertilized with CF in both years (Table 3). LD did not decrease fruit size and tomato yield – their values were similar, with no significant differences between control, or slightly higher than those of the control in both years. Nkoa (2014) reported some conflicting results regarding crop yields with respect to anaerobic digestate fertilizer treatments. Yu *et al.* (2010) reported that livestock digestates showed lower yields in tomatoes than the synthetically fertilized control. Moreover, no difference was found in the size of tomatoes treated with control, mineral N fertilizer and LD (Barzee *et al.*, 2019). However, Panuccio *et al.* (2021), who assessed the effectiveness of two digestate fractions (liquid and solid) on tomato production, demonstrated that digestate treatments decreased the number and size of tomato fruit compared with control. Our results, which were consistent with the results of SD used in this study, were in agreement with those described by Barzee *et al.* (2019) and Li *et al.* (2023). In contrast to the present study, Zheng *et al.* (2019) reported that compared with CF, LD increased tomato yield. Comparing studies on tomato treated with various digestates is difficult because of the varying cultivars, environments and agronomic practices used in each study. The variation in yield can be explained by changes in the physical and chemical properties of the digestates, which are mostly associated with the origin of the substrates (feedstock), operating parameters used in the digester and type of solid–liquid separation (Albuquerque *et al.*, 2012; Akhiar *et al.*, 2017). According to Barzee *et al.* (2019), the mode of application, split fertilization and time of application play a large role in nutrient availability, with some fertigation systems capable of reducing nutrient loss. Finally, in the present study, SD led to higher yield than LD, which can be explained by the higher nutrient (total nitrogen, phosphorus and potassium) and organic matter content of SD leading to better plant nutrition (Table 2). The results of the

Table 5. Effect of digestate fractions on the technological traits of processing tomato cultivars

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
PT (mm)	C	8.08	8.41	8.25 a	6.51	6.32	6.41 b	7.72 a	6.47 b
	CF	6.95	7.65	7.30 b	5.83	6.37	6.10 c		
	SD	7.58	8.05	7.82 ab	6.99	6.77	6.88 a		
	LD	6.95	8.07	7.51 b	6.22	6.75	6.48 b		
	Mean	7.39 b	8.05 a		6.39	6.55			
LSD 0.05	Variety	0.432**			NS				
	Treatment	0.611*			0.269**				
	Interaction	NS			0.381**				
FF (N)	C	28.0	28.5	28.2	36.3	28.3	32.3 b	28.9 a	33.7 b
	CF	28.7	28.3	28.5	36.8	34.9	35.9 a		
	SD	28.0	30.9	29.5	33.9	32.6	33.3 b		
	LD	28.8	29.5	29.2	33.7	32.9	33.3 b		
	Mean	28.4	29.3		35.2 a	32.2 b			
LSD 0.05	Variety	NS			1.385**				
	Treatment	NS			1.959*				
	Interaction	NS			2.771**				
DM (%)	C	5.62	5.48	5.55 c	6.08	6.15	6.12 b	5.85 a	6.30 b
	CF	6.15	5.49	5.82 b	6.95	6.40	6.68 a		
	SD	6.14	5.67	5.90 b	6.13	6.13	6.13 b		
	LD	6.30	5.92	6.11 a	6.20	6.30	6.25 b		
	Mean	6.05 a	5.64 b		6.34	6.25			
LSD 0.05	Variety	0.136**			NS				
	Treatment	0.193**			0.251**				
	Interaction	NS			0.355*				
TSS (°Brix)	C	4.83	5.00	4.92 b	4.67	4.47	4.57 b	5.32 a	4.70 b
	CF	5.73	5.20	5.47 a	5.17	5.23	5.20 a		
	SD	5.43	5.43	5.43 a	4.37	4.53	4.45 b		
	LD	5.53	5.37	5.45 a	4.87	4.30	4.58 b		
	Mean	5.38	5.25		4.77	4.63			
LSD 0.05	Variety	NS			NS				
	Treatment	0.278**			0.200**				
	Interaction	NS			0.282**				
TA (g citric acid/100 ml)	C	0.43	0.41	0.42 c	0.46	0.29	0.38 b	0.47 a	0.39 b
	CF	0.49	0.41	0.45 b	0.56	0.34	0.45 a		
	SD	0.50	0.50	0.50 a	0.45	0.30	0.38 b		
	LD	0.56	0.42	0.49 a	0.41	0.31	0.36 b		
	Mean	0.50 a	0.43 b		0.47 a	0.31 b			
LSD 0.05	Variety	0.014**			0.021**				
	Treatment	0.019**			0.029**				
	Interaction	0.027**			0.042**				
pH	C	4.43	4.47	4.45 a	4.50	4.64	4.57	4.43 a	4.61 b

(Continued)

Table 5. (Continued.)

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
	CF	4.46	4.51	4.49 a	4.58	4.67	4.63		
	SD	4.39	4.37	4.38 b	4.56	4.68	4.62		
	LD	4.37	4.43	4.40 b	4.60	4.61	4.61		
	Mean	4.41 b	4.45 a		4.56 b	4.65 a			
LSD 0.05	Variety	0.025*			0.031**				
	Treatment	0.035**			NS				
	Interaction	NS			0.062*				

C, control; CF, chemical fertilizer; SD, solid digestate; LD, liquid digestate; FT, pericarp thickness (mm); FF, fruit firmness; DM, fruit dry matter; TSS, total soluble solid; TA, titratable acidity.
* $P < 0.05$; ** $P < 0.01$; NS, not significant.

present study also showed that the effects of fertilizer treatment on FD, FL, MFW and fruit number per plant varied with genotype (Table 3).

Effect of fertilizer treatments on fruit quality parameters

Colour is probably the first quality factor considered by tomato product buyers. Therefore, an attractive deep red colour is a major quality attribute for tomato products (Garcia and Barrett, 2006). In the present study, fertilizer treatments had an unstable effect on colour traits, which varied with year and genotype (Table 4). These results were consistent with those reported by Morra *et al.* (2021) and Tallou *et al.* (2022), where digestate treatment significantly affected the colour traits of tomato fruits and fertilizer treatment had an unstable effect. By contrast, Barzee *et al.* (2019) found no clear differences in colour traits among tomato plants treated with control, mineral N fertilizer and digestate. One of the most important aspects of tomato fruit quality is its colour, which forms through a complex carotenoid pigment system that is influenced by genetics as well as the environment (López Camelo and Gómez, 2004).

The current paper showed that SD treatment yielded higher PT values in both years and the effects of fertilizer treatment on PT varied with genotype. PT in tomato fruit indicates firmness, thereby revealing whether fruits are suitable for canning, storage and long-distance transportation (Avdikos *et al.*, 2021).

Fruit firmness is one of the most important quality traits of tomatoes processed by the canning industry (Bilalis *et al.*, 2018). In the current work, CF treatment yielded the highest FF value, and the FF of tomato fruits treated with different fertilizer treatments showed variation with genotypes.

Similar results were reported by Viskelis *et al.* (2015), who compared the FF of conventional and organic tomato fruits and demonstrated that conventional fruits presented significantly higher FF values than organic ones, but only in some cultivars. In contrast to our study, a previous study showed that LD treatment yielded higher FF than CF and C treatments (no fertilization) (Zheng *et al.*, 2019).

In the present study, LD treatment yielded the highest DM value during the first year, whereas CF treatment yielded the highest DM value in the second year. Our results showed that the effects of fertilizer treatment on DM content varied with genotype. Data regarding the effects of digestates on DM content are not consistent in the

literature. Morra *et al.* (2021) found that the DM content of tomato was unaffected by fertilizer treatment (SD combined with reduced rates of NP fertilizers and control). However, Przygocka-Cyna *et al.* (2021) reported that treatment with a digestate combined with bio-mass ash increased the DM content of tomato compared with CF and control (no fertilization).

For the tomato processing business, TSS content (Brix) is economically important because even a slight increase in its value can result in a huge improvement in product output and reduce the cost of dehydrating puree into sauce and paste (Young *et al.*, 1993). The results of current study showed that CF, SD and LD treatments yielded the highest TSS values during the first year, whereas CF treatment yielded the highest TSS value during the second year. The findings of the present study also demonstrated that the effects of fertilizer treatment on TSS content varied with genotype. Ronga *et al.* (2020) evaluated the effects of different digestate fertilizers (LD, combined LD–biochar, pelleted digestate and control) and reported that LD induced the highest TSS (Brix) in processing tomato under the organic farming system. Moreover, Yu *et al.* (2010) and Zheng *et al.* (2019) reported that digestate treatments increased TSS content compared with control and conventional and/or CF management. Panuccio *et al.* (2021) demonstrated that LD treatment increased the TSS content of tomato fruit compared with control and SD treatment.

Because TA enhances the flavour of tomato products, it is an important attribute of processing tomatoes (Bilalis *et al.*, 2018). In this study, the highest TA values were observed with solid and LD treatments during the first year and with CF treatment during the second year. The effects of fertilizer treatment on TA varied with genotype. These results were consistent with those reported by Zheng *et al.* (2019), Panuccio *et al.* (2021) and Tallou *et al.* (2022), where digestate treatment increased TA values compared with control and/or CF treatments.

In the present study, SD and LD treatments decreased tomato fruit pH compared with control and CF treatments, except during the second year. Moreover, the effects of fertilizer treatment on pH varied with genotype (Table 5). These results were similar to the findings of Panuccio *et al.* (2021) and Ronga *et al.* (2020), who indicated that LD treatment decreased the pH value of tomato fruits. In contrast to the current research, Barzee *et al.* (2019) reported that there was no difference in the pH values of tomato fruits treated with control, mineral N fertilizer and digestate.

Table 6. Effect of digestate fractions on the health-related compounds of processing tomato cultivars

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
Lycopene (mg/kg)	C	56.2	71.0	63.6 d	48.9	60.5	54.7 d	72.9 a	61.6 b
	CF	80.7	87.8	84.3 a	67.6	72.8	70.2 a		
	SD	68.5	63.7	66.1 c	58.2	54.6	56.4 c		
	LD	69.6	85.6	77.6 b	59.1	71.1	65.1 b		
	Mean	68.7 b	77.0 a		58.5 b	64.8 a			
LSD 0.05	Variety	1.198**			1.215**				
	Treatment	1.694**			1.718**				
	Interaction	2.396**			2.429**				
Vitamin C (mg/100 g)	C	11.6	12.4	12.0 c	10.1	10.9	10.5 c	15.0 a	12.7 b
	CF	13.9	14.6	14.3 b	12.0	12.4	12.2 b		
	SD	15.6	18.2	16.9 a	13.2	14.8	14.0 a		
	LD	15.1	18.4	16.8 a	12.9	15.2	14.1 a		
	Mean	14.1 b	15.9 a		12.1 b	13.3 a			
LSD 0.05	Variety	0.606**			0.681**				
	Treatment	0.857**			0.963**				
	Interaction	1.212*			NS				
TPC (mg GAE/100 g FW)	C	52.7	49.2	50.9 a	47.9	43.3	45.6 c	48.3 a	48.1 b
	CF	49.0	49.2	49.1 ab	55.9	53.1	54.5 a		
	SD	54.8	42.3	48.6 b	43.6	45.3	44.4 c		
	LD	46.5	42.8	44.7 c	50.2	45.5	47.9 b		
	Mean	50.8 a	45.9 b		49.4 a	46.8 b			
LSD 0.05	Variety	1.660**			1.501**				
	Treatment	2.348**			2.122**				
	Interaction	3.320**			3.001*				
AA (μmol TE/g FW)	C	4.21	3.72	3.96	2.66	2.69	2.68 ab	3.90 a	2.67 b
	CF	3.95	3.81	3.88	3.22	2.60	2.91 a		
	SD	3.94	3.90	3.92	2.37	2.23	2.30 b		
	LD	4.28	3.41	3.84	2.77	2.82	2.79 a		
	Mean	4.10 a	3.71 b		2.75	2.59			
LSD 0.05	Variety	0.355*			NS				
	Treatment	NS			0.388*				
	Interaction	NS			NS				

C, control; CF, chemical fertilizer; SD, solid digestate; LD, liquid digestate; TPC, total phenolic content; AA, antioxidant activity.

* $P < 0.05$; ** $P < 0.01$; NS, not significant.

Effect of fertilizer treatments on fruit health-related compounds

Lycopene is the most prevalent carotenoid in the tomato fruit because of its AA against chronic diseases (Viskeliš *et al.*, 2015). In this study, the highest lycopene content was achieved with CF treatment. Regarding vitamin C content, SD and LD treatments yielded higher vitamin C values in both years. The effects of fertilizer treatment on lycopene and vitamin C contents varied with genotype. These results were consistent with those of Panuccio

et al. (2021), who reported that LD increased lycopene content compared with the control, and both LD and SD increased vitamin C content compared with the control. Some studies have indicated that the lycopene content of tomatoes is most significantly influenced by a range of genetic and environmental factors, including the cultivar, growing season and cultivation conditions (Toor *et al.*, 2006; Rickman Pieper and Barrett, 2008).

In the present study, control and CF treatments yielded the highest TPC value during the first year, whereas CF treatment

Table 7. Effect of digestate fractions on the physiological disorders of processing tomato cultivars

Years		2019			2022			Y × T × V	
Parameters	Treatment	cv. 'Arte'	cv. 'Zeplin'	Mean	cv. 'Arte'	cv. 'Zeplin'	Mean	2019	2022
Fruit affected by SS (%)	C	0.6	8.7	4.7 a	8.0	9.7	8.8 a	3.3 a	6.1 b
	CF	1.4	4.4	2.9 b	2.3	4.7	3.5 b		
	SD	2.5	1.0	1.8 c	4.0	2.0	3.0 b		
	LD	2.2	5.4	3.8 a	7.0	11.3	9.2 a		
	Mean	1.7 b	4.9 a		5.3 b	6.9 a			
LSD 0.05	Variety	0.571**			1.004**				
	Treatment	1.807**			1.420**				
	Interaction	1.142**			2.009**				
Fruit affected by BER (%)	C	1.9	6.3	4.1 a	1.2	1.1	1.2 a	2.8 a	0.8 b
	CF	0.9	2.4	1.7 b	0.7	0.7	0.7 b		
	SD	2.5	0.4	1.5 b	0.2	0.2	0.2 c		
	LD	3.4	4.0	3.7 a	0.9	1.5	1.2 a		
	Mean	2.2 b	3.3 a		0.8	0.9			
LSD 0.05	Variety	0.837*			NS				
	Treatment	1.183**			0.271**				
	Interaction	1.674**			NS				

C, control; CF, chemical fertilizer; SD, solid digestate; LD, liquid digestate; SS, sunscald; BER, blossom-end rot.

* $P < 0.05$; ** $P < 0.01$; NS, not significant.

yielded the highest TPC value during the second year. The effects of fertilizer treatment on TPC varied with genotype. Regarding AA, CF and LD treatments yielded higher in comparison with other treatments for only the second year of the experiment. These results were consistent with those reported by Panuccio *et al.* (2021), who indicated that TPC and AA levels increased with LD in a concentration-dependent manner compared with control. TPC and AA are influenced by genotype, environmental conditions, cultural practices, postharvest handling and processing techniques and their interactions (Ceglie *et al.*, 2016; Panuccio *et al.*, 2019).

Effect of fertilizer treatments on fruit physiological disorders

In the present study, SD reduced the incidence of SS and BER and their incidence was similar or even lower than those obtained with CF treatment in both years. LD did not increase these fruit physiological disorders, and their incidence was similar, with no significant differences, to that observed in the control group or slightly lower than those observed in the control group in both years. This result was consistent with that of Ronga *et al.* (2020), who reported that LD increased the incidence of BER compared with control. The differences observed in this study could be attributed to genotypic variability as well as the chemical, biochemical and biological properties of the soil, which have both direct and indirect influences on nutrient availability, thereby affecting BER occurrence. Djangsou *et al.* (2019) reported that fruit calcium deficiency and translocation cannot be considered as the sole primary factors of BER incidence in tomato and that the occurrence of BER is associated with a multitude of environmental, genetic, agronomical and physiological factors, which play complex roles and interactions. Regarding SS, the different

results can be attributed to the genetic background of tomato varieties as well as the higher N content and higher leaf-shoot dry weight of plants treated with SD and CF treatments (data not published) than others. These results were consistent with those of Parisi *et al.* (2022), who highlighted the effects of N fertilization on the enhancement of vegetative biomass and revealed that N fertilizer provided good fruit covering and better protection against SS damage than the control.

Effect of fertilizer treatments on the yield, quality, health-related compounds and physiological disorders based on the multi-criteria decision-making method

The TOPSIS method was used based on the year averages of four fertilizer treatments and 15 parameters (Table 8). In the present study, fertilizer treatments showed significant differences in terms of yield, quality, health-related compounds and physiological disorder parameters (Fig. 2). Regarding the parameters considered in the TOPSIS method, the SD treatment of cv. 'Zeplin' showed the highest yield and quality, whereas the C treatment of cv. 'Arte' and cv. 'Zeplin' showed the lowest yield and quality. The closest to the ideal solution for the decision maker who had to choose between fertilizer treatments was determined as the CF treatment with 0.6991 for cv. 'Arte' and SD treatment with 0.7765 for cv. 'Zeplin'.


Conclusion

In conclusion, the application of SD yielded similar or higher fruit size and yield parameters than the application of CF in both years. Furthermore, the application of both fractions significantly maintained or improved fruit quality. However, their effects on fruit

Table 8. Results for the ideal solution for different varieties and treatments based on TOPSIS

cv. ‘Arte’ (taking the average of 2019 and 2022)										cv. ‘Zeplin’ (taking the average of 2019 and 2022)						
		Treatment	Score	Rank						Treatment	Score	Rank				
		C	0.1957	4						C	0.0809	4				
		CF	0.6991	1						CF	0.6719	2				
		SD	0.6929	2						SD	0.7765	1				
		LD	0.2497	3						LD	0.3719	3				
Parameters	TFW	MFW	FN	BY	TSS	TA	pH	PT	FF	a*/b* fruit skin	a*/b* pulp	Lycopene	Vitamin C	BER	SS	
Weights	0.20	0.05	0.05	0.10	0.10	0.05	−0.05	0.05	0.05	0.05	0.05	0.10	0.05	−0.05	−0.05	

TFW, total fruit weight; MFW, mean fruit weight; FN, number of fruits per plant; BY, Brix yield; TSS, total soluble solid; TA, titratable acidity; PT, pericarp thickness; FF, fruit firmness; BER, blossom-end rot; SS, sunscald.

**Figure 2.** Bar plot showing the relative proximity to the ideal solution for fertilizer treatments (C, control; CF, chemical fertilizer; SD, solid digestate; LD, liquid digestate).

quality traits varied with year and variety. Moreover, the 2-year average results of the TOPSIS method revealed that the SD treatment of cv. 'Zeplin' showed the highest performance in terms of yield, quality, health-related compounds and physiological disorders. These results prove that SD can be successfully applied as the main fertilizer source to specialty crops, such as processing tomatoes, using the industry standard.

Acknowledgements. We are grateful to Ege University Planning and Monitoring Coordination of Organizational Development and Directorate of Library and Documentation for their support in editing and proofreading this manuscript. We also thank Professor Ozlem Alpu (Eskisehir Osmangazi University, Faculty of Science, Department of Statistics, Eskisehir, Turkey) for the TOPSIS analysis method.

Author contributions. O. A. and F. S. conceived the presented idea. O. A. and B. B. planned and carried out the experiments. M. T. and A. R. O. performed the numerical calculations for the suggested experiment. O. A. and F. S. conducted data collection. O. A. and F. S. performed the statistical analyses. O. A. wrote the manuscript with help from F. S.

Funding statement. This study was funded by the PROJECT titled 'Research on the Use of Digestates as Fertilizer Source in Agricultural Production: processing tomato and silage maize in the frame of the program Ege University Service and Consultancy Agreement' under the grant

agreement number '20180204'. SÜTAŞ Dairy Products Inc. had no role in the design, analysis or writing of this article.

Competing interests. None.

Ethical standards. Not applicable.

References

- Akdeniz B, Demirbükür Kavak B and Ağdatlioğlu N (2012) Use of factorial experimental design for analyzing the effect of storage conditions on color quality of sun-dried tomatoes. *Scientific Research and Essays* 7, 477–489.
- Akhiar A, Battimelli A, Torrijos M and Carrere H (2017) Comprehensive characterization of the liquid fraction of digestates from full-scale anaerobic co-digestion. *Waste Management* 59, 118–128.
- Albuquerque JA, de la Fuente C, Ferrer-Costa A, Carrasco L, Cegarra J, Abad M and Bernal MP (2012) Assessment of the fertiliser potential of digestates from farm and agroindustrial residues. *Biomass and Bioenergy* 40, 181–189.
- Antoniou N, Monlau F, Sambusiti C, Ficara E, Barakat A and Zabaniotou A (2019) Contribution to circular economy options of mixed agricultural wastes management: coupling anaerobic digestion with gasification for enhanced energy and material recovery. *Journal of Cleaner Production* 209, 505–514.
- AOAC (1990) *Official Method of Analysis*. Arlington, USA: Association of Official Analytical Chemists.
- AOAC (1995) *Official Methods of Analyses*, 14th Edn. Washington, DC: Association of Official Analytical Chemists.
- Avdikos ID, Tagiakas R, Tsouvaltzis P, Mylonas I, Xynias IN and Mavromatis AG (2021) Comparative evaluation of tomato hybrids and inbred lines for fruit quality traits. *Agronomy* 11, 609.
- Barra Caracciolo A, Bustamante M, Nogues I, Di Lenola M, Luprano M and Grenni P (2015) Changes in microbial community structure and functioning of a semiarid soil due to the use of anaerobic digestate derived composts and rosemary plants. *Geoderma* 245–246, 89–97.
- Barzee TJ, Edalati A, El-Mashad H, Wang D, Scow K and Zhang R (2019) Digestate biofertilizers support similar or higher tomato yields and quality than mineral fertilizer in a subsurface drip fertigation system. *Frontiers in Sustainable Food Systems* 3, 58.
- Benzie IF and Strain JJ (1996) The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': the FRAP assay. *Analytical Biochemistry* 239, 70–76.
- Bergstrand KJ, Löfkvist K and Asp H (2020) Dynamics of nutrient availability in tomato production with organic fertilizers. *Biological Agriculture and Horticulture* 36, 200–212.
- Bhatt AH and Tao L (2020) Economic perspectives of biogas production via anaerobic digestion. *Bioengineering* 7, 74.
- Bilalis D, Krokida M, Roussis I, Papastilianou P, Travlos I, Cheimona N and Dede A (2018) Effects of organic and inorganic fertilization on yield

- and quality of processing tomato (*Lycopersicon esculentum* Mill.). *Folia Horticulturae* **30**, 321–332.
- Bolzonella D, Fatone F, Gottardo M and Frison N** (2018) Nutrients recovery from anaerobic digestate of agro-waste: techno-economic assessment of full scale applications. *Journal of Environmental Management* **216**, 111–119.
- Carricondo-Martínez I, Berti F and Salas-Sanjuán MDC** (2022) Different organic fertilisation systems modify tomato quality: an opportunity for circular fertilisation in intensive horticulture. *Agronomy* **12**, 174.
- Ceglie FG, Amodio ML and Colelli G** (2016) Effect of organic production systems on quality and postharvest performance of horticultural produce. *Horticulturae* **2**, 4.
- Chea L, Erika C, Naumann M, Smit I, Horneburg B and Pawelzik E** (2021) Morphological, leaf nutrient, and fruit quality characteristics of diverse tomato cultivars under organic low-input management. *Sustainability* **13**, 12326.
- Cristina G, Camelin E, Tommasi T, Fino D and Pugliese M** (2020) Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy soil and on a peat substrate: effects on tomato plants growth and on soil properties. *Journal of Environmental Management* **269**, 110767.
- Czekala W** (2022) Digestate as a source of nutrients: nitrogen and its fractions. *Water* **14**, 4067.
- Davis AR, Fish WW and Perkins-Veazie P** (2003) A rapid spectrophotometric method for analyzing lycopene content in tomato and tomato products. *Postharvest Biology and Technology* **28**, 425–430.
- Djangsou H, Francia E, Ronga D and Buti M** (2019) Blossom end-rot in tomato (*Solanum lycopersicum* L.): a multi-disciplinary overview of inducing factors and control strategies. *Scientia Horticulturae* **249**, 49–58.
- Doyeni MO, Baksinskaite A, Suproniene S and Tilvikiene V** (2021) Effect of animal waste based digestate fertilization on soil microbial activities, greenhouse gas emissions and spring wheat productivity in loam and sandy loam soil. *Agronomy* **11**, 1281.
- Duman I, Gümtiş M and Erkan S** (2003) The determination of agronomical and technological properties of processing tomato varieties to be introduced to the production in Marmara region (Turkey) and their reactions to infections by the agents of diseases. *Acta Horticulturae* **613**, 345–349.
- FAOSTAT-Food and Agriculture Organization** (2022) Available at <https://www.fao.org/faostat/en/#data/QC> (accessed 7 November 2023).
- Ferdous Z, Ullah H, Datta A, Anwar M and Ali A** (2018) Yield and profitability of tomato as influenced by integrated application of synthetic fertilizer and biogas slurry. *International Journal of Vegetable Science* **24**, 445–455.
- Garcia E and Barrett DM** (2006) Evaluation of processing tomatoes from two consecutive growing seasons: quality attributes, peelability and yield. *Journal of Food Processing and Preservation* **30**, 20–36.
- Hernández T, Chocano C, Moreno JL and García C** (2014) Towards a more sustainable fertilization: combined use of compost and inorganic fertilization for tomato cultivation. *Agriculture, Ecosystems and Environment* **196**, 178–184.
- Karaçalı I** (2012) *Storage and Marketing of Horticultural Products*. Ege University Agricultural Faculty Publication Number: 494. Izmir: Turkey.
- Koszal M, Parafiniuk S, Szparaga A, Bochniak A, Kocira S, Atanasov AZ and Kovalyshyn S** (2020) Impact of digestate application as a fertilizer on the yield and quality of winter rape seed. *Agronomy* **10**, 878.
- Lee ME, Steiman MW and St. Angelo SK** (2021) Biogas digestate as a renewable fertilizer: effects of digestate application on crop growth and nutrient composition. *Renewable Agriculture and Food Systems* **36**, 173–181.
- Li F, Yuan Y, Shimizu N, Magaña J, Gong P and Na R** (2023) Impact of organic fertilization by the digestate from by-product on growth, yield and fruit quality of tomato (*Solanum lycopersicon*) and soil properties under greenhouse and field conditions. *Chemical and Biological Technologies in Agriculture* **10**, 70.
- López Camelo AF and Gómez PA** (2004) Comparison of color indexes for tomato ripening. *Horticultura Brasileira* **22**, 534–537.
- Makádi M, Szegi T, Tomócsik A, Orosz V, Michéli E, Ferenczy A, Posta K and Biró B** (2016) Impact of digestate application on chemical and microbiological properties of two different textured soils. *Communications in Soil Science and Plant Analysis* **47**, 167–178.
- Mayerová M, Šimon T, Stehlík M, Madaras M, Koubová M and Smatanová M** (2023) Long-term application of biogas digestate improves soil physical properties. *Soil and Tillage Research* **231**, 105715.
- McGuire RG** (1992) Reporting of objective color measurements. *Hortscience* **27**, 1254–1255.
- Morra L, Cozzolino E, Salluzzo A, Modestia F, Bilotto M, Baiano S and del Piano L** (2021) Plant growth, yields and fruit quality of processing tomato (*Solanum lycopersicon* L.) as affected by the combination of biodegradable mulching and digestate. *Agronomy* **11**, 100.
- Moura LED and Golyński A** (2018) Critical points of industrial tomato from field to processing. *Horticultura Brasileira* **36**, 521–525.
- Nkoa R** (2014) Agricultural benefits and environmental risks of soil fertilization with anaerobic digestates: a review. *Agronomy for Sustainable Development* **34**, 473–492.
- Pan Z, Qi G, Adriamanohiarisoamanana FJ, Yamashiro T, Iwasaki M, Nishida T, Tangtaweewipat S and Umetsu K** (2018) Potential of anaerobic digestate of dairy manure in suppressing soil-borne plant disease. *Animal Science Journal* **89**, 1512–1518.
- Panthée DR, Labate JA, McGrath MT, Breksa AP and Robertson LD** (2013) Genotype and environmental interaction for fruit quality traits in vintage tomato varieties. *Euphytica* **193**, 169–182.
- Panuccio MR, Papalia T, Attinà E, Giuffrè A and Muscolo A** (2019) Use of digestate as an alternative to mineral fertilizer: effects on growth and crop quality. *Archives of Agronomy and Soil Science* **65**, 700–711.
- Panuccio MR, Mallamaci C, Attinà E and Muscolo A** (2021) Using digestate as fertilizer for a sustainable tomato cultivation. *Sustainability* **13**, 1574.
- Panwar NL, Kaushik SC and Kothari S** (2011) Role of renewable energy sources in environmental protection: a review. *Renewable and Sustainable Energy Reviews* **15**, 1513–1524.
- Parisi M, Burato A, Pentangelo A and Ronga D** (2022) Towards the optimal mineral N fertilization for improving peeled tomato quality grown in southern Italy. *Horticulturae* **8**, 697.
- Petropoulos SA, Fernandes Á, Xyrafis E, Polyzos N, Antoniadis V, Barros L and Ferreira CFR** (2020) The optimization of nitrogen fertilization regulates crop performance and quality of processing tomato (*Solanum lycopersicum* L. cv. Heinz 3402). *Agronomy* **10**, 715.
- Przygocka-Cyna K, Barłóg P, Spizewski T and Grzebisz W** (2021) Bio-fertilizers based on digestate and biomass ash as an alternative to commercial fertilizers – the case of tomato. *Agronomy* **11**, 1716.
- RG** (2010) Implementing regulation on the use of domestic and urban treatment sludge in soil: official Gazette of the Republic of Turkey; 27661. Available at <http://www.resmigazete.gov.tr/eskiler> (accessed 10 March 2019).
- Rickman Pieper J and Barrett DM** (2008) Effects of organic and conventional production systems on quality and nutritional parameters of processing tomatoes. *Journal of the Science of Food and Agriculture* **89**, 177–194.
- Rocha-Meneses L, Luna-delRisco M, González CA, Moncada SV, Moreno A, Sierra-Del Rio J and Castillo-Meza LE** (2023) An overview of the socio-economic, technological, and environmental opportunities and challenges for renewable energy generation from residual biomass: a case study of biogas production in Colombia. *Energies* **16**, 5901.
- Ronga D, Caradonia F, Parisi M, Bezzi G, Parisi B, Allesina G, Pedrazzi S and Francia E** (2020) Using digestate and biochar as fertilizers to improve processing tomato production sustainability. *Agronomy* **10**, 138.
- Shen W, Hu M, Qian D, Xue H, Gao N and Lin X** (2021) Microbial deterioration and restoration in greenhouse-based intensive vegetable production systems. *Plant and Soil* **463**, 1–18.
- Tallou A, Aziz F, Garcia AJ, Salcedo FP, El Minaoui FE and Amir S** (2022) Bio-fertilizers issued from anaerobic digestion for growing tomatoes under irrigation by treated wastewater: targeting circular economy concept. *International Journal of Environmental Science and Technology* **19**, 2379–2388.
- Thaipong K, Boonprakob U, Crosby K, Cisneros-Zevallos L and Hawkins Byrne D** (2006) Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis* **19**, 669–675.
- Toor RK, Savage GP and Heeb A** (2006) Influence of different types of fertilisers on the major antioxidant components of tomatoes. *Journal of Food Composition and Analysis* **19**, 20–27.
- Van Midden C, Harris J, Shaw L, Sizmur T and Pawlet M** (2023) The impact of anaerobic digestate on soil life: a review. *Applied Soil Ecology* **191**, 105066–105078.

- Viskelis P, Radzevicius A, Urbonaviciene D, Viskelis J, Karkleliene R and Bobinas C** (2015) Biochemical parameters in tomato fruits from different cultivars as functional foods for agricultural, industrial, and pharmaceutical uses. In ElShemy H (ed.), *Plants for the Future*. London: InTech Open, pp. 45–77.
- Young TE, Juvik JA and Sullivan JG** (1993) Accumulation of the components of total solids in ripening fruits of tomato. *Journal of the American Society for Horticultural Science* **118**, 286–292.
- Yu FB, Luo XP, Song CF, Zhang MX and Shan SD** (2010) Concentrated biogas slurry enhanced soil fertility and tomato quality. *Acta Agriculturae Scandinavica, Section B – Plant Soil Science* **60**, 262–268.
- Yu Y, Li P, Zhang J, Li J and Yu R** (2021) Comprehensive insights into the organic fractions on solid–liquid separation performance of anaerobic digestates from food waste. *Science of the Total Environment* **800**, 149608.
- Zheng W and Wang SY** (2001) Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry* **49**, 5165–5170.
- Zheng J, Li X, Zhang Y, Zhang P and Wang J** (2019) Effects of digestate application on tomato growth, yield, quality, and soil nitrogen content via integrated hole irrigation. *Journal of Biobased Materials and Bioenergy* **13**, 620–634.