

EPP0169

Deep learning approach to evaluate sex differences in response to neuromodulation in Major Depressive Disorder

S. Seenivasan^{1*}, M. Adamson² and A. Phillips¹

¹Palo Alto Veterans Affairs, Psychiatry, Palo Alto, United States of America and ²Palo Alto Veterans Affairs, Polytrauma, Palo Alto, United States of America

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.480

Introduction: Identifying the factors that mediate treatment response to rTMS in MDD patients can guide clinicians to administer more appropriate, reliable, and personalized interventions.

Objectives: The present study aimed to investigate sex differences in response to repetitive transcranial magnetic stimulation (rTMS) in Major Depressive Disorder (MDD) patients.

Methods: In this paper, we developed a novel pipeline based on convolutional LSTM-based deep learning (DL) to classify 25 female and 25 male subjects based on their rTMS treatment response.

Results: Five different classification models were generated, namely pre/post-rTMS female (model 1), pre/post-rTMS male (model 2), pre-rTMS female responder vs. pre-rTMS female non-responders (model 3), pre-rTMS male responder vs. pre-rTMS male non-responder (model 4), and pre-rTMS responder vs. non-responder of both sexes (model 5), achieving 93.3%, 98%, 95.2%, 99.2%, and 96.6% overall test accuracy, respectively.

Conclusions: These results indicate the potential of our approach to be used as a response predictor especially regarding sex-specific antidepressant effects of rTMS in MDD patients.

Disclosure: No significant relationships.

Keywords: EEG; rTMS; deep learning; sex differences

EPP0168

EEG Markers of Suicidal Ideation in Depressive Female Adolescents

A. Iznak^{1*}, E. Iznak², T. Medvedeva³, E. Damyanovich¹ and I. Oleichik⁴

¹Mental Health Research Centre, Laboratory Of Neurophysiology, Moscow, Russian Federation; ²Mental Health Research Centre, Laboratory Of Neurophysiology, Москва, Russian Federation; ³Mental Health Research Centre, Department Of Clinical Psychology, Moscow, Russian Federation and ⁴Mental Health Research Centre, Department Of Endogenous Mental Disorders And Affective Conditions, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.481

Introduction: Timely detection of suicidal thoughts is one of the ways to prevent suicide. Use the psychiatric interview only for this purpose in adolescents is often insufficient due to poor compliance. Thus, the search for objective neurophysiological markers of suicidal ideation in adolescents seems to be actual.

Objectives: The aim of the study was to reveal the relationships between pre-treatment EEG parameters and intensity of suicidal ideation in depressive female adolescents.

Methods: 72 female depressive patients (all right-handed, age 16–25, mean 18.2±2.6 years old) were enrolled in the study. Pre-treatment total HDRS-17 scores varied from 13 to 43 (mean 26.9±7.5). Multichannel eyes closed EEG was recorded, and absolute spectral power was calculated in 8 narrow frequency sub-bands. All patients answered the inventory on intensity of suicidal thoughts. Spearman's correlations between pre-treatment EEG parameters and suicidal ideation scores were analyzed.

Results: Scores of intensity of suicidal ideation correlated positively ($p < 0.05$) with values of EEG alpha2 (9–11 Hz) spectral power in F7, F8, F4, C3, C4, T4, P4 and O2 EEG leads, as well as with values of EEG delta (2–4 Hz) spectral power in F7, F3 and C3 EEG leads ($p < 0.05$).

Conclusions: The intensity of suicidal ideation in depressive female adolescents associates with wide propagation of EEG alpha2, especially over the right hemisphere, and with EEG signs of decreased functional state of anterior regions of the left hemisphere. The study supported by RBRF grant No.20-013-00129a.

Disclosure: No significant relationships.

Keywords: female adolescents; suicidal ideation scores; quantitative EEG

EPP0169

An 8-year longitudinal study of long-acting injectable (LAI) antipsychotics. Prescription trends and therapeutic drug monitoring to inform precision dosing

D. Piacentino^{1*}, F. Carpi², G. Giupponi² and A. Conca²

¹National Institutes of Health, National Institute On Drug Abuse/ national Institute On Alcohol Abuse And Alcoholism, Bethesda, United States of America and ²Central Hospital of Bozen, Department Of Psychiatry, Bozen, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.482

Introduction: Despite the widespread use of long-acting injectable (LAI) antipsychotics in schizophrenia and other disorders, there is a lack of longitudinal studies evaluating prescription trends and the usefulness of therapeutic drug monitoring (TDM) to inform dosing. Indeed, LAI prescription varies greatly among different areas of the world and over the years.


Objectives: Assess trends in LAI prescription in 2013–2020 at the Psychiatry Department of Bozen, Italy, and (2) analyze the correlation between dose of prescribed LAIs and blood levels measured via TDM.

Methods: Parametric statistics.

Results: LAIs were administered to 471 patients (\bar{x} age±SD=47.2±16.3 years; 56.3% men). The pie chart shows LAI treatment duration, i.e., from 1 to 7 consecutive years. The most used LAIs were haloperidol in 2013–2014 (26.5–31.8%) and paliperidone in 2015–2020 (22.5–25.7%). Dose adjustments were rather frequent, whereas the switch between LAI, due to moderate-to-side effects or unsatisfactory improvement of clinical status, was infrequent (41 cases/8 years). LAI interruption for the same reasons or non-compliance was even more infrequent (10 cases), but in 8 cases it happened for opposite reasons, i.e., achievement of patients' stabilization and good compliance. The Table shows doses and plasma levels of LAIs. Correlations between doses and plasma levels

were: haloperidol: $r=-0.037$, $p=0.620$; paliperidone: $r=0.290$; $p=0.000$; risperidone: $r=0.219$, $p=0.006$; fluphenazine $r=0.358$, $p=0.000$; aripiprazole: $r=-0.051$, $p=0.610$; olanzapine: $r=-0.090$, $p=0.634$.

Proportion of patients with different consecutive years of LAI antipsychotic use



LAI	Dose ($\bar{x} \pm SD$)	Dose (range)	Plasma levels ($\bar{x} \pm SD$)	Plasma levels (range)
Paliperidone	62.8 \pm 33.4	12.5-200.0	24.8 \pm 14.0	0.20-96.0
Haloperidol dose	93.6 \pm 29.2	37.5-300.0	24.8 \pm 14.1	0.20-98.0
Risperidone dose	36.8 \pm 11.1	12.5-50.0	21.4 \pm 1.1	3.7-84.0
Fluphenazine dose	24.5 \pm 1.0	12.5-100.0	0.75 \pm 0.08	0-6.0
Aripiprazole dose	341.4 \pm 12.8	100.0-400.0	195.0 \pm 198.0	14.7-889.0
Olanzapine dose	323.5 \pm 83.9	150.0-405.0	28.0 \pm 3.4	8.0-75.0

Conclusions: Haloperidol and paliperidone were the most used LAIs. Drug prescription trends and doses were stable over time. A significant positive correlation between dose and plasma level was found for paliperidone, fluphenazine, and aripiprazole.

Disclosure: No significant relationships.

Keywords: psychopharmacology; Precision Psychiatry; Long-acting injectable antipsychotics; Therapeutic drug monitoring

EPP0170

Depressive Symptoms and Suicidal Ideation in Spanish Medical Students

J.A. Blanco^{1*}, M. Rodríguez², I. Santos Carrasco^{1,3}, M. Queipo De Llano De La Viuda¹, L. Gallardo Borge¹, P. Marqués Cabezas¹, M. J. Mateos Sexmero¹ and J. Blanco Vilches¹

¹University Clinical Hospital of Valladolid, Psychiatry, Valladolid, Spain; ²University of Valladolid, Psychiatry, Valladolid, Spain and

³Clinical Hospital of Valladolid, Psychiatry, Valladolid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.483

Introduction: Medical students have higher rates of depression, anxiety and suicidal ideation over the general population. The onset of these disorders can be a risk factor with unfavorable impact in both medical care and their lives during the years of clinical specialization

Objectives: To assess the prevalence and factors involved in depression, anxiety and suicidal behavior in medical students of the University of Valladolid (Spain). The results are compared with a previous study conducted 5 years earlier

Methods: We used an online self-administered questionnaire that included demographic variables, academic information, sanitary data, Beck Depression Inventory (BDI), Generalized Anxiety Disorder 7 (GAD7), and MINI International Neuropsychiatric Interview for suicide. Chi-Square Test was used for categorical variables, Student's t-test for quantitative variables and Spearman's Coefficient to evaluate correlations between variables

Results: 362 students of all courses enrolled in Medicine at the University of Valladolid completed the survey. There were no differences between male and female students, both with high rates of moderate-severe depression (27% vs 30,4%), anxiety (42,9% vs 54,5%), and moderate-severe suicide risk (14,2% vs 10,7%). Previous study (n=584) also showed no differences between sexes but with lower rates of moderate-severe depression (14,3% vs 16,3%). 11% reported suicidal thoughts in the past month (11,6% previous study). There was a significant inverse correlation between medical career satisfaction and BDI scores

Conclusions: Five years later, rates of depression and suicide risk could have increased in medical students at the University of Valladolid. We urgently recommend the implementation of mental health prevention programs in this population

Disclosure: No significant relationships.

Keywords: medical students; Depression; Suicide

EPP0171

Depression and suicidality amongst infertile women: a hidden pandemic?

B. Ghosh Dastidar

Calcutta National Medical College, Psychiatry, Kolkata, India

doi: 10.1192/j.eurpsy.2022.484

Introduction: The psychosocial impact of infertility has been well researched and documented. However very little research has been conducted to assess the causative relationship between infertility and serious psychiatric illness such as suicide.

We conducted a cross-sectional study to investigate suicidal risk (suicidal ideation/ suicide attempts) amongst 100 infertile women undergoing infertility treatment at an IVF Centre based in Kolkata, India. Allied parameters such as depression, anxiety, quality of life were also studied.

Objectives: The goal of the presence study was to assess the suicidal risk (suicidal ideation / attempted suicide) amongst infertile women undergoing infertility/ ART treatment. And to examine the possible etiological factors behind increased levels of suicidal risk amongst infertile women.

Methods: Mini International Neuropsychiatric Interview was administered to 100 female patient's undergoing IVF treatment and 100 control patient's , visiting an IVF Centre based in Kolkata. Chi square test, independent t test and Z test used for statistical analysis.

Results: According to the results obtained by assessment of MINI scale , the prevalence of major depressive disorder (50 %) followed