SEROQUEL® (quetiapine furnarate) Tablets

RINGE SUMMARY of PRESCRIBING INFORMATION

HUNCATIONS AND USAGE: Bipolar Mania: SEROULEL is indicated for the short-term tristment of acute manic pissodes associated with lipidant disorder, as either monotherapy or adjunct therapy to lithium or divalprose. The efficacy of SEROULEL in acute flops or mania was established in two 3-week monotherapy thats and one 3-week adjunct therapy that if bipolar i patients initially recyclated for 10 or 7 days for acute mania. Effectiveness for more than 3-week is not ober systematically evaluated in clinical trials. Therefore, the physician who elects to use SEROULEL in catching the sense of the singular production of the sense of the se

periodically re-evaluate the long-term usefulness of the drug for the individual patient.

CONTRIBUCATIONS EPROUNCE is contraindicated in individuals with a known hypersensitivity to this medication or any of its ingredients.

WARHINGS: Neurolepic Malignant Syndrome (MMS) has been reported in association with administration of analogy-totic drugs, including SEROUELE. Rare cases of MMS have been reported with SEROUELE. Clinical manifestations of MS are hypersyrea, muscle right, a Hard or melal status, and evidence of antomicin liciability frequient passe or blood pressure, behyvaride, disphoresis, and cardiac dysriptiminal, Additional signs may include elevated crass-with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation includes both serious medical liness (e.g., presentions), systemic mitted, endiponists evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation include to other authorities of complications of the contrained of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation include to other authorities of course, here is the contrained of madequarety freated ecrapyramidal signs and symptomis (EPS). Other important considerations in the differential diagnosis include central articularities; crossly, heat stroke, faring liveral and many contrained and articularities of the contrained of the contrained the contrained the contrained the contrained of the contrained t

periodically during treatment. Any patient treated with ahybical antispsychotics should be mointered for symptoms of hyper-plycemia during treatment with ahybical antispsychotics should undergo leasing blood plucose testing. In some some player plycemia during treatment with ahybical antispsychotics was discontinued, however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect dury.

PRECALTINISE, General: Ordinatate hypoterations: ESPOLUEL. may induce orthostatic hypoteristic respited with discherses, tachycardia and, in some patients, syncope, especially during the initial dose-firtation period, probably reflecting its General: Ordinatate for a probably reflecting its General: Ordinatate during the probably reflecting its General: Ordinatate cradent in patients with known cardiovascular disease, linked or ordinatate with SEROOUE, compared with 0% (0407) on placebo and about 0.4% (2627) on active control drugs. SEROOUEL storogram with season and season of catheratics was observed in seasotialized with general control of the placeboard or conditions without would predispose patients to hypoterison and synopoe may be minimized by linking the intrinsion schedule is appropriate. Catarnetic The development of catheracts was observed in seasotialized with general catheracts. The servelopment of catheracts was observed in seasotialized with general catheracts. The servelopment of catheracts was observed in seasotialized with general catheracts. The servelopment of catheracts was observed in seasotialized with general catheracts. The servelopment of catheracts was observed in seasotialized with general catheracts. The servelopment of catheracts was observed in seasotialized with general catheracts. The servelopment of catheracts was observed in associations with general catheracts. The servelopment of catheracts was observed in associations with general catheracts. The servelopment is a seasotial catheract that the servelopment is a seasotial catheract that the servelo

screlling patients (per designates unitariates) internal to accordance to the product of the pro

SEROQUEL® (quetiapine furnarate) Tablets

Servicus. Servicus de l'acception de


Steven Johnson syndrome (S.S.). Beautiful Statistance Class: SERDOUEL is not a controlled substance. Physicial BMD ANUSE AND DEPENDENCE: Controlled Substance Class: SERDOUEL is not a controlled substance. Physicial BMD Anus BMD

osserved costey for signs of missiles of assists of selectivetts, e.g., development of interrace, increases in obese, and up-seeding behavior, and increases in the control selection of the control of t

SEROQUEL is a trademark of the AstraZeneca group of companies.

C AstraZeneca 2004 Rev 01/04

Manufactured for: AstraZeneca Pharmaceuticals LP Wilmington, Delaware 19850-5437



SEROQUEL is indicated for the treatment of acute manic episodes associated with bipolar I disorder and the treatment of schizophrenia. Patients should be periodically reassessed to determine the need for continued treatment.

Prescribing should be consistent with the need to minimize the risk of tardive dyskinesia, seizures, and orthostatic hypotension. A rare condition referred to as neuroleptic malignant syndrome (NMS) has been reported with this class of medications, including SEROQUEL.

There have been reports of diabetes mellitus and hyperglycemia-related adverse events associated with the use of atypical antipsychotics, including SEROQUEL.

The most commonly observed adverse events associated with the use of SEROQUEL in clinical trials were somnolence, dry mouth, dizziness, constipation, asthenia, abdominal pain, postural hypotension, pharyngitis, SGPT increase, dyspepsia, and weight gain.





AstraZeneca Pharmaceuticals LF

To prevent medication errors, write "SEROQUEL" clearly on your Rx pad. Spell "SEROQUEL" clearly over the phone. Redefine Success

www.SEROQUEL.com Please see Brief Summary of Prescribing Information on following page.