



RESEARCH ARTICLE

A note on the splittings of finitely presented Bestvina–Brady groups

Yu-Chan Chang

Department of Mathematics and Computer Science, Wesleyan University, Middletown, CT, USA
Email: yuchanchang74321@gmail.com

Received: 11 August 2024; **Revised:** 25 October 2024; **Accepted:** 30 October 2024;

First published online: 19 December 2024

Keywords: Bestvina–Brady groups; splittings

2020 Mathematics Subject Classification: Primary - 20F65; Secondary - 20E06, 20E08

Abstract

We show that when a finitely presented Bestvina–Brady group splits as an amalgamated product over a subgroup H , its defining graph contains an induced separating subgraph whose associated Bestvina–Brady group is contained in a conjugate of H .

1. Introduction

Let Γ be a finite simplicial graph with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. The associated *right-angled Artin group* (RAAG), denoted by A_Γ , is generated by $V(\Gamma)$, and two generators v and w commute whenever they are connected by an edge. A common question in group theory is: when does a group split as an amalgamated product or an HNN extension over a subgroup? For RAAGs, the splittings over infinite cyclic subgroups and abelian subgroups were characterized by Clay [3] and by Groves and Hull [6], respectively. Recently, Hull [7] generalized the splittings of RAAGs over abelian subgroups to non-abelian subgroups.

Let $\phi: A_\Gamma \rightarrow \mathbb{Z}$ be a homomorphism that sends all the generators to 1. The kernel of ϕ is called the *Bestvina–Brady group* and is denoted by BB_Γ . We only focus on finitely presented Bestvina–Brady groups, which is equivalent to saying that the flag complexes on the defining graphs are simply connected [1]. The author in [2] characterized the splittings of finitely presented Bestvina–Brady groups over abelian subgroups. In this note, we prove a result for the splittings of finitely presented Bestvina–Brady groups over non-abelian subgroups.

Theorem 1.1. *Let Γ be a finite simplicial graph with no cut vertices and whose associated flag complex is simply connected. Suppose that BB_Γ splits as an amalgamated product over a subgroup H . Then Γ contains an induced subgraph Λ that separates Γ and BB_Λ is contained in a conjugate of H .*

In other words, Theorem 1.1 says that if BB_Γ acts on a tree which is not a line, then there is an induced subgraph Λ of Γ such that Λ separates Γ and BB_Λ fixes an edge of T .

If Γ contains an induced subgraph Λ such that $\Gamma \setminus \Lambda$ has more than one connected component, then A_Γ splits over A_Λ and BB_Γ splits over BB_Λ . In the language of Bass–Serre Theory, all the vertex groups and edge groups of this splitting for A_Γ are finitely presented, but this is not always the case for the corresponding splitting for BB_Γ ; see Example 3.3. We remark that Γ contains a cut vertex if and only if BB_Γ splits as a free product.

The proof of Theorem 1.1 uses basic facts about groups acting on trees, and the idea of the proof is similar to those in [6], [2], and [7].




Figure 1. A directed triangle.

2. Preliminaries

2.1. Bestvina–Brady groups

Let Γ be a finite simplicial graph. The main result in [1] states that Γ is connected if and only if BB_Γ is finitely generated, and that the flag complex on Γ is simply connected if and only if BB_Γ is finitely presented. In the latter situation, Dicks and Leary found an explicit presentation:

Theorem 2.1. ([5, Corollary 3]) *Let Γ be a finite simplicial directed graph. If the flag complex on Γ is simply connected, then BB_Γ is generated by all the directed edges of Γ , and the relators are of the form ee^{-1} , where e^{-1} denotes the edge e with the opposite orientation, and $ef = g = fe$, where e, f , and g form a directed triangle; see Figure 1.*

2.2. Group acting on trees

Let G be a group acting on a tree T without inversions. We always assume that actions are minimal and nontrivial. An element $g \in G$ is called *elliptic* if it fixes a point in T ; otherwise, it is called *hyperbolic*. When $g \in G$ is elliptic, the set of points fixed by g is a subtree of T and is denoted by $\text{Fix}(g)$. When $g \in G$ is hyperbolic, it fixes a line in T on which it acts by translation. This line is called the *axis* of g and is denoted by $\text{Axis}(g)$.

Lemma 2.2. ([4, Lemma 1.1, Corollary 1.5], [6, Lemma 1.1]) *Let G be a group acting on a tree, and let g and h be commuting elements in G .*

- (1) *If h is hyperbolic, then $\text{Axis}(h) \subseteq \text{Fix}(g)$.*
- (2) *If both g and h are hyperbolic, then $\text{Axis}(g) = \text{Axis}(h)$.*
- (3) *If both g and h are elliptic, then $\text{Fix}(g) \cap \text{Fix}(h) \neq \emptyset$.*

3. Proof of Theorem 1.1

Throughout this section, we identify edges of a graph with elements in the associated Bestvina–Brady group.

Definition 3.1. *Let Γ be a finite simplicial graph. A **triangle path** P_Δ between two distinct edges e and f in Γ is a sequence of triangles $\Delta_1, \dots, \Delta_n$ such that*

- *the edges e and f are contained in Δ_1 and Δ_n , respectively;*
- *the triangles Δ_i and Δ_{i+1} share a unique edge for each $i = 1, \dots, n-1$;*
- *the triangles Δ_i and Δ_j do not share a common edge if $j \neq i-1$ or $j \neq i+1$.*

*The edge shared by Δ_i and Δ_{i+1} is called an **intermediate edge**.*




Figure 2. Two triangle paths and their intermediate edges (red edges). The graph on the right illustrates that all the triangles in a triangle path share a common vertex.

Two examples of triangle paths and their intermediate edges are given in Figure 2. Notice that triangle paths between two edges may not be unique, and all the triangles in a triangle path can share a common vertex.

Recall that a subgraph Λ of Γ separates two vertices (or two edges) if these two vertices (or edges) lie in different connected components of $\Gamma \setminus \Lambda$.

Lemma 3.2. *Let Γ be a finite simplicial graph without cut vertices and whose associated flag complex is simply connected. Let Λ be an induced subgraph of Γ . If every triangle path between e_1 and e_2 in $E(\Gamma)$ has an intermediate edge in $E(\Lambda)$, then Λ separates e_1 from e_2 .*

Proof. Since Γ has no cut vertices and the associated flag complex is simply connected, every edge of Γ is contained in a triangle, and there is a triangle path between any two edges in Γ . Let $e_1 = (u_1, v_1)$ and $e_2 = (u_2, v_2)$. Suppose that Λ does not separate e_1 from e_2 . Then, without loss of generality, there is an edge path p between the vertices u_1 and u_2 . This edge path p is contained in some triangle path P_Δ between e_1 and e_2 , and therefore, every vertex of p , possibly except for the two end vertices, is a vertex of an intermediate edge of P_Δ . Since Λ contains an intermediate edge of P_Δ , removing Λ will disconnect the path p . Thus, the path p cannot exist. Hence, the subgraph Λ separates e_1 from e_2 . \square

We now prove the main theorem.

Proof of Theorem 1.1. Let BB_Γ act on a tree T , and let $e_h \in E(\Gamma)$ be hyperbolic. Since BB_Γ splits as an amalgamated product, the tree T is not a path. Let Λ be the induced subgraph of Γ such that $E(\Lambda)$ consists of all the elliptic edges of Γ that fix $\text{Axis}(e_h)$ pointwise. Let e be an edge of $\text{Axis}(e_h)$. Then BB_Λ fixes e and is contained in a conjugate of H .

We now show that Λ separates Γ . We claim that there is an edge $f \in E(\Gamma) \setminus E(\Lambda)$ such that every triangle path between e_h and f has an intermediate edge in $E(\Lambda)$. Suppose to the contrary that for every edge $f' \in E(\Gamma) \setminus E(\Lambda)$, there is a triangle path $P_\Delta = \{\Delta_1, \dots, \Delta_n\}$ between e_h and f' such that none of its intermediate edges is in $E(\Lambda)$. Denote by $\{f_1, \dots, f_{n-1}\}$ the set of intermediate edges of P_Δ , where f_i is the edge shared by Δ_i and Δ_{i+1} . Since e_h and f_1 are contained in Δ_1 , they are commuting elements. Since e_h is hyperbolic, the element f_1 is also hyperbolic. Otherwise, it follows from Lemma 2.2 (1) that $f_1 \in E(\Lambda)$. Therefore, Lemma 2.2 (2) implies $\text{Axis}(e_h) = \text{Axis}(f_1)$. Similarly, since f_1 is hyperbolic and commuting with f_2 , the element f_2 is also hyperbolic and has the axis $\text{Axis}(f_2) = \text{Axis}(f_1)$. Continuing with the same argument, we have that the edges $e_h, f_1, \dots, f_{n-1}, f'$ are all hyperbolic with the same axis $\text{Axis}(e_h)$. Now, every edge in $E(\Gamma) \setminus E(\Lambda)$ is hyperbolic and has the axis $\text{Axis}(e_h)$, which is fixed by $E(\Lambda)$ pointwise. Thus, the set $E(\Gamma)$ fixes $\text{Axis}(e_h)$, contradicting the fact that T is not a path. This proves the claim. Therefore, the subgraph Λ separates e_h from f by Lemma 3.2.

Next, suppose that every edge of Γ is elliptic. Since the action of BB_Γ on T has no global fixed points, it follows from [8, p.64, Corollary 2] that there are two edges e_α and e_β in $E(\Gamma)$ such that the intersection $\text{Fix}(e_\alpha) \cap \text{Fix}(e_\beta)$ is empty. Let L be the geodesic in T between $\text{Fix}(e_\alpha)$ and $\text{Fix}(e_\beta)$, and let e be an edge of L . Let Λ be an induced subgraph of Γ such that every edge of Λ fixes e . Then, BB_Λ fixes e and is contained in a conjugate of H . We now show that Λ separates Γ . Let $P_\Delta = \{\Delta_1, \dots, \Delta_n\}$ be a triangle path between e_α and e_β . Denote by $\{f_1, \dots, f_{n-1}\}$ the set of intermediate edges of P_Δ , where f_i is the edge shared by Δ_i and Δ_{i+1} . For convenience, we write $f_0 = e_\alpha$ and $f_n = e_\beta$. Since f_i and f_{i+1} are contained in the




Figure 3. A splitting of a finitely presented Bestvina–Brady group over a finitely generated but not finitely presented subgroup.

triangle Δ_{i+1} , they are commuting elliptic elements. It follows from Lemma 2.2 (3) that the intersection $\text{Fix}(f_i) \cap \text{Fix}(f_{i+1})$ is nonempty for $i = 0, \dots, n-1$. Thus, there is a path L' in T from $\text{Fix}(f_0) = \text{Fix}(e_\alpha)$ to $\text{Fix}(f_n) = \text{Fix}(e_\beta)$ lying entirely in $\bigcup_{i=0}^n \text{Fix}(f_i)$. Since T is a tree, we have $L' = L$. Then the edge e belongs to $\text{Fix}(f_i)$ for some i . That is, there is an intermediate edge f_i of P_Δ that belongs to $E(\Lambda)$. Since the choice of the triangle path P_Δ between e_α and e_β is arbitrary, the subgraph Λ separates e_α from e_β by Lemma 3.2. \square

We end this section with one example.

Example 3.3. Let Γ be the graph shown in Figure 3. Let W be the set of vertices that are adjacent to either u or v but different from u and v . Let Λ , Γ_1 , and Γ_2 be the induced graphs on W , $V(\Gamma) \setminus \{u, v\}$, and $W \cup \{u, v\}$, respectively. Then Λ is an induced separating subgraph of Γ and $\text{BB}_\Gamma \cong \text{BB}_{\Gamma_1} *_{\text{BB}_\Lambda} \text{BB}_{\Gamma_2}$. However, the groups BB_Γ and BB_{Γ_2} are finitely presented, while BB_{Γ_1} and BB_Λ are finitely generated but not finitely presented.

Acknowledgements. The author thanks Lorenzo Ruffoni for his encouragement and helpful discussions. The author thanks the referee for the constructive comments and suggestions.

References

- [1] M. Bestvina and N. Brady, Morse theory and finiteness properties of groups, *Invent. Math.* **129**(3) (1997), 445–470.
- [2] Y. C. Chang, Abelian splittings and JSJ-decompositions of finitely presented Bestvina–Brady groups, *J Group Theory* **26**(4) (2023), 677–692.
- [3] M. Clay, When does a right-angled Artin group split over \mathbb{Z} ? *Internat. J. Algebra Comput.* **24**(6) (2014), 815–825.
- [4] M. Culler and K. Vogtmann, A group-theoretic criterion for property FA, *Proc. Amer. Math. Soc.* **124**(3) (1996), 677–683.
- [5] W. Dicks and I. J. Leary, Presentations for subgroups of Artin groups, *Proc. Amer. Math. Soc.* **127**(2) (1999), 343–348.
- [6] D. Groves and M. Hull, Abelian splittings of right-angled Artin groups, in *Hyperbolic geometry and geometric group theory*, vol. **73**, of Adv. Stud. Pure Math. (Mathematical Society of Japan, Tokyo, 2017), 159–165. doi: [10.2969/aspm/07310159](https://doi.org/10.2969/aspm/07310159)
- [7] M. Hull, Splittings of right-angled Artin groups, *Internat. J. Algebra Comput.* **31**(7) (2021), 1429–1432.
- [8] J.-P. Serre, *Trees*, (Springer-Verlag, Berlin-New York, 1980). Translated from the French by John Stillwell.