

Nutritional portrait of processed foods purchased in Québec (Canada), 2016-2022

Julie Perron¹, Alicia Corriveau¹, Sonia Pomerleau¹, Clara-Jane Rhéaume¹, Marie-Ève Labonté^{1,2},
Véronique Provencher^{1,2}

¹Centre NUTRISS - Nutrition, Santé et Société, Institut sur la nutrition et les aliments fonctionnels (INAF), Université Laval, Québec, Canada.

²École de nutrition, Université Laval, Québec, Canada

Corresponding author: Véronique Provencher, PhD, Institut sur la nutrition et les aliments fonctionnels (INAF), Université Laval, 2440, boulevard Hochelaga, Québec, Québec (Canada), G1V 0A6, Email: veronique.provencher@fsaa.ulaval.ca; Telephone : 418.656.2131 p.404607

Short title: Food supply and purchases in Québec (Canada)



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its

DOI 10.1017/S1368980025000588

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

No disclosure: Part of the results presented in this paper are included in a report that has been published on the Observatory's website (www.offrealimentaire.ca) as well as in a scientific paper published in a French journal ([*Cahiers de nutrition et de diététique*](#)).

Acknowledgements: The authors are grateful to all undergraduate students for their involvement in data collection, as well as to Pierre Gagnon, Laurélie Trudel and the Observatory's Scientific Committee and Board of Directors for their support.

Financial Support: The Food Quality Observatory received financial support from the *ministère de la Santé et des Services sociaux du Québec* (MSSS), the *ministère de l'Agriculture, des Pêcheries et de l'Alimentation du Québec* (MAPAQ), *Québec en forme*, *ministère de l'Économie et de l'Innovation*, Canada Foundation for Innovation and the Institute of Nutrition and Functional Foods (INAF) at *Université Laval*. This project is funded as part of the *Politique gouvernementale de prévention en santé* of the *Gouvernement du Québec*.

Conflict of Interest: All authors have no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Authorship: Conceptualization: VP, MEL; Carry out the study: SP, JP, AC; Statistical analysis: C-JR; Writing: JP and AC; Review and editing: all authors

Ethical Standards Disclosure: N/A

ABSTRACT

Objective: The Food Quality Observatory synthesizes the nutritional composition of 15 processed food categories commonly purchased in Québec (Canada). We assessed how the new Canadian front-of-pack (FoP) labelling regulation of a ‘high in’ symbol, to be implemented as of January 1st 2026, would be potentially reflected in these categories and how simulations of reformulation would impact the presence of the symbol.

Design: Nutritional information was obtained by collecting food products available in supermarkets and grocery stores in the province of Québec (2016-2022). Sales data were obtained from *NielsenIQ* company. 15 food categories have been selected, and 3 levels of reformulation were simulated.

Setting: The nutritional values of 5132 products were merged with sales data. 3941 products were successfully cross-referenced.

Results: Sixty percent of all products sold ($n=2336/3941$) would carry the ‘high in’ symbol reflecting a high content in sodium, saturated fat and/or total sugar (39%, 16%, and 17%, respectively). For certain food categories, a slight reduction (5 to 15%) in sodium, saturated fat or total sugar content would allow removing the ‘high in’ symbol in a large number of products. For example, a 5% reduction of the sodium content in sliced breads would allow 22 percentage point (pp) fewer products to display the symbol.

Conclusions: This study presents a portrait of processed foods purchased in Québec (Canada) and the distribution of the FOP ‘high in’ symbol. Such portrait generates important data to monitor the food supply’s nutritional quality, which can ultimately contribute to improve the nutritional quality of processed foods.

INTRODUCTION

A food supply that offers products with high levels of sodium, saturated fat and/or sugar is considered a major cause of non-communicable diseases⁽¹⁾. Several strategies have been implemented to facilitate consumers' choices towards more nutritious food and to improve the food supply's nutritional quality. Among these strategies, front-of-pack (FoP) labelling has been identified as effective to promote healthy food environments and to help consumers make healthier food choices^(2, 3). In 2016, Chile adopted a FoP warning symbol for food products exceeding specified amounts of energy, saturated fat, sugar or sodium⁽⁴⁾. Since then, Peru, Mexico, Israel, Brazil, Venezuela, Uruguay, Argentina and Colombia have enacted policies requiring FoP warning labels similar to Chile's^(5, 6). A comparable approach has been recently adopted in Canada with a FoP nutrition symbol approved in 2022, which will be mandatory by 2026⁽⁷⁾. This symbol will have to be displayed on packaging of products exceeding at least one of the thresholds established for total sugar, sodium and saturated fat, based on the food category to which they belong. For example, for products with a reference amount of 30 g or less, the threshold is 10% of the recommended daily value (DV), specific to each nutrient, compared to 15% of DV for products with a reference amount of more than 30 g.

FoP nutrient warning is increasing across the globe since it has been shown to be easy to understand, not to mention that it is an effective strategy to help consumers identify products high in nutrients to limit and therefore to discourage them from purchasing these products^(5, 8, 9). An underlying aim of such a FoP label is to motivate food industries to reformulate their products to avoid the symbol on their packaging. In 2022, the WHO published a policy brief which supports reformulation of foods and beverages⁽¹⁰⁾. Data from this report support that consumers accept reformulated products, and that reformulation leads to better nutritional intakes and better health. Reformulation of sodium, saturated fat and sugar contents can indeed contribute to ensuring access to more nutritious food and shifting towards healthier consumption patterns. The advantage with food reformulation is that consumers do not need to modulate their buying behavior or make conscious efforts to choose healthier options.

Sodium is the most reformulated nutrient since 62 countries have reformulation strategies to reduce sodium in packaged foods⁽¹¹⁾. As an example, the United Kingdom (UK) launched a

successful voluntary salt reduction program that has set five levels of progressively more stringent sodium targets since 2004^(12, 13). Seven years after, this program has led to reductions of up to 45% in sodium levels in some products and has led to a 15% drop in population salt intakes. However, such a voluntary program in Canada has not produced the expected results⁽¹⁴⁾. Sugar reformulation strategies are less common. In the UK, a 3.5% reduction in total sugar of products sold was observed four years after the implementation of the voluntary sugar reduction program in 2016⁽¹⁵⁾. Reformulation to reduce saturated fat by the food industry has been much less studied and has little success, while WHO suggests that saturated fat be replaced by unsaturated fat instead of reducing saturated fat alone⁽¹⁰⁾.

To our knowledge, very few studies have measured the impact of FoP nutrient symbol strategies on food reformulation⁽¹⁶⁾ and none of these studies have been conducted in Canada. It is thus of relevance to monitor the nutritional composition of the food supply prior to implementing such a strategy, in order to ultimately measure the impact of the FoP nutrition symbol on consumer choice and industry reformulation of foods. While the nutritional composition of packaged processed foods and beverages in Canada has been previously studied based on products available on shelves^(17, 18), no study has yet measured the nutritional composition of processed foods that are actually purchased in the country. Therefore, the aims of the present study were first to characterize the nutritional composition of 15 processed food categories commonly purchased in the province of Québec (Canada). We then verified the extent to which the Canadian FoP nutrition symbol would be displayed in those food categories and verify the number of nutrients concerned. Finally, we simulated the effects of various level of reformulation (5%, 10% and 15% reduction) for sodium, saturated fat and total sugar on the distribution of the FoP nutrition symbol.

METHODS

Data collection

Hosted by the Institute of nutrition and functional foods at *Université Laval*, the Food Quality Observatory (Observatory) was launched in 2016. The Observatory aims at monitoring the evolution of the food supply, hence contributing to the collective effort of improving the quality and accessibility of food. Consultations with the Observatory knowledge users and the scientific

committee were conducted to prioritize the processed food categories to monitor ($n=15$)⁽¹⁹⁾. To be retained, the food categories had to have an impact on health (i.e., high in sugar, sodium and/or saturated fat), show a variability in terms of nutritional quality, have a high household penetration rate and show a potential for improvement. The following food categories were thus selected for the analyses: ready-to-eat (RTE) breakfast cereals, sliced breads, luncheon meats, ready-to-serve soups, pizzas, granola bars, frozen meals, pasta sauces, yogurts and dairy desserts, sausages, cookies, crackers, salty snacks, processed cheeses, and flavoured milks and plant-based beverages. Note that luncheon meats, yogurt and dairy desserts, sausages and processed cheeses also included related plant-based alternatives.

To reach the objectives described above, a database containing the nutritional value and labelling information of each product was created by the Observatory ($n=10$ out of 15 categories), *Protégez-Vous* ($n=4$ out of 15 categories) – a Québec-based non-profit organization specialized in consumer information and products testing – and Health Canada ($n=1$ out of 15 categories). Data collections were done in supermarkets, grocery stores and specialty grocery stores from the Greater Québec City area, the Greater Montreal area (Québec, Canada) or across Canada (for pizzas only) between September 2016 and February 2022. All information on the product packaging (e.g., brand, Nutrition Facts table, list of ingredients, nutrition and health claims, serving size) was coded in the database using double coders. Nutritional value variables listed were as follows: energy (kcal), total fat (g), saturated fat (g), total sugar (g), fibre (g), protein (g) and sodium (mg).

This nutritional value database was then merged with a sales database (provided by *NielsenIQ* company⁽²⁰⁾) by using unique product codes (UPC). For each product, the database included the following data: sales in Canadian dollars (CAD\$), sales in kilograms (kg) and sales per unit. Sales information comes from the optical reading of the products purchased in the main food chains of Québec markets (or Canada markets for pizzas).

Statistical analyses

Means and standard deviations of nutritional values and price per serving of products were first generated. The averages were weighted by sales volume in kg to better represent what the population buys – and indirectly consumes – by giving a higher weight to the most popular

products of a given food category and a lower weight to the products which were less purchased. Since the analyses weighted for sales were conducted based on the combined database, the number of products analysed was lower than the number of food items found on shelves.

In order to determine if each product would carry the Canadian FoP nutrition symbol, the 15% of the daily value (DV) threshold was used for most products⁽⁷⁾. The 30% of the DV threshold was used for frozen meals and pizzas since they are considered as mixed dishes. The 10% of the DV threshold was used for products for which the reference amount was less than or equal to 30g. The highest amount between the serving size indicated on the package and the reference amount of the product⁽²¹⁾ was used for the calculation. Exemptions of the FoP nutrition symbol have also been considered (e.g., yogurts with high calcium content).

Simulations were performed by modelling different scenarios where food products underwent reformulation to reduce specific components (sodium, saturated fat, or sugar), by varying percentages of 5%, 10%, or 15%. These reductions were then presented for their impact on the distribution of FoP labels. Statistical analyses were conducted using SAS software version 9.4.

RESULTS

A total of 5132 different products were identified from the 15 selected food categories. Among them, 3941 products with sales information were successfully cross-referenced with the products identified in the food supply. According to *NielsenIQ* database, the products offered from which sales data were available represented an average market coverage of 79%.

Table 1 shows the variety of products offered, the number of products with purchases data and the market coverage with *NielsenIQ* database according to each food category, with cookies and salty snacks representing the largest food categories. (Add Table 1)

Table 2 shows the nutritional value of the products purchased (weighted by sales volume) according to each food category. (Add Table 2) Purchased pizzas, sausages and frozen meals had the highest content in saturated fat per serving. Purchased flavoured milks and plant-based beverages, RTE breakfast cereals and cookies had the highest sugar content per serving. Purchased pizzas, ready-to-serve soups and frozen meals had the highest sodium content per serving.

Figure 1 shows the current proportion of products which would carry a FoP nutrition symbol for 0, 1, 2 or 3 nutrients if no changes were made in their food composition. (Add Figure 1) In total,

60% of products purchased (n=2336/3941) would have the FoP nutrition symbol for at least one nutrient. More specifically, 39% would have the symbol for sodium, 17% for total sugar and 16% for saturated fat (see Table 3). Almost half of products purchased (46%) would carry the symbol for only one nutrient, 13% for two nutrients and 0.5% for the three nutrients. Yogurt and dairy desserts, RTE breakfast cereals, crackers and granola bars were the food categories that would most often carry no symbol. On the other hand, sausages, pizzas and cookies were the food categories which would most often carry the symbol for at least two nutrients.

Table 3 simulates the effects on the distribution of the FoP nutrition symbol of potential reductions (5%, 10% or 15%) in the sodium, saturated fat and total sugar content in each food category. (Add Table 3) Among salient results, a theoretical improvement of only 5% in sodium content of sliced breads would lead to a 22-percentage point (pp) reduction, going from 70% to 48%, of products with the FoP nutrition symbol for sodium. Moreover, a theoretical improvement of only 5% in the saturated fat content of pizzas would lead to a 12-pp reduction, going from 71% to 59%, of products with the FoP nutrition symbol for this nutrient. Finally, a theoretical reduction of 10% in total sugar in cookies would lead to a 28-pp reduction, going from 77% to 49%, of products with the symbol for sugar. On the other hand, a 15% reduction in sodium content for ready-to-serve soups and luncheon meats or a 15% reduction in saturated fat for sausages would not be enough to modify the FoP distribution in those food categories.

DISCUSSION

This novel study provides an overview of the nutritional composition, distribution of potential FoP nutrition symbols and reformulation simulations of processed foods purchased in the province of Québec (Canada). Few studies have examined different food categories simultaneously, and those that have done so, have not included weighted sales data in their analyses due to limited access.

Nutritional composition

First, results showed that the vast majority of food categories analyzed in the present study have a high sodium content. Per 100g, processed cheeses, luncheon meats and sausages were the categories with the highest sodium content, an observation similar to what has been reported in the literature. Indeed, according to another Canadian study carried out on 7234 food items from

13 food categories, cheese products offered had indeed the highest average sodium content (1471mg/100g), followed by packaged deli meats (1092mg/100g)⁽²²⁾. In Argentina, an analysis conducted on 1320 products from 14 food categories has revealed that sausages (1050mg/100g) and ready meals offered (941mg/100g) were among the food categories with the highest sodium content⁽²³⁾.

Moreover, our study revealed that, per 100g, purchased cookies, granola bars, and RTE breakfast cereals were the categories with the highest sugar content. Similarly, a Canadian study showed that cookies (32g/100g), granola bars (30g/100g) and RTE breakfast cereals (21g/100g) offered had the highest sugar content⁽²⁴⁾. In the New Zealand food supply, the granola bar category ranked third in terms of sugar content, following jams and confectionery⁽²⁵⁾.

In the present study, processed cheeses, cookies, and sausages were among the food categories with the highest content of saturated fat, again per 100g. Although fewer studies have been carried out on saturated fat, a New Zealand study reported that the highest saturated fat content was in cheeses (18g/100g), biscuits (8g/100g) and processed meats (6g/100g)⁽²⁵⁾.

In summary, food categories high in sodium, sugar or saturated fat are similar elsewhere in Canada and around the world. Moreover, for all nutrients of concern, a large variability was observed between food categories and within each category (e.g., sodium content varies from 112mg to 373mg/100g in ready-to-serve soups containing vegetables, starch and proteins⁽²⁶⁾, sugar content varies between 3g to 15g/100g for stirred yogurts⁽²⁷⁾, and saturated fat content varies between 2g to 8g/100g for deli meat pizzas⁽²⁸⁾). Such findings suggest that a reduction is possible and realistic for a large proportion of products.

FoP nutrition symbol

In total, unless any changes are made in the nutritional composition, more than half of products purchased and analyzed in the present study would carry the Canadian FoP nutrition symbol for at least one nutrient, with sodium being the most common nutrient displayed. Accordingly, a research team in the province of Ontario (Canada) reported that a total of 66% of the food offered on shelves would carry the FoP nutrition symbol (32% for sodium, 28% for sugar and 28% for saturated fat)⁽¹⁷⁾. However, the comparison with our study is limited, as it did not incorporate weighted sales data, and it covered a wider range of food categories. In 2019 in

Chile, at the beginning of the FoP warning implementation (final stage), 83% of near 5500 food products had the symbol for at least one nutrient⁽²⁹⁾.

In the current study, one product out of eight would have the FoP nutrition symbol for two nutrients while a minority of products (0.5%) would have it for all nutrients. In comparison, in another Canadian study, the majority of consumed foods that would display a FoP nutrition symbol would be for only one nutrient accounting for 18% of energy intakes⁽³⁰⁾. Foods that would display a FoP symbol for two and for three nutrients accounted for 6% and 0.1% of food energy intakes respectively. The present study demonstrated that luncheon meats, ready-to-serve soups and pizzas were the categories which would carry the FoP nutrition symbol for almost all products purchased. In Chile, sweet baked products, soups and savoury snacks were the food categories with the most prevalent warning symbol before the implantation of the regulation (100%, 98% and 94%, respectively)⁽³¹⁾.

Simulations of reformulation

The present study demonstrates that reductions in the order of 5, 10 or 15% in sodium, saturated fat or total sugar can have a major impact on products with a potential Health Canada FoP nutrition symbol, with a large proportion of them being able to avoid such a symbol. Considering that reformulated products without nutritional warning were perceived in a Chilean study as healthier and had higher purchase intention scores than their regular counterparts with warning⁽³²⁾, this can be a motivating call to action for food industries. Since those food categories are largely consumed by Canadians, these little changes can reduce purchases of nutrients of public health concern and ultimately reduce their intake in the population. Modelling studies have shown that modest reductions, if applied on all products of most purchased food categories, have the potential to improve population health⁽³³⁻³⁵⁾. As observed above, sodium is the nutrient for which reformulations should be attempted in priority. In a systematic review, sodium reformulation was carried out more often in breads, sauces and processed meats⁽³⁵⁾. In Chile, most frequent reductions of FoP prevalence in “high in sodium” concerned savory spreads, cheeses, ready-to-eat meals, soups, and sausages⁽³¹⁾.

However, this does not dismiss the fact that reformulation is challenging. Numerous considerations are raised on technical feasibility, shelf life, palatability and food safety, which

can be discouraging for the food industry^(36, 37). Moreover, some adverse effects of FoP warning symbol and reformulation may occur. Indeed, some countries noted an increase in the use of non-nutritive sweeteners as a replacement for sugar to avoid the warning symbol^(38, 39). Since non-nutritive sweeteners have been associated with adverse metabolic effects, particularly in youth (e.g., altered sweetness perceptions, gut microbiota dysbiosis and disrupts glucose homeostasis)⁽⁴⁰⁻⁴²⁾, some countries now display a warning on products for the presence of non-nutritive sweeteners⁽⁴³⁾. Similarly, the WHO discourages the use of non-nutritive sweeteners in replacement of sugar⁽⁴⁴⁾. As another collateral effect, some stakeholders anticipate the introduction of additives in recipes to avoid the FoP (e.g., potassium salts or monosodium glutamate to compensate for the diminution of sodium). The long-term effects of these additives on health are not yet known.

This study has several strengths. While the access to sales data is expensive, the combination of nutritional composition with sales data to monitor the actual foods purchased is an important strength of the present study. Another strength is that the sales data cover a large proportion of the food supply for all food categories studied with an average of approximately 80% overall. Moreover, this study is among the first to propose simulations of reformulation specific to each food category. However, the current study also has some limitations. Although the 15 food categories identified are widely bought by consumers, they do not cover all processed food categories. Also, since data collection and analysis of a single food category lasts approximately nine months, it was not possible to monitor all 15 food categories during the same year. Furthermore, the nutritional database is an overview at a given time, which may not represent the whole portrait of the food supply during the entire year, and does not evolve in real time. Different products may not have been identified, such as products that entered the market after data collection or those purchased at another moment during the year but that were discontinued before the data collection. Additionally, not all products were successfully matched to sales data. In fact, the sales database available through the *NielsenIQ* company does not include products from certain private labels of specific grocery stores. Moreover, even if food sales data can give an overview of food intakes^(45, 46), it is not possible to ensure that products purchased are truly eaten by the consumers who bought them.

In conclusion, this study presents an initial portrait of 15 food categories highly consumed in Québec (Canada), which represents the starting point to monitor the nutritional composition of food purchased in the province. In accordance with current food-related policies, monitoring the nutritional composition of food purchased after the implementation of the FoP Canadian regulation will be essential over the next few years⁽⁴⁷⁾. In that context, the Observatory will continue to monitor the composition of processed food purchases every five years. Based on these findings, it becomes possible to identify areas of improvement regarding the nutritional composition of processed foods, which is of great relevance for policymakers and public health nutrition advocates in the aim to encourage healthier food choices.

The full report is freely available on www.observatoire.inaf.ulaval.ca.

REFERENCES

1. Lane MM, Davis JA, Beattie S *et al.* (2021) Ultraprocessed food and chronic noncommunicable diseases: A systematic review and meta-analysis of 43 observational studies. *Obesity Reviews* 22, e13146.
2. Feteira-Santos R, Fernandes J, Virgolino A *et al.* (2020) Effectiveness of interpretive front-of-pack nutritional labelling schemes on the promotion of healthier food choices: a systematic review. *JBIC Evidence Implementation* 18, 24-37.
3. World Health Organization (2017). Tackling Ncds "best buys" and other recommended interventions for the prevention and control of noncommunicable diseases. Available at: <https://apps.who.int/iris/handle/10665/259232> (accessed December 2023).
4. DOF (2015) Reglamento sanitario de los alimentos. Diario Oficial de la Republica N°4193. http://web.minsal.cl/sites/default/files/decreto_etiquetado_alimentos_2015.pdf (accessed October 2020).
5. Crosbie E, Gomes FS, Olvera J *et al.* (2023) A policy study on front-of-pack nutrition labeling in the Americas: emerging developments and outcomes. *The Lancet Regional Health - Americas* 18, 100400.
6. Kanter R, Vanderlee L Vandevijvere S (2018) Front-of-package nutrition labelling policy: global progress and future directions. *Public health nutrition* 21, 1399-1408.

7. Government of Canada (2022) Summary of amendments published in the Canada Gazette, Part II: Nutrition symbols, other labelling provisions, vitamin D and hydrogenated fats or oils. Available at: <https://www.canada.ca/en/health-canada/services/food-labelling-changes/front-package/summary-amendments-gazette-nutrition-symbols-labelling.html> (accessed December 2023).
8. Taillie LS, Hall MG, Popkin BM *et al.* (2020) Experimental studies of front-of-package nutrient warning labels on sugar-sweetened beverages and ultra-processed foods: a scoping review. *Nutrients* 12, 569.
9. Croker H, Packer J, Russell SJ *et al.* (2020) Front of pack nutritional labelling schemes: a systematic review and meta-analysis of recent evidence relating to objectively measured consumption and purchasing. *Journal of human nutrition and dietetics* 33, 518-537.
10. World Health Organization (2022) Reformulation of food and beverage products for healthier diets: policy brief. Available online: <https://www.who.int/publications/i/item/9789240039919> (accessed December 2023).
11. Rosewarne E, Santos JA, Trieu K *et al.* (2022) A global review of national strategies to reduce sodium concentrations in packaged foods. *Advances in Nutrition* 13, 1820-1833.
12. Niblett P, Coyle N, Little E *et al.* (2020) Salt targets 2017: Second progress report. A report on the food industry's progress towards meeting the 2017 salt targets. Available online: <https://www.gov.uk/government/publications/salt-targets-2017-second-progress-report> (accessed December 2023). *Public Health England*.
13. He F, Brinsden H MacGregor G (2014) Salt reduction in the United Kingdom: a successful experiment in public health. *Journal of human hypertension* 28, 345-352.
14. Health Canada, Sodium Reduction in Processed Foods in Canada: An Evaluation of Progress toward Voluntary Targets from 2012 to 2016. 2018. Available at: <https://www.canada.ca/content/dam/hc-sc/documents/services/food-nutrition/legislation-guidelines/guidance-documents/guidance-food-industry-reducing-sodium-processed-foods-progress-report-2017/publ-eng.pdf> (accessed October 2020).
15. UK Parliament (2024) House of Lords Library. Eating less sugar: Reformulating food and drink products and government policy. Available at: <https://lordslibrary.parliament.uk/eating-less-sugar-reformulating-food-and-drink-products-and-government->

- [policy/#:~:text=Reduction%20and%20reformulation%20voluntary%20programme,in%20processed%20foods%20and%20drinks](#) (accessed January 2024).
16. Kanter R, Reyes M, Vandevijvere S *et al.* (2019) Anticipatory effects of the implementation of the Chilean Law of Food Labeling and Advertising on food and beverage product reformulation. *Obesity Reviews* 20, 129-140.
 17. Vergeer L, Vanderlee L, Ahmed M *et al.* (2020) A comparison of the nutritional quality of products offered by the top packaged food and beverage companies in Canada. *BMC public health* 20, 650.
 18. Mulligan C, Lee JJ, Vergeer L *et al.* (2022) Evaluating the potential implications of canadian front-of-pack labelling regulations in generic and branded food composition databases. *BMC public health* 22, 1866.
 19. Turcotte M, Trudel L J L (2016-2017) Consultation du comité des utilisateurs de connaissances de l'Observatoire de la qualité de l'offre alimentaire. *Observatoire de la qualité de l'offre alimentaire*.
 20. Nielsen IQ MarketTrack (2016-2021) Quebec All Channels, 52 weeks
 21. Government of Canada (2022) Table of Reference Amounts for Food. Available at: <https://www.canada.ca/en/health-canada/services/technical-documents-labelling-requirements/table-reference-amounts-food.html> (accessed December 2023).
 22. Arcand J, Au JT, Schermel A *et al.* (2014) A comprehensive analysis of sodium levels in the Canadian packaged food supply. *American journal of preventive medicine* 46, 633-642.
 23. Allemandi L, Tiscornia MV, Ponce M *et al.* (2015) Sodium content in processed foods in Argentina: Compliance with the national law. *Cardiovascular diagnosis and therapy* 5, 197.
 24. Bernstein JT, Schermel A, Mills CM *et al.* (2016) Total and Free Sugar Content of Canadian Prepackaged Foods and Beverages. *Nutrients* 8.
 25. Mackay Sally NMC, Swinburn B *et al.*, (2019) State of the food supply: New Zealand 2019. The University of Auckland, Auckland, New Zealand.
 26. Perron J, Pomerleau S, Gagnon P *et al.* (2020). Portrait des soupes prêtes-à-servir disponibles au Québec 2017. Observatoire de la qualité de l'offre alimentaire. 59 p.

27. Perron J, Pomerleau S, Gilbert-Moreau J *et al.* (2020). Portrait des yogourts et desserts laitiers disponibles au Québec - 2018-2019. Observatoire de la qualité de l'offre alimentaire. 55p.
28. Perron J, Pomerleau S, Gagnon P *et al.* (2021). Portrait des pizzas disponibles au Canada 2017. Observatoire de la qualité de l'offre alimentaire. 50 p.
29. Kanter R, Reyes M, Swinburn B *et al.* (2018) The Food Supply Prior to the Implementation of the Chilean Law of Food Labeling and Advertising. *Nutrients* 11.
30. Lee JJ, Ahmed M, Ng A *et al.* (2023) Nutrient intakes and top food categories contributing to intakes of energy and nutrients-of-concern consumed by Canadian adults that would require a 'high-in' front-of-pack symbol according to Canadian labelling regulations. *Plos one* 18, e0285095.
31. Reyes M, Smith Taillie L, Popkin B *et al.* (2020) Changes in the amount of nutrient of packaged foods and beverages after the initial implementation of the Chilean Law of Food Labelling and Advertising: A nonexperimental prospective study. *PLoS Med* 17, e1003220.
32. Schnettler B, Ares G, Sepúlveda N *et al.* (2019) How do consumers perceive reformulated foods after the implementation of nutritional warnings? Case study with frankfurters in Chile. *Food quality and preference* 74, 179-188.
33. Alessandrini R, He FJ, Ma Y *et al.* (2021) Potential impact of gradual reduction of fat content in manufactured and out-of-home food on obesity in the United Kingdom: a modeling study. *The American Journal of Clinical Nutrition* 113, 1312-1321.
34. Trieu K, Coyle DH, Afshin A *et al.* (2021) The estimated health impact of sodium reduction through food reformulation in Australia: A modeling study. *PLoS medicine* 18, e1003806.
35. Federici C, Detzel P, Petracca F *et al.* (2019) The impact of food reformulation on nutrient intakes and health, a systematic review of modelling studies. *BMC nutrition* 5, 1-21.
36. Fanzo J, McLaren R, Bellows A *et al.* (2023) Challenges and opportunities for increasing the effectiveness of food reformulation and fortification to improve dietary and nutrition outcomes. *Food Policy* 119, 102515.

37. Onyeaka H, Nwaiwu O, Obileke K *et al.* (2023) Global nutritional challenges of reformulated food: A review. *Food science & nutrition* 11, 2483-2499.
38. Lowery CM, Mora-Plazas M, Gómez LF *et al.* (2020) Reformulation of packaged foods and beverages in the Colombian food supply. *Nutrients* 12, 3260.
39. Zancheta Ricardo C, Corvalán C, Smith Taillie L *et al.* (2021) Changes in the use of non-nutritive sweeteners in the Chilean food and beverage supply after the implementation of the food labeling and advertising law. *Frontiers in nutrition* 8, 773450.
40. Shum B & Georgia S (2021) The effects of non-nutritive sweetener consumption in the pediatric populations: what we know, what we don't, and what we need to learn. *Frontiers in Endocrinology* 12, 625415.
41. Toews I, Lohner S, de Gaudry DK *et al.* (2019) Association between intake of non-sugar sweeteners and health outcomes: systematic review and meta-analyses of randomised and non-randomised controlled trials and observational studies. *bmj* 364.
42. Swithers SE (2015) Artificial sweeteners are not the answer to childhood obesity. *Appetite* 93, 85-90.
43. White M & Barquera S (2020) Mexico adopts food warning labels, why now? *Health Systems & Reform* 6, e1752063.
44. World Health Organization (2023). WHO advises not to use non-sugar sweeteners for weight control in newly released guideline. Available at: <https://www.who.int/news/item/15-05-2023-who-advises-not-to-use-non-sugar-sweeteners-for-weight-control-in-newly-released-guideline> (accessed December 2023).
45. Ransley JK, Donnelly JK, Khara TN *et al.* (2001) The use of supermarket till receipts to determine the fat and energy intake in a UK population. *Public health nutrition* 4, 1279-1286.
46. Martin SL, Howell T, Duan Y *et al.* (2006) The feasibility and utility of grocery receipt analyses for dietary assessment. *Nutrition journal* 5, 10.
47. Vanderlee L, McLaughlin A, Vaillancourt C *et al.* Actions to cultivate healthier food environments in Canada: Federal government policies and progress, 2023. Available at: www.informascanada.com/food-epi-canada-2023 (accessed December 2023).

Table 1. Availability of products among the selected food categories (n=15)

Food categories (year of collection)	Availability of products	Products offered matched with sales data	Supply coverage of the <i>NielsenIQ</i> database*
	Offer (n)	Purchases (n)	Market coverage (%)
RTE breakfast cereals (2016)	331	308	90
Sliced breads (2016)	294	262	75
Luncheon meats (2017)	361	317	62
Ready-to-serve soups (2017)	223	180	92
Pizzas (2017)	155	155	80
Frozen meals (2018)	386	275	70
Granola bars (2018)	310	240	75
Yogurts and dairy desserts (2018)	380	325	86
Pasta sauces (2019)	322	210	88
Cookies (2019)	694	494	87
Sausages (2019)	289	214	63
Crackers (2020)	439	223	94
Salty snacks (2020)	627	503	91
Processed cheeses (2020)	118	87	72
Flavoured milks and plant- based beverages (2022)	203	148	73
Total	5132	3941	79

n=number of unique products per food category

*Source: Nielsen IQ MarketTrack (2016-2022), Quebec All Channels, 52 weeks




Figure 1. Distribution of requirement to display FoP “high-in” nutrition symbol among purchased food categories

Table 2. Mean nutritional value of purchased food categories per portion

	Portion	Calories (kcal)	Lipids (g)	Saturated fat (g)	Carbohydrates (g)	Fibre (g)	Total sugar (g)	Protein (g)	Sodium (mg)
RTE breakfast cereals (n=308)*	55g	210±20	2.6±2.6	0.8±1.6	44±4	4.4±3.8	12.8±5.5	4.7±1.7	216±125
Sliced breads (n=262)	2 slices (73g)	189±34	2.3±0.9	0.5±0.2	35±7	2.7±1.7	3.0±2.4	7.3±1.6	331±83
Luncheon meats (n=317)	55g	99±45	6.2±5.1	2.2±1.8	2±1	0.0±0.2	0.6±0.7	8.7±2.4	542±148
Ready-to-serve soups (n=180)	250ml	130±51	3.5±4.2	1.1±1.2	19±6	3.9±3.0	3.5±2.7	5.3±2.6	715±132
Pizzas (n=155)	200g	474±51	19.5±5.7	7.5±2.5	55±6	3.3±1.2	6.7±3.0	19.8±3.4	1066±193
Frozen meals (n=275)	1 meal (278g)	331±100	10.2±6.6	3.7±2.8	44±13	3.1±1.4	6.8±6.0	15.8±5.4	698±237
Granola bars (n=240)	1 bar (34g)	143±30	5.1±2.4	1.6±1.0	23±5	2.0±1.3	9.9±3.7	2.5±1.6	94±42
Yogurt and dairy desserts (n=325)	1 unit (110g)	76±29	1.5±1.4	0.9±0.8	12±5	0.1±0.4	9.2±4.1	3.7±1.5	53±19
Pasta sauces (n=210)	125ml	73±37	3.0±3.7	1.1±2.2	9±2	1.9±0.8	5.3±1.4	2.5±1.2	476±110
Cookies (n=494)	30g	141±28	5.7±1.7	2.3±1.4	21±5	0.8±0.7	10.4±3.2	1.6±0.7	87±43

Sausages (n=214)	55g (cooked/pre-cooked) or 75g (raw)	156±33	12.6±3.5	4.2±1.5	4±1	0.1±0.3	0.5±0.9	8.2±1.8	483±78
Crackers (n=223)	23g	90±14	3.4±1.4	0.8±0.6	13±2	0.7±0.7	0.9±1.0	1.7±0.5	150±50
Salty snacks (n=503)	50g	262±19	14.7±3.7	1.9±1.3	29±4	1.9±1.0	1.3±1.8	3.2±1.0	332±131
Processed cheeses (n=87)	30g	76±12	5.2±1.7	3.1±1.1	4±2	0.0±0.1	2.2±1.1	3.8±1.3	365±130
Flavoured milks and plant-based beverages (n=148)	250ml	123±49	3.3±1.1	1.1±0.9	18±10	0.7±0.8	14.7±10.4	5.3±3.2	149±47

Mean ± SD

*n=number of products with sales data

Table 3. Distribution of FoP nutrition symbol according to nutrients of interest and simulations of reformulation of 5, 10 and 15%

	Proportion of products with the FoP nutrition symbol for at least one nutrient (weighted for sales)	Proportion of products with FoP nutrition symbol for sodium (weighted for sales)				Proportion of products with FoP nutrition symbol for saturated fat (weighted for sales)				Proportion of products with FoP nutrition symbol for sugar (weighted for sales)			
Food category	Actual	Actual (%)	-5% (%) ^a	-10% (%) ^b	-15% (%) ^c	Actual	-5% (%) ^a	-10% (%) ^b	-15% (%) ^c	Actual	-5% (%) ^a	-10% (%) ^b	-15% (%) ^c
RTE breakfast cereals (n=308)	32.6	2.9	0.7	0.2	0.0	7.4	6.8	6.8	6.8	23.4	14.6	8.1	3.8
Sliced breads (n=262)	70.8	69.6	47.5	41.5	19.1	0.0	0.0	0.0	0.0	2.4	1.8	1.8	0.0
Luncheon meats (n=317)	99.8	99.8	98.3	97.3	96.7	43.6	39.8	39.3	32.6	0.0	0.0	0.0	0.0
Ready-to-serve soups (n=180)	99.8	99.8	99.6	99.5	99.0	9.2	5.6	5.6	5.3	1.7	1.7	0.7	0.5
Pizzas (n=155)	99.9	93.6	91.6	85.2	79.4	71.0	59.4	52.3	38.1	0.0	0.0	0.0	0.0
Frozen meals (n=275)	64.7	63.7	51.0	45.0	33.8	20.1	17.4	16.3	12.3	2.1	1.8	1.5	1.5
Granola bars (n=240)	40.5	0.0	0.0	0.0	0.0	22.9	16.9	13.4	7.1	30.2	24.3	20.6	16.3
Yogurt and dairy	25.4	0.0	0.0	0.0	0.0	0.5	0.5	0.5	0.4	25.1	24.5	20.8	18.9

desserts (n=325)													
Pasta sauces (n=210)	77.1	77.1	74.5	64.1	58.4	11.0	10.3	10.3	10.3	0.0	0.0	0.0	0.0
Cookies (n=494)	84.4	0.0	0.0	0.0	0.0	57.7	44.9	44.6	44.5	76.7	65.9	49.1	44.9
Sausages (n=214)	97.5	97.0	96.8	96.6	92.7	85.7	85.6	85.6	83.3	0.0	0.0	0.0	0.0
Crackers (n=223)	33.2	24.5	19.7	7.9	5.5	11.9	10.7	10.7	7.5	0.1	0.0	0.0	0.0
Salty snacks (n=503)	44.5	38.4	35.2	29.6	20.7	9.9	6.6	6.6	6.6	0.3	0.3	0.3	0.1
Processed cheeses (n=87)	39.5	12.8	11.9	11.9	11.9	39.5	39.5	39.5	39.5	0.0	0.0	0.0	0.0
Flavoured milks and plant-based beverages (n=148)	52.9	3.3	0.0	0.0	0.0	11.3	9.8	9.8	0.9	50.2	48.3	48.3	48.3
Total	59.2	39.1	32.4	29.5	22.6	16.3	14.3	13.9	11.8	17.4	15.5	13.4	12.0

^aProportion of products with FoP nutrition symbol with -5% reformulation

^bProportion of products with FoP nutrition symbol with -10% reformulation

^cProportion of products with FoP nutrition symbol with -15% reformulation