

Plastic pollution and human–primate interactions: A growing conservation concern

Evelyn D. Anca¹  and Janette Wallis²

¹Faculty of Humanities and Social Sciences, Oxford Brookes University, Oxford, UK and ²Kasokwa-Kityedo Forest Project, Masindi, Uganda

Perspective

Cite this article: Anca ED and Wallis J (2024). Plastic pollution and human–primate interactions: A growing conservation concern. *Cambridge Prisms: Plastics*, 2, e10, 1–8 <https://doi.org/10.1017/plc.2024.10>

Received: 30 December 2023
Revised: 02 April 2024
Accepted: 21 April 2024

Keywords:

plastic pollution; primates; conservation; human–wildlife interaction; transport and wildlife vectors

Corresponding author:

Evelyn D. Anca;
Email: evelynanca@gmail.com

Abstract

As an anthropogenic creation, plastic pollution is a form of human–wildlife interaction and an emerging conservation threat to a growing number of species in both terrestrial and marine environments. Although plastic pollution has spread worldwide and a growing body of literature shows its effects on human health, little is known about its impact on our closest living relatives, nonhuman primates, and their habitats. With over 60% of primate species already under threat of extinction, plastic pollution in their habitats poses a unique problem, exposing them to physical harm, synthetic chemicals, and pathogens through ingestion, entanglement, and oral manipulation. Moreover, through its presence in soil, air, and waterways, plastic pollution leads to environmental degradation and reduces the quality and ecological functionality of primate habitats. This perspective article covers what is known so far about plastic pollution as a conservation threat to nonhuman primates. It is a call for primatologists to address plastic pollution in our research and conservation initiatives. By collecting data on plastic pollution's presence and assessing its impact on primates and their habitats, we can develop safe protocols and prevention strategies to combat the threat of plastic pollution in the Anthropocene.

Impact statement

Plastic pollution is a worldwide environmental problem that impacts a wide range of species, ecosystems and landscapes. But, while much attention is given to the consequences of plastic pollution in marine environments (e.g., sea turtles and marine birds eating plastic while foraging), relatively little attention has been given to plastic's effect on terrestrial wildlife. In fact, as the impact of plastic on human health is gaining attention, the effect on our closest relatives and vulnerable species group, nonhuman primates, remains unknown. This article raises awareness of the existing and potential impacts of plastic pollution on the conservation of primates and their habitats and highlights existing evidence of exposure, sources and knowledge gaps. This article provides a starting point for biologists, ecologists, primatologists, and conservationists, summarising the reasons for concern and urgency in exploring the impact of plastic pollution on primates.

Introduction

Plastic pollution is one of the most pressing environmental crises of our time, affecting a growing number of wildlife species worldwide (Santos et al., 2021). As plastic is becoming part of the landscape, turning into a stratigraphic marker likely to enter fossil records, the term *Plasticene* has been suggested as a stage within the Anthropocene and an era in our geological history (Haram et al., 2020; Rangel-Buitrago et al., 2022). The consequences of plastic pollution are most often discussed within the topic of ocean health, as plastic accounts for ~80% of marine pollution alongside an average of 9 million metric tons flowing into the world's ocean every year (Jambeck et al., 2015; Fava, 2022). Numerous plastic pollution awareness campaigns include photographs of sea turtles or marine birds that have become caught in plastic or accidentally ingested it. But while mismanaged plastic waste can eventually reach the oceans, plastic originates on land; it is created, used, and discarded well before reaching the ocean, yet the effects of plastic pollution on terrestrial and freshwater fauna and flora remain understudied and under-monitored (Alimi et al., 2018; de Souza Machado et al., 2018; Bucci et al., 2020). With increasing research on plastic pollution in recent years ecologists are encouraged to focus more attention on plastic pollution's impacts on terrestrial ecosystems and species (de Souza Machado et al., 2018; Blettler and Mitchell, 2021). In marine environments, a concentration of 1.21×10^5 microplastic particles per m^{-3} has been proposed as a minimum threshold; higher levels could lead to significant ecological risks (Everaert et al., 2020; Tekman et al., 2022). A similar threshold has not been

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

 Cambridge
Prisms

 CAMBRIDGE
UNIVERSITY PRESS

determined for terrestrial ecosystems and no comprehensive understanding of the potential ecological risks has been published (Koelmans et al., 2022). Recent studies have documented interactions and ingestion of plastic in various terrestrial mammals and freshwater species (Andrade et al., 2019; Blettler and Mitchell, 2021; Thrift et al., 2022; Ayala et al., 2023). Yet, little is known about the scale and consequences of microplastic exposure, physiological accumulation, and impact on their ecological functions.

Created by humans, plastic and its eventual disposal into the environment may lead to direct and indirect human–wildlife interaction. Pollution, among other human activities, can lead to animal mortality (Gross et al., 2021; Narayan and Rana, 2023), and plastic pollution is no exception. In the environment, plastic undergoes weathering processes (e.g., UV exposure, physical abrasion), fragmentation into smaller particles such as microplastic <5 mm (hereafter, MP) and nanoplastic <100 nm (hereafter, NP) and can be dispersed, transferred and deposited throughout various ecosystems (Barnes, 2009; Alimi et al., 2018). While mismanaged plastic continues to persist and fragment in the environment, it creates a physical hazard when ingested by wildlife, possibly transferring chemicals, and distributing microorganisms up the food chain (Barnes, 2009). The presence and accumulation of plastic in the bodies of humans have been linked to adverse health effects, ranging from reproductive health to immune response and cancer (Wright and Kelly, 2007; Gore et al., 2015). In humans, MP and NP have been found in the blood (Leslie et al., 2022), digestive tract (Schwabl et al., 2019), lungs (Amato-Lourenço et al., 2021), and heart (Yang et al., 2023). Humans ingest between 0.1 and 5 g of microplastic every week, not intentionally, but simply because it is in our environment and food sources (Senathirajah et al., 2021). Most plastic products are manufactured with endocrine-disrupting chemicals (EDCs) with estrogenic activity, such as bisphenol A (BPA) and phthalates, that have adverse effects on human and animal health (Yang et al., 2011). BPA exposure in humans and animals is found to affect development, immune and metabolic systems, and is linked to behavioural changes, obesity, inflammation and reproductive health and fertility in both sexes (Molina-Lopez et al., 2023). Phthalates and BPA have also been linked to decreased fertility, cardiovascular health, and immunological and metabolic disorders in humans (Ramadan et al., 2020; Wang and Qian, 2021). Because we have documented proof of the adverse effects of plastic (and its additives) on human health, we should expand our concern to other species also exposed to microplastics in their environment. This is especially important for our closest living relatives, the nonhuman primates (hereafter, ‘primates’ or ‘NHP’). Many primates are already currently threatened with extinction (Wolfe et al., 2007; Siepel, 2009; Estrada et al., 2017) and plastic pollution only serves to exacerbate their dire situation.

Throughout their evolution, humans and nonhuman primates have shared habitats and formed relationships, some of which remain today, for example, for consumption, tourism, pet keeping and traditional medicine (Fuentes, 2006; Estrada et al., 2017). Primates are the most endangered group of large mammals, constituting 533 species (723 taxa), of which over ~60% are threatened with extinction and 93% with declining populations (Estrada and Garber, 2022; IUCN, 2023). The threats faced by the world’s primates are mostly the results of human activities, for example, habitat loss, fragmentation, disease transmission, hunting and climate change. Pollution is an emerging threat to primates that we are yet to fully understand and include in conservation work (Wich and Marshall, 2016; Estrada et al., 2017). Out of the range of unsustainable

anthropogenic activities, plastic pollution may act as an accelerator to these threats and is already affecting primates.

Plastic pollution is now drawing the attention of primatologists as an emerging threat that must be explored and quantified (Wallis, 2015; Chapman and Peres, 2021). The flexibility and intelligence of primates in adjusting to human habitats often lead to growing interaction with human property and waste, more inclusive diets, increasing access and consumption of human foods in anthropogenic habitats (McLennan et al., 2017). Due to physiological similarities, phylogenetic relationship and cross-species pathogen and disease transmission between humans and primates (Wallis and Lee, 1999; Wolfe et al., 2007), research is needed to understand how primates respond to microplastic accumulation in their bodies. Primates and their population stability also serve as a proxy for ecosystem health and hold a key role in forest regeneration, another reason for protecting them and their habitats from plastic pollution (Chapman, 1995; Stevenson et al., 2002; Estrada et al., 2017).

Sources of plastic pollution in primate habitats

In areas where humans and primates live in close proximity, pollution created by human activities may disperse into primate habitats. Asia and Africa (including Madagascar) are home to 65% of primate species (IUCN, 2023), and more specifically, Southeast Asia and West Africa, regions with high primate species richness, are major hotspots for mismanaged plastic waste and contribution to ocean plastic from land (Maijer et al., 2020). Future projections on the distribution of plastic waste show that even if mitigation efforts occur, Africa and Asia will still cope with disproportionate levels of mismanaged waste (Lebreton and Andrady, 2019).

Exploitation and resource extraction

Extractive industries are known for their destructive impacts on natural habitats. One of the effects of natural resource extractions such as logging, mining and road expansion is environmental pollution and resource contamination (Fuwape, 2003; Kreif et al., 2020; Sun et al., 2021). These could occur as a direct consequence of resource extraction and also through human presence and the disposal of plastic and other solid waste (Fuwape, 2003; Sun et al., 2021). For example, in Uganda’s Kibale National Park the impact of a new road showed a high level of plastic pollution and bisphenol A, an EDC associated with plastic, was found in wild chimpanzee hair for the first time (Kreif et al., 2020).

Indigenous and local communities

The environmental injustice faced by indigenous and local communities (ILC) has consequences for biodiversity and primate conservation, as 71% of primate species’ ranges overlap with indigenous peoples’ lands (Estrada et al., 2022). ILC in developing countries are disproportionately affected by plastic pollution and environmental pollution, as the lack of social justice systems and access to environmental management bodies leads to increased exposure to pollution and its impact on the ecosystem, public health and livelihoods (Fernández-Llamazares et al., 2020; UNEP, 2021). In addition to pollution created by external factors, products packaged in plastic and brought into remote ILC may create an accumulation of solid waste and lead to improper disposal practices such as waste incineration and uncontained landfills (Verma et al., 2016; UNEP, 2021; Figure 1a), and contaminate the habitats of wildlife living alongside them. The burning of plastic waste releases



Figure 1. (a) Himalayan langur (*Semnopithecus ajax*) interacting with a plastic bag in Kanchula, India. Photo by Ryan Ura, The Himalayan Langur Project; (b) Bonnet macaque (*Macaca radiata*) in India. Photo by Janette Wallis; (c) Dead macaque (*Macaca* spp.) suffocated in a plastic bag found on the beach at a popular tourist site, Khao Sam Muk, Thailand. Photo by Phongphat Veeradeetanon; (d) Chimpanzee (*Pan troglodytes schweinfurthii*) interacting with a plastic tarp in Uganda. Photo by Janette Wallis; (e) Pig-tailed macaque (*Macaca nemestrina*) in a polluted mangrove forest, Jakarta, Indonesia, photo by Elisabetta Zavoli; (f) Lion-tailed macaque (*Macaca silenus*) picking food off a plastic bag in a village in India. Photo by Janette Wallis; (g) Long-tailed macaque (*Macaca fascicularis*) with a plastic cup over its head along Thomson Road, Singapore. Photo by Amos Chua; (h) Polluted river bank along the Ucayali River, Peru. Photo by Evelyn D Anca; (i) Celebes crested macaque (*Macaca nigra*) interacting with plastic in Tangkoko Nature Reserve, North Sulawesi, Indonesia. Photo by Meldi/Macaca Nigra Project; (j) White-fronted capuchin (*Cebus albifrons*) eating fruits from a plastic bag 'stolen' from tourists, Puerto Misahuallí, Ecuador. Photo by Adrián Ordieres.

toxic chemicals linked to air pollution and long-term health effects such as cancer and reproductive problems in humans (Verma et al., 2016). Frequent rainfall causes the substances released into the air when plastic is burned to become more incorporated into the soil and the food chain (Ágnes and Rajmund, 2016), and makes tropical and subtropical regions (where more primate species are found) more susceptible to this form of contamination.

Rivers

Land-based plastic pollution carried by rivers is the main source of ocean plastic pollution, with an estimated annual input of 1.15 to 2.41 million tonnes (Lebreton et al., 2017; Schmidt et al., 2017). The highest riverine plastic emissions to the ocean come from 20

countries, most of which are primate range countries (Meijer et al., 2020), and the top 20 most polluting rivers all flow in primate range countries (Lebreton et al., 2017). Because rivers are a key component in primate distribution and diversity (Harcourt, 2012; Naka et al., 2022), the plastic pollution carried by rivers can negatively impact primate ecosystems along the way, becoming trapped in mangroves and flooded forests (do Sul et al., 2014; Bijsterveldt et al., 2021; Figure 1e, h).

Tourism

According to the UNEP, tourism is a major contributor to plastic pollution globally (UNEP, 2023a). Plastic pollution brought by tourists can accumulate, attract wildlife, and potentially disperse

into natural habitats. In addition, feeding wild or habituated primates in their habitats as a touristic activity can increase their exposure to plastic as food packaging may be left behind, break down in the environment, and serve as a fomite for disease transmission, that is, pathogens left behind by humans on their discarded plastic can be transmitted to primates that pick up and inspect these items with their mouth and hands. Curious primates may also 'steal' items from tourists. The impact of waste from tourists and visitors on wild primates is documented in primate range countries such as Thailand (Jones, 2018), India (Krupa, 2021), Singapore (Cheung, 2020) and Ecuador (Figure 1j).

Research

NHP research may be a contributor to plastic pollution in their habitats, most of which are in forests. A survey of primate research field projects indicated reports of increased plastic pollution in and around study sites over recent years (Wallis and Cohen, 2016). Predictably, the survey respondents who worked at sites located deep within a protected forest reported little to no evidence of plastic, while researchers working in unprotected areas or forest fragments indicated a growing problem of plastic pollution. While the exact source of any plastic pollution may be difficult to identify, it is possible that research field staff may contribute to the problem by accidentally leaving plastic in areas where NHPs can access it. In fact, field station trash pits are often raided by nocturnal animals or bold diurnal ones (e.g., baboons) (J. Wallis, pers. obs.). Exposure to plastic is much more likely with more terrestrial than arboreal NHPs.

How does plastic pollution affect primates?

Habitat degradation

Pollution is a major source of habitat degradation, affecting the quality of water, soil and food sources. As a result, the habitat becomes less suitable to support life and leads to a loss of crucial ecosystem services (Adla et al., 2022).

Globally, ~50% of primate species with defined ranges potentially encounter mangrove forest in their habitat, and 147 species were observed to directly use it (Hamilton et al., 2022). Beyond serving as an essential habitat for various species of primates (Nowak, 2012; Gardner, 2016; Hamilton et al., 2022), mangroves hold a substantial role in shoreline protection, water quality, food, shelter and support for over 1,000 species (UNEP, 2023b). With 75% of the world's mangroves being under threat (Azoulay, 2023), one of the many dangers to their survival is pollution. Easily trapped by mangrove forests, plastic waste covers aerial roots and persists in the forest floor and sediment, affecting tree growth in density and size, causing stress reactions, as well as suffocation and death (Suyadi and Manullang, 2020; Bijsterveldt et al., 2021). As plastic pollution accumulates in mangrove forests, it exposes primates to physiological harm and also impacts the health, quality, and growth of their habitats (Figure 1e).

Ecosystem health in primate habitats often starts with the soil. Microplastic accumulation in soil was found to affect its composition and ecological functions (Sajjad et al., 2022). Consumed by soil's flora and fauna and affecting plant growth, it also attracts fungal communities and pathogens travelling up the food chain (Gkoutselis et al., 2021). Microplastic stress in plants was found to affect physiological growth, development and nutrient uptake (Jia et al., 2023). Airborne MP and NP are released into the

atmosphere and travel through wind and rain deposited in soils and water sources contributing to further contamination in surrounding areas (Brahney et al., 2020). Primates, arboreal or terrestrial, heavily depend on plants and fruiting trees in their natural habitats (Chapman and Onderdonk, 1998), and plastic pollution negatively affecting the soil can have an impact on this important co-dependency.

Plastic pollution also reaches rainforests and savannahs. MP was detected in neotropical rainforest and savannah soil in Oaxaca, Mexico, falling in a primate habitat (Álvarez-Lopezello et al., 2021). Plastic pollution was documented in the tropical rainforest along the Ucayali River, Peru where over 10 species of primate live in sympatry and near human settlements (Anca et al., 2023; Figure 1h; Shanee et al., 2023). Microplastic has reached the Himalayas, contaminating the lakes, rivers and downstream communities of humans and wildlife, such as the Himalayan langurs (*Semnopithecus ajax*) (Napper et al., 2020; Talukdar et al., 2023; Figure 1a).

Ingestion and entanglement

As in marine wildlife, plastic waste affects a range of non-marine species through ingestion and entanglement. But while most marine plastic ingestion is accidental (e.g., a sea turtle mistaking a plastic bag for jellyfish), primates are highly intelligent and can carefully explore plastic items for play, exploration, foraging, and unintentionally consume plastic (Wallis, 2015; Figure 1b). As plastic breaks down into MP or NP, and leaches particles and chemicals into air, food and beverages in a wide range of temperatures (Uadia et al., 2019; Mortula et al., 2021), ingestion or inhalation is almost inevitable. Therefore, primates living in close proximity to human settlements, foraging for food in human waste or provisioned by the public or tourists, and manipulating plastic can lead to accidental ingestion of plastic's fragmentation particles and chemical additives or lead to physical injuries (Figure 1c,d,g). Entanglement has been documented in several instances for example: a macaque fatally suffocated inside a plastic bag in Thailand (Sheralyn, 2019; Figure 1c); a macaque hand was trapped in a plastic bottle causing bleeding in Chonburi, Thailand (Yahoo News UK, 2019); and a black howler monkey (*Alouatta caraya*) was found entangled in a fishing net (Blettler and Mitchell, 2021). Ingestion of MP and NP in humans mainly results from the consumption of food contaminated with further exposure through inhalation and skin contact (Domech and Marcos, 2021), and in similar ways could be ingested by NHPs. MP and NP were found in a wide range of human foods, from fruits and vegetables to fish, salt, bottled water, soft drinks and processed foods (Conti et al., 2020; Kwon et al., 2020; Shruti et al., 2020; Lin et al., 2022). Due to the leaching of MP and chemical additives, packaged and ultra-processed foods are concerning sources of MP ingestions and exposure to EDCs (Yang et al., 2011; Buckley et al., 2019; Jadhav et al., 2021). Foraging for food in human garbage dumps can expose primates to plastic through oral interaction with plastic items or food packaging and ingestion of food contaminated with plastic and its chemical additives (Figure 1a,b,f). First documentations of MP in primate digestive system were found in Juruá red howler (*Alouatta juara*) gut content, in the Brazilian Amazon (de Souza Jesus, 2023) and pig-tailed macaque (*Macaca nemestrina*) stool in Indonesia (Suyadi, 2023). In another case, a plastic clothes peg was found in a primate intestine in Bengaluru, India, causing blockage and infection (Prasher, 2023).

Disease transmission

Plastic waste dispersed in the environment and uncontained/unofficial landfills and garbage dumps may lead to primates foraging on human food that may be contaminated with plastic or pathogens that may cause illness (Sapolsky and Share, 2004; Lappan et al., 2020). Plastic has great absorption capabilities taking up a range of environmental pollutants, organic matter, and biomolecules (Rochman et al., 2013). This makes it an effective means of spreading microorganisms (Meng et al., 2021). Thus, plastic can serve as a fomite – transmitting pathogens, including influenza and COVID-19, resulting in disease from humans to NHP and vice versa (Devaux, 2019; Meng et al., 2021). In the post-COVID-19 era, human-primate interactions require careful consideration in our avoidance of disease transmission/risk of zoonosis (Lappan et al., 2020). Plastic pollution therefore should be seen as a form of indirect human-primate interaction and its role in disease transmission should not be overlooked. Wallis and Lee (1999) highlight the need to prevent disease transmission as a major conservation concern; because NHPs are closely related to our own species, they are susceptible to many of the same diseases we carry (Wolfe et al., 2007; Harper et al., 2013). Thus, any pathogen able to be transmitted via plastic is of greater danger to NHPs than to, for example, marine wildlife.

Habitat changes

Behavioural changes, shifts in diet, and modified ranging patterns have been observed in primates as a result of anthropogenic activity (McLennan et al., 2017). Evidence of habitat shift as a result of alternative food sources in open garbage dumps near human settlements was seen in lion-tailed macaques (*Macaca silenus*), posing a risk of dependence on human food (Dhawale and Sinha, 2022). Similarly, olive baboons (*Papio anubis*) were found to shift sleeping sites and foraging exclusively on garbage dumps (Sapolsky and Share, 2004). For many primate species, any substantial shift in habitat use or range can ultimately impact their social structure, reproductive opportunities, and long-term survival.

Chemical additives to plastic

As in humans, exposure to MP, NP, and added EDCs can have long and transgenerational impacts on reproductive health in primates. Studies on primates in laboratory settings show that exposure to EDCs affect reproductive health, cognition, behaviour, and growth in rhesus (*Macaca mulatta*) and long-tailed (*Macaca fascicularis*) macaques (Hunt et al., 2012; Annamalai and Namasivayam, 2015). Bisphenols, chemicals found in plastic, were detected in wild chimpanzee hair in Kibale National Park (Krief et al., 2020). Another study (Krief et al., 2022) showed that captive chimpanzees' exposure to chemical pollutants was even higher, linked to consumption of food and water stored in plastic and interaction with plastic toys. The exposure of wild primate populations to EDCs, transmitted through air or through consumption of contaminated food and water can act as an overlooked 'silent killer' with transgenerational impacts on reproductive health and population stability.

Conclusions

Ecosystem health in primate habitats is vital to the protection and persistence of their populations (Estrada and Garber, 2022).

Primate populations are under growing threat of unsustainable human activities, among them plastic pollution (Chapman and Peres, 2021; Estrada and Garber, 2022). The alarming rate at which plastic pollution contaminates ecosystems makes it urgent to understand how exposure can affect the health of primates as we evaluate the threats they face in the Anthropocene. As plastic pollution continues to spread far from its source of production, primates' exposure to plastic and its associated chemical additives is almost inevitable. We encourage primatologists to incorporate the study of plastic pollution in research and conservation efforts. By collecting data that measures and evaluates any possible threat created by plastics, we can better address the concerns and develop mitigation measures to reduce harm. Despite global efforts to minimise damage from plastic pollution and a growing body of literature warning about its impacts on ecosystems, wildlife, and humans, plastic production is on the rise, with an additional 6 million tonnes produced every year and the 460 million tonnes consumed globally in 2019 is projected to triple by 2060 (OECD, 2022). The accumulation of plastic pollution in the environment, in its many forms and derivatives, is inevitable at this point and expected to increase despite mitigation efforts (Borrelle et al., 2020). While entirely eliminating plastic pollution may not be possible in the near future, a significant reduction in the production of plastic, improved waste management and a shift towards a reuse model is needed on the local, national and global levels (Lau et al., 2020). In the *Plasticene*, where our plastic footprint has entered fossil records and humans and animals live with plastic in their bodies, mitigation and prevention can help reduce the exposure of all living things to adverse effects of plastic and its chemical additives while research and monitoring are crucial to understanding its consequences and implications for conservation.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/plc.2024.10>.

Open peer review. To view the open peer review materials for this article, please visit <http://doi.org/10.1017/plc.2024.10>.

Data availability statement. Data availability is not applicable to this article as no new data were created or analysed in this study.

Acknowledgements. We wish to thank the anonymous reviewers for their feedback and help in improving the quality of this manuscript. We also thank photographers and researchers who have kindly allowed us to use their photos in this paper.

Author contribution. Concept and writing: E.D.A.; Writing and editing: J.W.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing interest. The authors declare none.

References

- Adla K, Dejan K, Neira D and Dragana Š (2022) Degradation of ecosystems and loss of ecosystem services. In Prata JC, Ribeiro AI and Rocha-Santos T (eds.), *One Health*. Cambridge, MA: Academic Press, pp. 281–327. <https://doi.org/10.1016/B978-0-12-822794-7.00008-3>.
- Ágnes N and Rajmund KUTI (2016) The environmental impact of plastic waste incineration. *AARMS—Academic and Applied Research in Military and Public Management Science* 15(3), 231–237. <http://doi.org/10.23937/2572-4061.1510021>.
- Alimi OS, Farner Budarz J, Hernandez LM and Tufenkji N (2018) Microplastics and nanoplastics in aquatic environments: Aggregation, deposition,

- and enhanced contaminant transport. *Environmental Science & Technology* 52(4), 1704–1724. <https://doi.org/10.1021/acs.est.7b05559>.
- Álvarez-Lopezello J, Robles C and del Castillo RF** (2021) Microplastic pollution in neotropical rainforest, savanna, pine plantations, and pasture soils in lowland areas of Oaxaca, Mexico: Preliminary results. *Ecological Indicators* 121, 107084. <https://doi.org/10.1016/j.ecolind.2020.107084>.
- Amato-Lourenço LF, Carvalho-Oliveira R, Júnior GR, dos Santos Galvão L, Ando RA and Mauad T** (2021) Presence of airborne microplastics in human lung tissue. *Journal of Hazardous Materials* 416, 126124. <https://doi.org/10.1016/j.jhazmat.2021.126124>.
- Anca E, Shanee S and Svensson MS** (2023) Ethnoprimateology of the Shipibo of the upper Ucayali River, Perú. *Journal of Ethnobiology and Ethnomedicine* 19(1), 45. <https://doi.org/10.1186/s13002-023-00616-1>.
- Andrade MC, Winemiller KO, Barbosa PS, Fortunati A, Chelazzi D, Cincinelli A and Giarrizzo T** (2019) First account of plastic pollution impacting freshwater fishes in the Amazon: Ingestion of plastic debris by piranhas and other serrasalmids with diverse feeding habits. *Environmental Pollution* 244, 766–773. <https://doi.org/10.1016/j.envpol.2018.10.088>.
- Annamalai J and Namasivayam V** (2015) Endocrine disrupting chemicals in the atmosphere: Their effects on humans and wildlife. *Environment International* 76, 78–97. <https://doi.org/10.1016/j.envint.2014.12.006>.
- Ayala F, Zeta-Flores M, Ramos-Baldarrago S, Tume-Ruiz J, Rangel-Vega A, Reyes E, Quinde E, De-la-Torre GE, Lajo-Salazar L and Cárdenas-Alayza S** (2023) Terrestrial mammals of the Americas and their interactions with plastic waste. *Environmental Science and Pollution Research* 30(20), 57759–57770. <https://doi.org/10.1007/s11356-023-26617-x>.
- Azoulay A** (2023) Message from Ms Audrey Azoulay, Director-General of UNESCO, on the occasion of the International Day for the Conservation of the Mangrove Ecosystem. UNESCO, 26 July. Available at <https://unesdoc.unesco.org/ark:/48223/pf0000386144> (accessed 17 November 2023).
- Barnes DK, Galgani F, Thompson RC and Barlaz M** (2009) Accumulation and fragmentation of plastic debris in global environments. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364(1526), 1985–1998. <https://doi.org/10.1098/rstb.2008.0205>.
- Blettler MC and Mitchell C** (2021) Dangerous traps: Macroplastic encounters affecting freshwater and terrestrial wildlife. *Science of the Total Environment* 798, 149317. <https://doi.org/10.1016/j.scitotenv.2021.149317>.
- Borrelle SB, Ringma J, Law KL, Monnahan CC, Lebreton L, McGivern A, Murphy E, Jambeck J, Leonard GH, Hilleary MA and Eriksen M** (2020) Predicted growth in plastic waste exceeds efforts to mitigate plastic pollution. *Science* 369(6510), 1515–1518. <https://doi.org/10.1126/science.aba3656>.
- Brahney J, Hallerud M, Heim E, Hahnenberger M and Sukumaran S** (2020) Plastic rain in protected areas of the United States. *Science* 368(6496), 1257–1260. <https://doi.org/10.1126/science.aaz5819>.
- Bucci K, Tulio M and Rochman CM** (2020) What is known and unknown about the effects of plastic pollution: A meta-analysis and systematic review. *Ecological Applications* 30(2), 02044. <https://doi.org/10.1002/eap.2044>.
- Buckley JP, Kim H, Wong E and Rebholz CM** (2019) Ultra-processed food consumption and exposure to phthalates and bisphenols in the US National Health and nutrition examination survey, 2013–2014. *Environment International* 131, 105057. <https://doi.org/10.1016/j.envint.2019.105057>.
- Chapman CA** (1995) Primate seed dispersal: Coevolution and conservation implications. *Evolutionary Anthropology: Issues, News, and Reviews* 4(3), 74–82. <https://doi.org/10.1002/evan.1360040303>.
- Chapman CA and Onderdonk DA** (1998) Forests without primates: Primate/plant codependency. *American Journal of Primatology* 45(1), 127–141. [https://doi.org/10.1002/\(sici\)1098-2345\(1998\)45:1%3C127::aid-ajp9%3E3.0.co;2-y](https://doi.org/10.1002/(sici)1098-2345(1998)45:1%3C127::aid-ajp9%3E3.0.co;2-y).
- Chapman CA and Peres CA** (2021) Primate conservation: Lessons learned in the last 20 years can guide future efforts. *Evolutionary Anthropology: Issues, News, and Reviews* 30(5), 345–361. <https://doi.org/10.1002/evan.21920>.
- Cheung R** (2020) Wild monkey sporting bubble tea cup ‘helmet’ a worrying sight: Conservationists. *Asiaone*, 25 August. Available at <https://www.asiaone.com/singapore/wild-monkey-sporting-bubble-tea-cup-helmet-worrying-sight-conservationists> (accessed 17 November 2023).
- Conti GO, Ferrante M, Banni M, Favara C, Nicolosi I, Cristaldi A, Fiore M and Zuccarello P** (2020) Micro- and nano-plastics in edible fruit and vegetables. The first diet risks assessment for the general population. *Environmental Research* 187, 109677. <https://doi.org/10.1016/j.envres.2020.109677>.
- de Souza Jesus A, Nonato F, Cruz AN, Valsecchi J, El Bizri HR, Tregidgo D and Rabelo R** (2023) *Primeira evidência de ingestão de partículas plásticas por um primata arborícola*. Paper presented at the 19^o Simpósio sobre Conservação e Manejo Participativo na Amazônia, 3–7 July, Tefe, Brazil. (Unpublished)
- de Souza Machado AA, Kloas W, Zarfl C, Hempel S and Rillig MC** (2018) Microplastics as an emerging threat to terrestrial ecosystems. *Global Change Biology* 24(4), 1405–1416. <https://doi.org/10.1111/gcb.14020>.
- Devaux CA, Mediannikov O, Medkour H and Raouf D** (2019) Infectious disease risk across the growing human-non human primate interface: A review of the evidence. *Frontiers in Public Health* 7, 305. <https://doi.org/10.3389/fpubh.2019.00305>.
- Dhawal AK and Sinha A** (2022) Ranging patterns of the rainforest-adapted lion-tailed macaque *Macaca silenus* in a human-dominated landscape in the Anamalai hills of the Western Ghats, India. bioRxiv preprint 2022.08.04.502767. <https://doi.org/10.1101/2022.08.04.502767>.
- Domenech J and Marcos R** (2021) Pathways of human exposure to microplastics, and estimation of the total burden. *Current Opinion in Food Science* 39, 144–151. <https://doi.org/10.1016/j.cofs.2021.01.004>.
- Estrada A and Garber PA** (2022) Principal drivers and conservation solutions to the impending primate extinction crisis: Introduction to the special issue. *International Journal of Primatology* 43(1), 1–14. <https://doi.org/10.1007/s10764-022-00283-1>.
- Estrada A, Garber PA, Rylands AB, Roos C, Fernandez-Duque E, Di Fiore A, Nekaris KAI, Nijman V, Heymann EW, Lambert JE and Rovero F** (2017) Impending extinction crisis of the world’s primates: Why primates matter. *Science Advances* 3(1), e1600946. <https://doi.org/10.1126/sciadv.1600946>.
- Everaert G, De Rijcke M, Lonnevillle B, Janssen CR, Backhaus T, Mees J, van Sebillie E, Koelmans AA, Catarino AI and Vandegheuchte MB** (2020) Risks of floating microplastic in the global ocean. *Environmental Pollution* 267, 115499. <https://doi.org/10.1016/j.envpol.2020.115499>.
- Fava M** (2022) Ocean plastic pollution an overview: Data and statistics. UNESCO Ocean Literacy Portal, 9 May. Available at <https://oceanliteracy.unesco.org/plastic-pollution-ocean/> (accessed 21 September 2023).
- Fernández-Llamazares Á, Garteizgogea M, Basu N, Brondizio ES, Cabeza M, Martínez-Alier J, McElwee P and Reyes-García V** (2020) A state-of-the-art review of indigenous peoples and environmental pollution. *Integrated Environmental Assessment and Management* 16(3), 324–341. <https://doi.org/10.1002/ieam.4239>.
- Fuentes A** (2006) Human-nonhuman primate interconnections and their relevance to anthropology. *Ecological and Environmental Anthropology* 2(2), 1–11. Available at <https://digitalcommons.unl.edu/icwdmeea/1>. (accessed 21 September 2023).
- Fuwape JA** (2003) The impacts of forest industries and wood utilization on the environment. In *Paper Presented at the XII World Forestry Congress*, 21–28 September. Quebec, Canada: FAO. Available at <https://www.fao.org/3/xii/0122-a2.htm>. (accessed 17 November 2023).
- Gardner CJ** (2016) Use of mangroves by lemurs. *International Journal of Primatology* 37(3), 317–332. <https://doi.org/10.1007/s10764-016-9905-1>.
- Gkoutselis G, Rohrbach S, Harjes J, Obst M, Brachmann A, Horn MA and Rambold G** (2021) Microplastics accumulate fungal pathogens in terrestrial ecosystems. *Scientific Reports* 11(1), 13214. <https://doi.org/10.1038/s41598-021-92405-7>.
- Gore AC, Chappell VA, Fenton SE, Flaws JA, Nadal A, Prins GS, Toppari J and Zoeller RT** (2015) EDC-2: The Endocrine Society’s second scientific statement on endocrine-disrupting chemicals. *Endocrine Reviews* 36(6), E1–E150. <https://doi.org/10.1210/er.2015-1010>.
- Gross E, Jayasinghe N, Brooks A, Polet G, Wadhwa R and Hilderink-Koopmans F** (2021) *A Future for all: The Need for Human-Wildlife Coexistence*. Gland, Switzerland: WWF. Available at https://wwfint.awsassets.panda.org/downloads/a_future_for_all_the_need_for_human_wildlife_coexistence.pdf (accessed 25 March 2024).
- Hamilton SE, Presotto A and Lembo AJ** (2022) Establishing the relationship between non-human primates and mangrove forests at the global, national,

- and local scales. *PLoS One* 17(11), e0277440. <https://doi.org/10.1371/journal.pone.0277440>.
- Haram LE, Carlton JT, Ruiz GM and Maximenko NA (2020) A plasticene lexicon. *Marine Pollution Bulletin* 150, 110714. <https://doi.org/10.1016/j.marpolbul.2019.110714>.
- Harcourt AH and Wood MA (2012) Rivers as barriers to primate distributions in Africa. *International Journal of Primatology* 33, 168–183. <http://doi.org/10.1007/s10764-011-9558-z>.
- Harper KN, Zuckerman MK, Turner BL and Armelagos GJ (2013) Primates, pathogens, and evolution: A context for understanding emerging disease. In Brinkworth JF and Pechenikina K (eds.), *Primates, Pathogens, and Evolution*. New York: Springer, pp. 389–409. <https://doi.org/10.1007/978-1-4614-7181-3>.
- Hunt PA, Lawson C, Gieske M, Murdoch B, Smith H, Marre A, Hassold T and VandeVoort CA (2012) Bisphenol a alters early oogenesis and follicle formation in the fetal ovary of the rhesus monkey. *Proceedings of the National Academy of Sciences* 109(43), 17525–17530. <https://doi.org/10.1073/pnas.1207854109>.
- IUCN Primate Specialist Group (2023) Global non-human primate diversity, 15 February 2023. Available at http://www.primatesg.org/primates_diversity_by_region/ (accessed 30 October 2023).
- Jadhav EB, Sankhla MS, Bhat RA and Bhagat DS (2021) Microplastics from food packaging: An overview of human consumption, health threats, and alternative solutions. *Environmental Nanotechnology, Monitoring & Management* 16, 100608. <http://doi.org/10.1016/j.enmm.2021.100608>.
- Jambeck JR, Geyer R, Wilcox C, Siegler TR, Perryman M, Andrady A, Narayan R and Law KL (2015) Plastic waste inputs from land into the ocean. *Science* 347(6223), 768–771. <https://doi.org/10.1126/science.1260352>.
- Jia L, Liu L, Zhang Y, Fu W, Liu X, Wang Q, Tanveer M and Huang L (2023) Microplastic stress in plants: Effects on plant growth and their remediations. *Frontiers in Plant Science* 14, 1226484. <https://doi.org/10.3389/fpls.2023.1226484>.
- Jones M (2018) Monkeys choke inside carrier bags in Thailand's plastic waste crisis. *The Mirror*, 10 August. Available at <https://www.mirror.co.uk/news/world-news/thailand-plastic-waste-crisis-monkeys-13062464> (accessed 30 October 2023).
- Koelmans AA, Redondo-Hasselerharm PE, Nor NH, de Ruijter VN, Mintenig SM and Kooi M (2022) Risk assessment of microplastic particles. *Nature Reviews Materials* 7(2), 138–152. <https://doi.org/10.1038/s41578-021-00411-y>.
- Krief S, Iglesias-González A, Appenzeller BM, Okimat JP, Fini JB, Demeneix B, Vaslin-Reimann S, Lardy-Fontan S, Guma N and Spirhanzlova P (2020) Road impact in a protected area with rich biodiversity: The case of the Sebitali road in Kibale National Park, Uganda. *Environmental Science & Pollution Research* 27(22), 27914. <https://doi.org/10.1007/s11356-020-09098-0>.
- Krief S, Iglesias-González A, Appenzeller BM, Rachid L, Beltrame M, Asalu E, Okimat JP, Kane-Maguire N and Spirhanzlova P (2022) Chimpanzee exposure to pollution revealed by human biomonitoring approaches. *Ecotoxicology and Environmental Safety* 233, 113341. <https://doi.org/10.1016/j.ecoenv.2022.113341>.
- Krupa PL (2021) Monkey see monkey do. *Bangalore Mirror*, 7 October. Available at <https://bangaloremirror.indiatimes.com/bangalore/cover-story/monkey-see-monkey-do/articleshow/86826708.cms> (accessed 19 December 2023).
- Kwon JH, Kim JW, Pham TD, Tarafdar A, Hong S, Chun SH, Lee SH, Kang DY, Kim JY, Kim SB and Jung J (2020) Microplastics in food: A review on analytical methods and challenges. *International Journal of Environmental Research and Public Health* 17(18), 6710. <https://doi.org/10.3390/ijerph17186710>.
- Lappan S, Malaivijitnond S, Radhakrishna S, Riley EP and Ruppert N (2020) The human–primate interface in the New Normal: Challenges and opportunities for primatologists in the COVID-19 era and beyond. *American Journal of Primatology* 82(8), e23176. <https://doi.org/10.1002/ajp.23176>.
- Lau WW, Shiran Y, Bailey RM, Cook E, Stuchtey MR, Koskella J, Velis CA, Godfrey L, Boucher J, Murphy MB and Thompson RC (2020) Evaluating scenarios toward zero plastic pollution. *Science* 369(6510), 1455–1461. <https://doi.org/10.1126/science.aba9475>.
- Lebreton L and Andrady A (2019) Future scenarios of global plastic waste generation and disposal. *Palgrave Communications* 5(1), 1–11. <https://doi.org/10.1057/s41599-018-0212-7>.
- Lebreton LC, Van Der Zwet J, Damsteeg JW, Slat B, Andrady A and Reisser J (2017) River plastic emissions to the world's oceans. *Nature Communications* 8(1), 15611. <https://doi.org/10.1038/ncomms15611>.
- Leslie HA, Van Velzen MJ, Brandsma SH, Vethaak AD, Garcia-Vallejo JJ and Lamoree MH (2022) Discovery and quantification of plastic particle pollution in human blood. *Environment International* 163, 107199. <https://doi.org/10.1016/j.envint.2022.107199>.
- Lin Q, Zhao S, Pang L, Sun C, Chen L and Li F (2022) Potential risk of microplastics in processed foods: Preliminary risk assessment concerning polymer types, abundance, and human exposure of microplastics. *Ecotoxicology and Environmental Safety* 247, 114260. <https://doi.org/10.1016/j.ecoenv.2022.114260>.
- McLennan MR, Spagnoletti N and Hockings KJ (2017) The implications of primate behavioral flexibility for sustainable human–primate coexistence in anthropogenic habitats. *International Journal of Primatology* 38, 105–121. <https://doi.org/10.1007/s10764-017-9962-0>.
- Meijer LJ, Van Emmerik T, Van Der Ent R, Schmidt C and Lebreton L (2020) More than 1000 rivers account for 80% of global riverine plastic emissions into the ocean. *Science Advances* 7(18), eaaz5803. <https://doi.org/10.1126/sciadv.aaz5803>.
- Meng J, Zhang Q, Zheng Y, He G and Shi H (2021) Plastic waste as the potential carriers of pathogens. *Current Opinion in Food Science* 41, 224–230. <http://doi.org/10.1016/j.cofs.2021.04.016>.
- Molina-López AM, Bujalance-Reyes F, Ayala-Soldado N, Mora-Medina R, Lora-Benítez A and Moyano-Salvago R (2023) An overview of the health effects of Bisphenol a from a one health perspective. *Animals* 13(15), 2439. <https://doi.org/10.3390/ani13152439>.
- Mortula MM, Atabay S, Fattah KP and Madbully A (2021) Leachability of microplastic from different plastic materials. *Journal of Environmental Management* 294, 112995. <https://doi.org/10.1016/j.jenvman.2021.112995>.
- Naka LN, Werneck FP, Rosser N, Pil MW and Boubli JP (2022) The role of rivers in the origins, evolution, adaptation, and distribution of biodiversity. *Frontiers in Ecology and Evolution* 10, 1035859. <https://doi.org/10.3389/fevo.2022.1035859>.
- Napper IE, Davies BF, Clifford H, Elvin S, Koldewey HJ, Mayewski PA, Miner KR, Potocki M, Elmore AC, Gajurel AP and Thompson RC (2020) Reaching new heights in plastic pollution—Preliminary findings of microplastics on Mount Everest. *One Earth* 3(5), 621–630. <https://doi.org/10.1016/j.oneear.2020.10.020>.
- Narayan E and Rana N (2023) Human–wildlife interaction: Past, present, and future. *BMC Zoology* 8(1), 5. <https://doi.org/10.1186/s40850-023-00168-7>.
- Nowak K (2012) Mangrove and peat swamp forests: Refuge habitats for primates and felids. *Folia Primatologica* 83(3–6), 361–376. <https://doi.org/10.1159/000339810>.
- OECD (2022) Global plastic waste set to almost triple by 2060, says OECD. OECD, 3 June. Available at <https://www.oecd.org/environment/global-plastic-waste-set-to-almost-triple-by-2060.htm> (accessed 16 November 2023).
- Prasher G (2023) Plastic goes wild. *Bangalore Mirror*, 2 July. Available at <https://bangaloremirror.indiatimes.com/bangalore/others/plastic-goes-wild/article-show/101422562.cms> (accessed 16 November 2023).
- Ramadan M, Cooper B and Posnack NG (2020) Bisphenols and phthalates: Plastic chemical exposures can contribute to adverse cardiovascular health outcomes. *Birth Defects Research* 112(17), 1362–1385. <https://doi.org/10.1002/bdr2.1752>.
- Rangel-Buitrago N, Neal W and Williams A (2022) The plasticene: Time and rocks. *Marine Pollution Bulletin* 185, 114358. <https://doi.org/10.1016/j.marpolbul.2022.114358>.
- Rochman CM, Hoh E, Hentschel BT and Kaye S (2013) Long-term field measurement of sorption of organic contaminants to five types of plastic pellets: Implications for plastic marine debris. *Environmental Science & Technology* 47(3), 1646–1654. <https://doi.org/10.1021/es303700s>.
- Sajjad M, Huang Q, Khan S, Khan MA, Liu Y, Wang J, Lian F, Wang Q and Guo G (2022) Microplastics in the soil environment: A critical review. *Environmental Technology & Innovation* 27, 102408. <https://doi.org/10.1016/j.eti.2022.102408>.
- Santos RG, Machovsky-Capuska GE and Andrades R (2021) Plastic ingestion as an evolutionary trap: Toward a holistic understanding. *Science* 373(6550), 56–60. <https://doi.org/10.1126/science.abh0945>.

- Sapolsky RM and Share LJ (2004) A pacific culture among wild baboons: Its emergence and transmission. *PLoS Biology* 2(4), e106. <https://doi.org/10.1371/journal.pbio.0020106>.
- Schmidt C, Krauth T and Wagner S (2017) Export of plastic debris by rivers into the sea. *Environmental Science & Technology* 51(21), 12246–12253. <https://doi.org/10.1021/acs.est.7b02368>.
- Schwabl P, Köppel S, Königshofer P, Bucsecs T, Trauner M, Reiberger T and Liebmann B (2019) Detection of various microplastics in human stool: A prospective case series. *Annals of Internal Medicine* 171(7), 453–457. <https://doi.org/10.7326/m19-0618>.
- Senathirajah K, Attwood S, Bhagwat G, Carbery M, Wilson S and Palanisami T (2021) Estimation of the mass of microplastics ingested – A pivotal first step towards human health risk assessment. *Journal of Hazardous Materials* 404, 124004. <https://doi.org/10.1016/j.jhazmat.2020.124004>.
- Shanee S, Fernández-Hidalgo L, Walford J, Fernandez-Hilarío R, Alarcon A, Llaja KGS and Allgas N (2023) Surveys of a high-diversity primate Community in a Forestry Concession, Ucayali, Peru. *Primate Conservation* 37, 1–14. Available at http://www.primate-sg.org/storage/pdf/PC37_Shanee_et_al_Peru_primate_community_survey.pdf.
- Sheralyn (2019) Heartbreaking photos show dead monkey inside plastic bag after it drowned & bled from nose. World of Buzz, 20 November. Available at <https://worldofbuzz.com/heartbreaking-photos-show-dead-monkey-inside-plastic-bag-after-it-drowned-bled-from-nose/> (accessed 16 November 2022).
- Shruti VC, Pérez-Guevara F, Elizalde-Martínez I and Kutralam-Muniasamy G (2020) First study of its kind on the microplastic contamination of soft drinks, cold tea and energy drinks-future research and environmental considerations. *Science of the Total Environment* 726, 138580. <https://doi.org/10.1016/j.scitotenv.2020.138580>.
- Siepel A (2009) Phylogenomics of primates and their ancestral populations. *Genome Research* 19(11), 1929–1941. <https://doi.org/10.1101/gr.084228.108>.
- Stevenson PR, Castellanos MC, Pizarro JC and Garavito M (2002) Effects of seed dispersal by three ateline monkey species on seed germination at Tinigua National Park, Colombia. *International Journal of Primatology* 23, 1187–1204. <https://doi.org/10.1023/A:1021118618936>.
- Sul JAI, Costa MF, Silva-Cavalcanti JS and Araújo MCB (2014) Plastic debris retention and exportation by a mangrove forest patch. *Marine Pollution Bulletin* 78(1–2), 252–257. <https://doi.org/10.1016/j.marpolbul.2013.11.011>.
- Sun Y, Li Y, Yu T, Zhang X, Liu L and Zhang P (2021) Resource extraction, environmental pollution and economic development: Evidence from prefecture-level cities in China. *Resources Policy* 74, 102330. <https://doi.org/10.1016/j.resourpol.2021.102330>.
- Suyadi (2023) *Plastic Pollution in Bukit Barisan Selatan National Park, Its Implications, and Public Awareness*. National Research and Innovation Agency Indonesia, BRIN.
- Suyadi and Manullang CY (2020) Distribution of plastic debris pollution and its implications on mangrove vegetation. *Marine Pollution Bulletin* 160, 111642. <https://doi.org/10.1016/j.marpolbul.2020.111642>.
- Talukdar A, Bhattacharya S, Bandyopadhyay A and Dey A (2023) Microplastic pollution in the Himalayas: Occurrence, distribution, accumulation and environmental impacts. *Science of the Total Environment* 874, 162495. <https://doi.org/10.1016/j.scitotenv.2023.162495>.
- Tekman MB, Walther BA, Peter C, Gutow L and Bergmann M (2022) *Impacts of Plastic Pollution in the Oceans on Marine Species, Biodiversity and Ecosystems*. Berlin: WWF. <https://doi.org/10.5281/zenodo.5898684>.
- Thrift E, Porter A, Galloway TS, Coomber FG and Mathews F (2022) Ingestion of plastics by terrestrial small mammals. *Science of the Total Environment* 842, 156679. <https://doi.org/10.1016/j.scitotenv.2022.156679>.
- Uadia PO, Makinwa TT and Akeshinro AA (2019) Leaching of bisphenol A (BPA) from plastic materials increases with storage time while its degradation increases with temperature. *Salem University Journal of Environmental Sciences* 1(1), 1–8.
- United Nations Environment Programme (UNEP)(2021) *Neglected: Environmental Justice Impacts of Marine Litter and Plastic Pollution*. Nairobi. Available at <https://wedocs.unep.org/bitstream/handle/20.500.11822/35417/EJIPP.pdf> (accessed 21 September 2023).
- United Nations Environment Programme (UNEP)(2023a) How can tourism fix its plastic problem? UNEP, 12 May. Available at <http://www.unep.org/news-and-stories/story/how-can-tourism-fix-its-plastic-problem>. (accessed 21 September 2023).
- United Nations Environment Programme (UNEP)(2023b) *Decades of mangrove forest change: What does it mean for nature, people and the climate?* Nairobi. Available at https://wedocs.unep.org/bitstream/handle/20.500.11822/42254/mangrove_forest_change.pdf?sequence=3&isAllowed=y.
- van Bijsterveldt CE, van Wesenbeeck BK, Ramadhani S, Raven OV, van Gool FE, Pribadi R and Bouma TJ (2021) Does plastic waste kill mangroves? A field experiment to assess the impact of macro plastics on mangrove growth, stress response and survival. *Science of the Total Environment* 756, 143826. <https://doi.org/10.1016/j.scitotenv.2020.143826>.
- Verma R, Vinoda KS, Papireddy M and Gowda ANS (2016) Toxic pollutants from plastic waste-a review. *Procedia Environmental Sciences* 35, 701–708. <https://doi.org/10.1016/j.proenv.2016.07.069>.
- Wallis J (2015) *Plastics endanger primates: A field report*. Plastic Pollution Coalition blog. Available at <https://www.plasticpollutioncoalition.org/blog/2015/8/23/plastic-endanger-primates-a-field-report> (accessed 21 September 2023).
- Wallis J and Cohen D (2016) *Plastic pollution and primates: A survey of field sites*. In *Paper Presented at the International Primatological Society Meeting, 21–27 August, Chicago, IL* (Unpublished).
- Wallis J and Lee DR (1999) Primate conservation: The prevention of disease transmission. *International Journal of Primatology* 20, 803–826. <https://doi.org/10.1023/A:102087970028>.
- Wang Y and Qian H (2021) Phthalates and their impacts on human health. *Healthcare* 9(5), 603. <https://doi.org/10.3390/healthcare9050603>.
- Wich SA and Marshall AJ (2016) *An Introduction to Primate Conservation*. Oxford: Oxford University Press.
- Wolfe ND, Dunavan CP and Diamond J (2007) Origins of major human infectious diseases. *Nature* 447(7142), 279–283. <https://doi.org/10.1038/nature05775>.
- Wright SL and Kelly FJ (2007) Plastic and human health: A micro issue? *Environmental Science & Technology* 51(12), 6634–6647. <https://doi.org/10.1021/acs.est.7b00423>.
- Yahoo News UK (2019) *Heartbreaking Moment Wild Monkey Is Found with Plastic Bottle Stuck on His Wrist*. Yahoo News UK, 15 March. <https://uk.news.yahoo.com/heartbreaking-moment-wild-monkey-found-100400006.html> (accessed 21 September 2023).
- Yang Y, Xie E, Du ZP, Han Z, Li Z, Zhao L, Qin R, Xue Y, Li F M and Hua K (2023) Detection of various microplastics in patients undergoing cardiac surgery. *Environmental Science & Technology* 57(30), 10911–10918. <http://doi.org/10.1021/acs.est.2c07179>.
- Yang CZ, Yaniger SI, Jordan VC, Klein DJ and Bittner GD (2011) Most plastic products release estrogenic chemicals: A potential health problem that can be solved. *Environmental Health Perspectives* 119(7), 989–996. <https://doi.org/10.1289/ehp.1003220>.