

Nonlinear drift-wave and energetic particle long-time behaviour in stellarators: solution of the kinetic problem

Alessandro Zocco^{1,†}, Alexey Mishchenko¹, Axel Könies¹,
Matteo Falessi² and Fulvio Zonca^{2,3}

¹Max Planck Institute for Plasma Physics, 17491 Greifswald, Germany

²Center for Nonlinear Plasma Science and ENEA C. R. Frascati, Frascati, Italy

³Institute for Fusion Theory and Simulation and Department of Physics, Zhejiang University, Hangzhou, PR China

(Received 2 November 2022; revised 9 May 2023; accepted 11 May 2023)

We propose a theoretical scheme for the study of the nonlinear interaction of drift-wave-like turbulence and energetic particles in stellarators. The approach is based on gyrokinetics, and features a separation of time and scales, for electromagnetic fluctuations, inspired by linear ballooning theory. Two specific moments of the gyrokinetic equation constitute the main equations of the system, which requires a full kinetic nonlinear solution. This is found iteratively, expanding in the smallness of the bounce-average radial drift frequency, and nonlinear $\mathbf{E} \times \mathbf{B}$ drift frequency, compared with the inverse time scales of the resonantly interacting energetic particles. Our analysis is therefore valid for neoclassically optimised stellarators. The resummation of all iterative and perturbative nonlinear kinetic solutions is discussed in terms of Feynman diagrams. Particular emphasis is put on the role of collisionlessly undamped large-scale structures in phase space, the kinetic equivalent of zonal flows, i.e. phase-space zonal structures, and on wave-like fluctuations generated by energetic particles.

Key words: fusion plasma, plasma nonlinear phenomena

1. Introduction

The purpose of any magnetic confinement fusion device is to access energetically favourable scenarios where fusion reactions are self-sustained by alpha particles. This has not been attained yet, but the study of energetic particle minorities, produced by auxiliary heating, gives us a glimpse of what could be the behaviour of a burning plasma.

A burning plasma is expected to be a kinetic turbulent environment in which energetic particles experience resonant interactions with fluctuating electromagnetic fields. These are believed to be faithfully described by gyrokinetic theory (Frieman & Chen 1982). Here, fluctuations are assumed to vary slowly along and strongly across the confining

† Email address for correspondence: alessandro.zocco@ipp.mpg.de

magnetic field, while evolving on a time scale which is slow when compared with the gyromotion, but fast when compared with the long-time behaviour of the plasma density and temperature background. Thus, within a gyrokinetic description, plasma transport properties are determined through a turbulence average (in time and space) (Frieman & Chen 1982; Abel & Cowley 2013) which allows nonlinear fluxes to act as sources in the evolution of mean quantities. Thereby, in a sense, nonlinear gyrokinetic simulations provide insight into the detailed properties of kinetic turbulence in magnetised plasmas, but do not fully characterise them from the point of view of transport, and therefore possible reactor performance, unless one knows how fluctuations, on a long time scale, impact mean quantities such as temperature and density.

Among the many approaches that one can pursue to attack the problem of turbulent transport in magnetically confined plasmas (which is naturally multiscale, and tackled with an asymptotic expansion for small Larmor radii), the one reviewed by Chen & Zonca (2019), perhaps more than others, kept the physics of fast particles at heart. In the first of a series of works there cited (Zonca *et al.* 2015), the question was proposed, and partially addressed, as to what would be the dominant contribution to the possible long-time behaviour of gyrokinetic fluctuations driven by wave–particle resonant interaction. By exploiting the existence of action-angle variables in axisymmetric toroidal geometry, Zonca *et al.* (2015) introduced a way of integrating perturbed particle orbits by using the unperturbed action-angle variables, thus capturing cumulative wave–particle resonance effects of bounce- and transit-averaged processes on unperturbed particle motion, for sufficiently small nonlinear dynamics. This approach allows one to describe several nonlinear wave–particle phenomena, such as phase locking, resonance detuning and broadening. It gives a powerful means of predicting chorus emission in planetary magnetospheres (Zonca, Tao & Chen 2021*b*), but it also gives the basis for the study of what the authors call phase-space zonal structures, i.e. the counterpart of zonal flows (Rosenbluth & Hinton 1998) or, more generally, zonal field structures in phase space. These are structures that are constant over magnetic flux surfaces, thus, in many circumstances, immune to collisionless Landau damping. They play a major role in the regulation of turbulence levels (Rogers, Dorland & Kotschenreuther 2000; Diamond *et al.* 2005).

Further advancing this line of research, Falessi & Zonca (2019) have focused on the long-time behaviour of phase-space zonal structures in axisymmetry. Their results have been successfully applied to particle-in-cell codes (Bottino *et al.* 2022), and completed with the evaluation of the mesoscale radial extent of fluctuations (Zonca *et al.* 2021*a*). These are evaluated as first-order corrections with a subsidiary limit of gyrokinetics that mimics the original radial envelope evaluation of ballooning modes (Connor, Hastie & Taylor 1978), Alfvén waves (Zonca & Chen 1992, 1993) and the ion-temperature-gradient-driven modes (Romanelli & Zonca 1993).

We are starting to see the first numerical results on the behaviour of energetic particles embedded into turbulence (Biancalani *et al.* 2021; Di Siena *et al.* 2021; Carlevaro *et al.* 2022; Hayward-Schneider *et al.* 2022; Lu *et al.* 2023), but, to date, the aforementioned fundamental theoretical aspects have only been covered in the context of tokamak physics (Zonca *et al.* 2015; Chen & Zonca 2016, 2019; Falessi & Zonca 2019). A generalisation to stellarators is not straightforward mainly for two reasons. The first being that within a multiscale gyrokinetic approach in stellarators, the strong variation of the strength of the equilibrium magnetic field on flux surfaces is very likely more important than the radial variation of thermodynamic equilibrium quantities (Zocco *et al.* 2016; Zocco, Aleynikova & Xanthopoulos 2018*a*; Zocco, Plunk & Xanthopoulos 2020), the latter being the most important mesoscale effect encountered in asymmetry, and taken into account

when evaluating the fluctuation radial envelope. The second problem encountered when extending the work of Falessi & Zonca (2019) to stellarators is the absence of a conserved toroidal canonical momentum, which complicates considerably the analysis.

Nevertheless, in this work, we find a way of extending to stellarators the kinetic theory first put forth by Zonca *et al.* (2015). Whilst realising that we are engaging with innumerable aspects of a complex endeavour, we decide to particularly focus on the kinetic solution of the problem, devising a method, given the electromagnetic fields, to solve iteratively the gyrokinetic equation while retaining the nonlinear wave–particle interaction. The method is diagrammatic, and applies not only to the physics of energetic particles, but also to the study of micro-instability-driven transport as the electrostatic ion temperature gradient at marginality (Zocco *et al.* 2018*b*), for instance, when wave–particle resonance effects are important.

This work is organised as follows. In § 2 we introduce the basic gyrokinetic equation and the equations for fluctuating fields and set up the relation between wave-like fluctuations and the underlying kinetics. In § 3 we introduce the coordinates and specify some geometric features that allow us to solve the gyrokinetic equation. In § 4 we introduce its iterative diagrammatic solution. In §§ 5 and 6 we apply the kinetic solution to the evaluation of transport due to wave-like fluctuating fields, and the phase-space zonal structure. In § 7 we conclude.

2. Nonlinear transport from finite- β fluctuations

In this section we highlight the mathematical formulation for the study of transport from electromagnetic fluctuations. Our starting point is the evaluation of the quasi-neutrality equation and the divergence of the perturbed plasma current (Frieman *et al.* 1980; Tang, Connor & Hastie 1980; Qin, Tang & Rewoldt 1998; Zonca & Chen 2014; Chen & Zonca 2016; Zocco, Helander & Connor 2015; Aleynikova & Zocco 2017):

$$\sum_s q_s \int d^3 \mathbf{v} F_{0s} \frac{q_s \phi}{T_{0s}} = \sum_s q_s \int d^3 \mathbf{v} \sum_k e^{ik \cdot \mathbf{r}} J_0(a_s) \delta G_{sk} \quad (2.1)$$

and

$$\begin{aligned} B \nabla_{\parallel} \frac{j_{\parallel, k}}{B} = & - \sum_s \frac{e_s^2}{T_{0s}} \int d^3 \mathbf{v} F_{0s} \left[(1 - J_0^2(a_s)) \frac{\partial}{\partial t} - i \Omega_{*s}^T J_0^2(a_s) \right] \phi_k \\ & - \sum_s \frac{e_s^2}{T_s} \int d^3 \mathbf{v} F_{0s} \left(\frac{\partial}{\partial t} - i \Omega_{*s}^T \right) 4 \mu B \frac{J_0(a_s) J_1(a_s)}{a_s} \frac{T_{0s}}{q_s} \frac{\delta B_{\parallel k}}{B} \\ & + i \frac{c}{B} \hat{\mathbf{b}} \cdot \boldsymbol{\kappa} \times \mathbf{k} \sum_s \int d^3 \mathbf{v} m_s (\mu B + v_{\parallel}^2) J_0(a_s) \delta G_{sk} \\ & + \sum_s q_s \int d^3 \mathbf{v} J_0(a_s) \sum_{k'} \frac{c}{B} \mathbf{b} \cdot \mathbf{k}'_{\perp} \times \mathbf{k}_{\perp} \left[J_0(a'_s) \left(\phi_{k'} - \frac{v_{\parallel}}{c} A_{\parallel k'} \right) \right. \\ & \left. + 4 \mu B \frac{J_1(a'_s)}{a'_s} \frac{T_{0s}}{q_s} \frac{\delta B_{\parallel k'}}{B} \right] \delta G_{s, k-k'} + B \sum_s \int d^3 \mathbf{v} \delta G_k \nabla_{\parallel} J_0(a_s), \quad (2.2) \end{aligned}$$

where F_{0s} is the equilibrium distribution function (taken to be Maxwellian), thus $f_s = F_{0s} + \delta f_s \equiv F_{0s} (1 - q_s \phi(\mathbf{r}, t) / T_{0s}) + \delta G_s(\mathbf{R}_s, \mu, \mathcal{E}, t) + \mathcal{O}(\epsilon^2)$, and $\delta f_s / F_{0s} \sim k_{\parallel} / k_{\perp} \sim \epsilon \equiv \rho_* = \rho_s / a$, where ρ_s is the Larmor radius and a a macroscopic length.

Here $\mathbf{R}_s = \mathbf{r} + \mathbf{v}_\perp \times \hat{\mathbf{b}}/\Omega_s$ is the gyrocentre position, where \mathbf{r} is the particle position, $\Omega_s = q_s B/(m_s c)$, $\hat{\mathbf{b}} = \mathbf{B}/B$, with \mathbf{B} the equilibrium magnetic field, and $\mu = v_\perp^2/(2B)$, $\mathcal{E} = v^2/2$ the velocity-space coordinates. The Bessel function, $J_0 = J_0(a_s)$, with $a_s^2 = 2B\mu k_\perp^2/\Omega_s^2$, relates the Fourier transform with respect to \mathbf{R}_s of δG_s with its Fourier transform with respect to \mathbf{r} (velocity-space integrals in (2.1)–(2.2) are taken at constant \mathbf{r}). We also have

$$\Omega_{*,s}^T = \frac{cT_{0s}}{Be_s} \hat{\mathbf{b}} \cdot \mathbf{k} \times \frac{\nabla F_{0s}}{F_{0s}}, \tag{2.3}$$

which takes account of diamagnetic effects, while $\boldsymbol{\kappa} = \hat{\mathbf{b}} \cdot \nabla \hat{\mathbf{b}}$ is the magnetic field curvature. Finally,

$$\frac{\delta B_{\parallel,k}}{B} = -\frac{4\pi}{c} \sum_s \int d^3 v m_s 2\mu B \frac{J_1(a_s)}{a_s} \delta G_{s,k} \tag{2.4}$$

and

$$j_{\parallel k} = \sum_s q_s \int d^3 v v_{\parallel} J_0(a_s) \delta G_{s,k}, \tag{2.5}$$

where $j_{\parallel} = (c/4\pi) \hat{\mathbf{b}} \cdot \nabla \times (\nabla \times \mathbf{A})$. We take $\boldsymbol{\kappa} \approx \nabla B/B$, consistent with linear cancellations due to the perpendicular pressure balance (Tang *et al.* 1980; Hasegawa & Sato 1981; Zocco *et al.* 2015; Chen & Zonca 2016; Aleynikova & Zocco 2017), thus allowing us to eliminate (but not neglect!) magnetic compressibility altogether. The final term in (2.2) can be neglected in most practical applications, since the current is mostly carried by the electrons (Connor *et al.* 1978; Zocco *et al.* 2015; Aleynikova & Zocco 2017). The kinetic information is contained in the velocity-space integrals of the function δG_s .

The function δG_s follows the nonlinear Frieman–Chen equation (Frieman & Chen 1982) for the non-adiabatic part of the perturbed distribution function, $\delta G_s = \delta f_s + F_{0s} e_s \phi/T_{0s}$:

$$\begin{aligned} & \left(\frac{\partial}{\partial t} + v_{\parallel} \nabla_{\parallel} + \mathbf{v}_{d,s} \cdot \nabla \right) \delta G_s \\ &= \frac{e_s F_{0s}}{T_{0s}} \frac{\partial}{\partial t} \langle \chi \rangle_{R_s} - \frac{c}{B} \mathbf{b} \cdot \nabla \langle \chi \rangle_{R_s} \times \nabla F_{0s} - \frac{c}{B} \mathbf{b} \cdot \nabla \langle \chi \rangle_{R_s} \times \nabla \delta G, \end{aligned} \tag{2.6}$$

where $\langle \chi \rangle_{R_s} = \sum_k \langle \chi \rangle_{R_s,k} \exp i\mathbf{k} \cdot \mathbf{R}_s$, with $\langle \chi \rangle_{R_s,k} = J_0(a_s)(\phi_k - v_{\parallel} A_{\parallel k}/c)$ being the gyroaveraged kinetic potential, $\nabla = \partial/\partial \mathbf{R}_s$, $\mathbf{v}_{d,s} = -v_{\parallel} \hat{\mathbf{b}} \times \nabla(v_{\parallel}/\Omega_s)$ and

$$\frac{c}{B} \mathbf{b} \cdot \nabla \chi \times \nabla \delta G = \frac{c}{B} [\chi, \delta G]. \tag{2.7}$$

In a stellarator context, it is wise to introduce Boozer coordinates (Boozer 1981) $\mathbf{B} = \nabla \psi \times \nabla \theta + \iota \nabla \varphi \times \nabla \psi = G(\psi) \nabla \varphi + I(\psi) \nabla \theta + K(\psi, \theta, \varphi) \nabla \psi$. Here, the magnetic equilibrium satisfies force balance:

$$\frac{4\pi}{c} \mathbf{J} \times \mathbf{B} = \frac{1}{\sqrt{g}} \left(\iota I' + G' - \iota \frac{\partial K}{\partial \theta} - \frac{\partial K}{\partial \varphi} \right) \nabla \psi = 4\pi \nabla p_0, \tag{2.8}$$

where $1/\sqrt{g} = B^2/(G + \iota I) \simeq B^2/G$ so that

$$\iota I' + G' - \iota \frac{\partial K}{\partial \theta} - \frac{\partial K}{\partial \varphi} = \frac{4\pi G}{B^2} \frac{dp_0}{d\psi}, \tag{2.9}$$

and p_0 and \mathbf{J} are the equilibrium pressure and plasma current, respectively.

It is furthermore useful to see that

$$\begin{aligned}
 \mathbf{v}_d \cdot \nabla \delta G &= -v_{\parallel} \mathbf{b} \cdot \nabla \left(\frac{v_{\parallel}}{\Omega} \right) \times \nabla \delta G \\
 &= -\frac{v_{\parallel}}{\sqrt{g}B} \left\{ G \left[\left(\frac{\partial}{\partial \psi} \frac{v_{\parallel}}{\Omega} \right) \frac{\partial}{\partial \theta} - \left(\frac{\partial}{\partial \theta} \frac{v_{\parallel}}{\Omega} \right) \frac{\partial}{\partial \psi} \right] \right. \\
 &\quad - I \left[\left(\frac{\partial}{\partial \psi} \frac{v_{\parallel}}{\Omega} \right) \frac{\partial}{\partial \varphi} - \left(\frac{\partial}{\partial \varphi} \frac{v_{\parallel}}{\Omega} \right) \frac{\partial}{\partial \psi} \right] \\
 &\quad \left. + K \left[\left(\frac{\partial}{\partial \varphi} \frac{v_{\parallel}}{\Omega} \right) \frac{\partial}{\partial \theta} - \left(\frac{\partial}{\partial \theta} \frac{v_{\parallel}}{\Omega} \right) \frac{\partial}{\partial \varphi} \right] \right\} \delta G \\
 &\equiv -v_{\parallel} \left[\frac{v_{\parallel}}{\Omega}, \delta G \right].
 \end{aligned} \tag{2.10}$$

2.1. Nonlinear fluctuations

Equations (2.1)–(2.2) accommodate finite-β Alfvénic and drift-wave ballooning-like instabilities. Here we set up a model for nonlinear transport associated with ballooning-type modes supported by them (Chen & Zonca 2016).

As in ballooning theory (Connor *et al.* 1978; Connor, Hastie & Taylor 1979), one introduces an eikonal representation for perturbed quantities (Antonsen & Lane 1980):

$$\delta Q(\psi, \theta, \varphi, t) = \delta \hat{Q}(\psi, \alpha(\theta, \varphi), \theta, t) \exp(iS(\psi, \alpha(\theta, \varphi), t)), \tag{2.11}$$

where $\alpha = \theta - \iota\varphi$,

$$\frac{1}{S} \frac{\partial S}{\partial t} \gg \frac{1}{\delta \hat{Q}} \frac{\partial \delta \hat{Q}}{\partial t} \tag{2.12}$$

and

$$\frac{1}{S} \nabla S \gg \frac{1}{\delta \hat{Q}} \nabla \delta \hat{Q}. \tag{2.13}$$

To leading order, (2.2) becomes a nonlinear drift-wave-ballooning equation:

$$\begin{aligned}
 B \nabla_{\parallel} \frac{c}{4\pi} k_{\perp}^2 \frac{\delta \hat{A}_{\parallel, k}}{B} - i\omega \sum_s \frac{e_s^2}{T_s} \int d^3 v F_{0s} \left[1 - J_0^2(a_s) - \frac{\Omega_s^T}{\omega} J_0^{*,s}(a_s) \right] \phi_k \\
 + \frac{c}{B} \hat{\mathbf{b}} \cdot \boldsymbol{\kappa} \times \mathbf{k}_{\perp} \sum_s \int d^3 v m_s (\mu B + v_{\parallel}^2) J_0(a_s) \delta G_{sk} \\
 = i \mathcal{N}_{F, k},
 \end{aligned} \tag{2.14}$$

where $\mathbf{k} = \partial_{\psi} S \nabla \psi + \partial_{\alpha} S \nabla \alpha$, $i\omega = \partial_t S$ and \mathcal{N}_F contains all the terms that are nonlinear in the field amplitudes:

$$\begin{aligned}
 \mathcal{N}_{F, k} &= \frac{c}{B} \sum_k \mathbf{b} \cdot \mathbf{k}'_{\perp} \times \mathbf{k}_{\perp} \sum_s q_s \int d^3 v J_0(a_s) J_0(a'_s) \hat{\phi}_{k'} \delta \hat{G}_{s, k-k'} \\
 &\quad - \frac{1}{B} \sum_k \mathbf{b} \cdot \mathbf{k}'_{\perp} \times \mathbf{k}_{\perp} \sum_s q_s \int d^3 v J_0(a_s) J_0(a'_s) \hat{A}_{\parallel, k'} v_{\parallel} \delta \hat{G}_{s, k-k'}.
 \end{aligned} \tag{2.15}$$

In the linear limit, $\mathcal{N}_{F, k} \rightarrow 0$, (2.14) reduces to the equation used by Tang *et al.* in the study of linear kinetic ballooning modes in the intermediate-frequency regime (Tang *et al.*

1980; Aleynikova & Zocco 2017). Similarly, quasi-neutrality remains virtually unchanged:

$$n_i \left(\sum_s \frac{q_s n_s}{\tau_s n_i} \right) \frac{q_s \delta \hat{\phi}_k}{T_{0e}} = \sum_s \frac{q_s}{n_s} \int d^3 v J_0(a_s) \delta \hat{G}_{s,k}. \quad (2.16)$$

Equations (2.14)–(2.16) constitute a local eigenvalue problem for the fields $\delta \hat{A}_{\parallel}$ and $\delta \hat{\phi}$ where nonlinearities are retained.

Antonsen & Lane (1980) have proposed a way of attacking this type of eigenvalue problems, by applying a method introduced by Weinberg (1962), and further explored by Dewar & Glasser (1983). One considers, for a given radial location and a given field line, the local eigenvalue

$$\omega_l = \omega_l(\psi, \alpha, S_{\psi}, S_{\alpha}), \quad (2.17)$$

and determines S_{ψ} and S_{α} by solving Hamilton equations

$$\frac{\partial \psi}{\partial \ell} = \frac{\partial \omega_l}{\partial S_{\psi}}, \quad \frac{\partial \alpha}{\partial \ell} = \frac{\partial \omega_l}{\partial S_{\alpha}} \quad (2.18)$$

and

$$\frac{\partial S_{\psi}}{\partial \ell} = -\frac{\partial \omega_l}{\partial \psi}, \quad \frac{\partial S_{\alpha}}{\partial \ell} = -\frac{\partial \omega_l}{\partial \alpha}, \quad (2.19)$$

where ℓ parametrises the trajectories that make ω_l constant. This determines S_{ψ} and S_{α} . A global eigenvalue could further be determined, but we do not proceed along these lines.

Instead, we point out that, to next order in the

$$\omega_l^{-1} \delta \hat{Q}^{-1} \partial_t \delta \hat{Q} \sim k_{\perp}^{-1} \delta \hat{Q}^{-1} \nabla \delta \hat{Q} \ll 1 \quad (2.20)$$

expansion, one obtains the equation for the global structure of field amplitudes. The outcome of this calculation is a nonlinear Schrödinger-type equation that integrates the description of finite- β fluctuations with nonlinear wave–particle effects. Zonca *et al.* (2021a) have recently derived the expressions for the radial envelope of fluctuations in tokamak geometry by implementing a specific ordering between linear and nonlinear time scales. The radial envelope of fluctuations, being a higher-order perturbative quantity, will follow an equation in which corrections to the radial variation of equilibrium quantities will need to be considered. These ‘radially global’ corrections to the turbulence description supported by the gyrokinetic equation, (2.6), need to be considered extremely carefully in a stellarator, since the strong variation of the modulus of the magnetic field on the magnetic surface introduces an intermediate equilibrium scale, $\rho \ll L_B \ll R$, which induces important ‘surface-global effects’ that have no real equivalent in axisymmetry (Zocco *et al.* 2016, 2020). Thus, the generalisation to non-axisymmetry of the analysis of Zonca *et al.* (2021a), while conceptually unproblematic, needs its own dedicated work. We therefore dedicate the rest of the article to the kinetic side of the problem, that is, the evaluation of the function δG_s .

3. Kinetics

In this section we solve the kinetic equation with nonlinear wave–particle interaction in stellarators. Before proceeding, a couple of technical comments are in order.

3.1. Choice of coordinates

Let us be more precise with the choice of coordinates. We introduce

$$\begin{pmatrix} \theta \\ \varphi \end{pmatrix} \rightarrow \begin{pmatrix} \theta_c \\ \alpha_c \end{pmatrix} = \begin{pmatrix} \theta \\ \varphi - q\theta \end{pmatrix}, \tag{3.1}$$

where $q = \iota^{-1}$, so that

$$f(\psi, \theta, \varphi) \mapsto \hat{f}(\psi, \theta_c(\theta), \alpha_c(\theta, \varphi)). \tag{3.2}$$

Then

$$\frac{\partial}{\partial \theta} \Big|_{\varphi} \hat{f}(\psi, \theta_c(\theta), \alpha_c(\theta, \varphi)) = \left(\frac{\partial}{\partial \theta_c} - q \frac{\partial}{\partial \alpha_c} \right) \hat{f}(\psi, \theta_c, \alpha_c) \tag{3.3}$$

and

$$\frac{\partial}{\partial \varphi} \Big|_{\theta} \hat{f}(\psi, \theta_c(\theta), \alpha_c(\theta, \varphi)) = \frac{\partial}{\partial \alpha_c} \hat{f}(\psi, \theta_c, \alpha_c). \tag{3.4}$$

The purpose of these coordinates is to reduce the parallel streaming term operator into a first-order partial derivative in one angle only:

$$v_{\parallel} \nabla_{\parallel} = \frac{v_{\parallel} \iota}{B \sqrt{g}} \frac{\partial}{\partial \theta_c}, \tag{3.5}$$

as can easily be verified. Then the gyrokinetic equation reads

$$\begin{aligned} & \frac{\partial \delta G}{\partial t} + \frac{v_{\parallel} \iota}{B \sqrt{g}} \frac{\partial}{\partial \theta_c} \delta G + \frac{v_{\parallel} G}{B \sqrt{g}} \left(\frac{\partial \rho_{\parallel}}{\partial \theta_c} \frac{\partial}{\partial \psi} - \frac{\partial \rho_{\parallel}}{\partial \psi} \frac{\partial}{\partial \theta_c} \right) \delta G \\ & + \iota^{-1} B v_{\parallel} \left(\frac{\partial \rho_{\parallel}}{\partial \psi} \frac{\partial}{\partial \alpha_c} - \frac{\partial \rho_{\parallel}}{\partial \alpha_c} \frac{\partial}{\partial \psi} \right) \delta G = - \frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} \frac{\partial}{\partial t} \chi \\ & - \frac{c}{\sqrt{g} B^2} \left[G \left(\frac{\partial \chi}{\partial \psi} \frac{\partial}{\partial \theta_c} - \frac{\partial \chi}{\partial \theta_c} \frac{\partial}{\partial \psi} \right) - \iota \sqrt{g} B^2 \left(\frac{\partial \chi}{\partial \psi} \frac{\partial}{\partial \alpha_c} - \frac{\partial \chi}{\partial \alpha_c} \frac{\partial}{\partial \psi} \right) \right] \\ & + \frac{c}{\sqrt{g} B^2} \left[G \frac{\partial \chi}{\partial \theta_c} \frac{\partial}{\partial \psi} + \iota \sqrt{g} B^2 \frac{\partial \chi}{\partial \alpha_c} \frac{\partial}{\partial \psi} \right] F_0, \end{aligned} \tag{3.6}$$

where we introduced $\rho_{\parallel} = v_{\parallel} / \Omega$, we are dropping the symbol of gyroaverage and the species index and we are working in real space. The $\mathbf{v}_d \cdot \nabla \theta$ term has generated the

$$- \frac{v_{\parallel} G}{B \sqrt{g}} \frac{\partial \rho_{\parallel}}{\partial \psi} \frac{\partial}{\partial \theta_c} \tag{3.7}$$

contribution, and this can be discarded if compared with the streaming term (a common approximation). However, $\partial_{\psi} \rho_{\parallel} \partial_{\theta_c} \sim \partial_{\theta_c} \rho_{\parallel} \partial_{\psi}$ so, formally, $\mathbf{v}_d \cdot \nabla \theta$ can be neglected for

$$k_{\theta_c} \frac{\partial \rho_{\parallel}}{\partial \psi} \ll k_{\psi} \frac{\partial \rho_{\parallel}}{\partial \theta_c}, \tag{3.8}$$

where k_{θ_c} and k_{ψ} are the characteristic fast-varying wavelengths of fluctuations, which are assumed to be of the same order. The physical meaning of this statement is that finite

banana width effects are caused by the poloidal variation of the equilibrium quantities that enter the definition of ρ_{\parallel} . We stress here that ρ_{\parallel} is an equilibrium quantity. While its ‘fast’ variation might impact mode structure, it is not allowed to interfere with the gyrokinetic ordering $k_{\parallel}/k_{\perp} \ll 1$. The term $\partial\rho_{\parallel}/\partial\alpha_c$ is purely non-axisymmetric, and cannot be neglected if $\partial_{\theta_c}\rho_{\parallel}\partial\psi \sim \partial_{\alpha_c}\rho_{\parallel}\partial\psi$. The gyrokinetic equation that we study is therefore

$$\begin{aligned} \frac{\partial\delta G}{\partial t} + \frac{v_{\parallel}\iota}{B\sqrt{g}}\frac{\partial}{\partial\theta_c}\delta G + \frac{v_{\parallel}G}{B\sqrt{g}}\frac{\partial\rho_{\parallel}}{\partial\theta_c}\frac{\partial}{\partial\psi}\delta G - \iota^{-1}Bv_{\parallel}\frac{\partial\rho_{\parallel}}{\partial\alpha_c}\frac{\partial}{\partial\psi}\delta G &= -\frac{e}{m}\frac{\partial F_0}{\partial\mathcal{E}}\frac{\partial}{\partial t}\chi \\ -\frac{c}{\sqrt{g}B^2}\left[G\left(\frac{\partial\chi}{\partial\psi}\frac{\partial}{\partial\theta_c}-\frac{\partial\chi}{\partial\theta_c}\frac{\partial}{\partial\psi}\right) - \iota\sqrt{g}B^2\left(\frac{\partial\chi}{\partial\psi}\frac{\partial}{\partial\alpha_c}-\frac{\partial\chi}{\partial\alpha_c}\frac{\partial}{\partial\psi}\right)\right]\delta G \\ +\frac{c}{\sqrt{g}B^2}\left[G\frac{\partial\chi}{\partial\theta_c}\frac{\partial}{\partial\psi}-\iota\sqrt{g}B^2\frac{\partial\chi}{\partial\alpha_c}\frac{\partial}{\partial\psi}\right]F_0, \end{aligned} \tag{3.9}$$

where we are basically assuming that equilibrium quantities vary more strongly in α and θ than in ψ . To better understand what this approximation means, we revert to the original Boozer coordinates to obtain

$$\begin{aligned} \frac{\partial\delta G}{\partial t} + \frac{v_{\parallel}}{B\sqrt{g}}\left(\frac{\partial}{\partial\varphi} + \iota\frac{\partial}{\partial\theta}\right)\delta G + \frac{v_{\parallel}}{B\sqrt{g}}\left(G\frac{\partial\rho_{\parallel}}{\partial\theta} - I\frac{\partial\rho_{\parallel}}{\partial\varphi}\right)\frac{\partial}{\partial\psi}\delta G \\ = -\left\{\frac{e}{m}\frac{\partial F_0}{\partial\mathcal{E}}\frac{\partial}{\partial t}\chi + \frac{c}{B}[\chi, \delta G] + \frac{c}{B}[\chi, F_0]\right\}. \end{aligned} \tag{3.10}$$

The terms left that are proportional to v_{\parallel} are recognisable as

$$\frac{v_{\parallel}}{B\sqrt{g}}\left(\frac{\partial}{\partial\varphi} + \iota\frac{\partial}{\partial\theta}\right) \rightarrow v_{\parallel}\nabla_{\parallel}, \tag{3.11}$$

while the last term of the first line is

$$\frac{v_{\parallel}}{B\sqrt{g}}\left(G\frac{\partial\rho_{\parallel}}{\partial\theta} - I\frac{\partial\rho_{\parallel}}{\partial\varphi}\right)\frac{\partial}{\partial\psi} \rightarrow \mathbf{v}_d \cdot \nabla\psi\frac{\partial}{\partial\psi}. \tag{3.12}$$

In field-following coordinates, the left-hand side of (3.10) would then be

$$\frac{\partial\delta G}{\partial t} + v_{\parallel}\nabla_{\parallel}\delta G + k_{\psi}\mathbf{v}_d \cdot \nabla\psi\delta G. \tag{3.13}$$

One can choose the poloidal angle to be the field-following coordinate, thus accounting for a slow variation of perturbations along the field line, in the spirit of the underlying gyrokinetic ordering. However, this fact does not preclude fluctuations from being most important at short wavelengths, that is, when

$$k_{\psi} \sim k_{\alpha_c} \sim k_{\theta_c} \sim n/R \sim m/R \sim \rho_i^{-1} \gg 1, \tag{3.14}$$

for any set of coordinates used, where n and m are some poloidal and toroidal wavenumbers. This discussion is largely concerned with the influence of the strong variations of magnetic equilibrium quantities in gyrokinetics, and is specific to stellarators. Alternatively, one could state that (3.10) can be used when the radial derivatives of the

fluctuating fields are much larger than any other derivative, and $\mathbf{v}_d \cdot \nabla \alpha_c$ does not play a role in their destabilisation.

The inclusion in (3.10) of the $\mathbf{v}_d \cdot \nabla \theta$ and $\mathbf{v}_d \cdot \nabla \varphi$ components of the magnetic drift, which would drive the toroidal branch of the ion-temperature-gradient-driven instability, would not be problematic. This could be done by simply upgrading the ordering of the radial derivatives of ρ_{\parallel} . However, in this work, we rely on linear drives provided by wave-particle resonant excitation, thus naturally energetic particle modes, and work with (3.10) as it is. From now on, we refer to the form of (3.9) as ‘formulation A’, and to (3.10) as ‘formulation B’. The first is more useful to treat explicitly the streaming term for the nonlinear investigation of wave-particle nonlinear interaction, the second is more practical for the study of the late-time behaviour of zonal flows in a flux tube.

4. Solution of the kinetic problem

We consider formulation B (equation (3.10)), and introduce $\delta G = \exp(-H\partial_{\psi})\delta g$, to obtain the compact form

$$\begin{aligned} & \frac{\partial}{\partial t} \delta g + v_{\parallel} \hat{\mathbf{b}} \cdot \nabla \delta g \\ &= -\bar{v}_{\psi} \frac{\partial}{\partial \psi} \delta g - e^{iQ} \left\{ \frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} \frac{\partial}{\partial t} \chi + \frac{c}{B} [\chi, e^{-iQ} \delta g] + \frac{c}{B} [\chi, F_0] \right\}, \end{aligned} \quad (4.1)$$

where $Q = -iH\partial_{\psi}$ (this radial derivative acts on fluctuating quantities only) and H solves for the magnetic differential equation

$$v_{\parallel} \nabla_{\parallel} H = \frac{v_{\parallel}}{\sqrt{g}B} \left[G(\psi) \frac{\partial}{\partial \theta} \left(\frac{v_{\parallel}}{\Omega} \right) - I(\psi) \frac{\partial}{\partial \varphi} \left(\frac{v_{\parallel}}{\Omega} \right) \right] - \bar{v}_{\psi}. \quad (4.2)$$

Here $\bar{v}_{\psi} = \bar{v}_{\psi}(\mathcal{E}, \psi) \equiv \overline{\mathbf{v}_d \cdot \nabla \psi}$, since $\overline{v_{\parallel} \nabla_{\parallel} H} = 0$. Equation (4.1) was also derived by Sugama & Watanabe (2005), for a specific helical magnetic equilibrium. Our equation is completely general from the point of view of geometry, provided a solution to the magnetic differential equation is given. For this, see Mishchenko, Helander & Könies (2008). Let us introduce the ‘rippling time’, τ , which satisfies the equation

$$\left(\frac{\partial}{\partial \varphi} + \iota \frac{\partial}{\partial \theta} \right) \tau(\theta, \varphi) = \tau_b, \quad (4.3)$$

where $\tau_b^{-1} = v_{\parallel} / (B\sqrt{g})$. In these new variables, the left-hand side of (4.1) becomes

$$\frac{\partial}{\partial t} \delta g + v_{\parallel} \hat{\mathbf{b}} \cdot \nabla \delta g = \left(\frac{\partial}{\partial t} + \frac{\partial}{\partial \tau} \right) \delta g. \quad (4.4)$$

We can then introduce the ripple-bounce expansion (Sugama & Watanabe 2005):

$$\delta g(\tau) = \sum_l h_l e^{i\omega_b \tau}, \quad h_l = T_b^{-1} \oint d\tau \delta g(\tau) e^{-i\omega_b \tau}, \quad (4.5)$$

with $\omega_b = 2\pi T_b^{-1}$ and $T_b = \oint d\theta' \tau_b(\theta'_c)$, where the \oint integral is from θ_1 to θ_2 and back to θ_1 , along a trapped particle trajectory, where θ_1 and θ_2 are two consecutive bounce points whose exact definition depends on the class of trapped particles considered (Sugama &

Watanabe 2005; Mishchenko, Helander & Könies 2008). Then, the ripple-bounce and Laplace transforms are applied to find

$$\int_0^\infty dt e^{-pt} \frac{1}{T_b} \oint d\tau e^{-il\omega_b\tau} \left(\frac{\partial}{\partial t} + \frac{\partial}{\partial \tau} \right) \delta g = (p + il\omega_b) \hat{h}_l(p) - h_l(0), \tag{4.6}$$

for the first term.

All other terms in the kinetic equation are better represented by Fourier modes. We introduce

$$\Phi_{mn} = \Xi_{mn}^l \Phi_l \tag{4.7}$$

and

$$\Phi_l = \Theta_l^{mn} \Phi_{mn}, \tag{4.8}$$

where

$$\Theta_l^{mn} = \frac{1}{\oint \tau_b} \oint d\tau \exp(-i(l\tau\omega_b + n\varphi(\tau) - m\theta(\tau))) \tag{4.9}$$

and

$$\Xi_{mn}^l = \int_0^{2\pi} d\theta \int_0^{2\pi} d\varphi \exp(i(l\tau(\theta, \varphi) + n\varphi - m\theta)). \tag{4.10}$$

We also use the radial local approximation:

$$\chi(\psi, \theta, \varphi, t) = \sum_k \Phi_k(\theta, \varphi, t) e^{ik\psi} \tag{4.11}$$

and

$$\delta g(\psi, \theta, \varphi, t) = \sum_k h_k(\theta, \varphi, t) e^{ik\psi}. \tag{4.12}$$

Then, the nonlinear equation expanded in bounce harmonics is

$$\frac{\partial h_k^l}{\partial t} + il\omega_b h_k^l = -ik\bar{v}_\psi h_k^l - \frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} q_{l'} \frac{\partial \Phi_k^{l-l'}}{\partial t} + \frac{\partial F_0}{\partial \psi} H_{l,l'} \Phi_k^{l'} + \mathcal{N}_{k'}^{ll''} \Phi_{k'}^{l'} h_{k-k'}^{l''}, \tag{4.13}$$

where

$$H_{l,l'} = i \frac{c}{\sqrt{g}B^2} (nI + mG) q_{l'} \Theta_{l-l'}^{mn} \Xi_{l'}^{mn}, \tag{4.14}$$

$$\mathcal{N}_{k'}^{ll''} = \frac{c}{\sqrt{g}B^2} (\mathcal{S}_{k'}^{ll''} + i\mathcal{R}^{ll''} k k'), \tag{4.15}$$

with

$$\mathcal{S}_{k'}^{ll''} = \{G[k'm' - (k - k')m] + I[k'n' - (k - k')n]\} \Xi_{l'}^{mn} \Xi_{l''}^{mn} \Theta_{l'}^{m+m', n+n'} \tag{4.16}$$

and

$$\mathcal{R}^{ll''} \Phi_{k'}^{l'} h_{k-k'}^{l''} = (mG + nI) Q_l \Xi_{mn}^l \Theta_l^{m+m'+m'', n+n'+n''} \Xi_{l''}^{m'n'} \Xi_{l''}^{m''n''} \Phi_{k'}^{l''} h_{k-k'}^{l''} \tag{4.17}$$

and q_l are the coefficients of the bounce expansion of $\exp[iQ]$. We use the Einstein convention for repeated indices.

Dynamically, for any l , we choose

$$p \sim l\omega_b \gg k\bar{v}_\psi, \omega_{E \times B}. \tag{4.18}$$

This means that wave–particle resonance effects are kept to all orders. However, subsidiary expansions in $p/(l\omega_b)$ are allowed. In particular, one could consider the ion sound approximation, $p \gg l\omega_{b,i}$, and derive the fluid ion-temperature-gradient-driven instability (after neglecting nonlinearities). The bounce-averaged radial drift is supposed to be small since we are not interested in unoptimised machines. Our ordering is most appropriate for turbulence driven by high-frequency resonant phenomena, such as those caused by energetic particle (or electron) instabilities, or for weak turbulence close to marginality (where resonant effects are important). Strongly driven fluid regimes would require consideration of $p \sim \omega_{E \times B}$, and at present we are not in the position to conclude if our solution can efficiently describe them.

Let us now introduce a perturbative nonlinear solution. In order to explore all time scales supported by our ordering, and in particular long-time turbulent behaviour, such solution must be known to all orders. We treat perturbatively the stellarator contribution and the nonlinearity, in a fashion somewhat similar to that of Al'tshu'l & Karpman (1966) for the study of nonlinear oscillations in collisionless plasmas. As already stated, the same type of treatment has proved fruitful for the study of phase-space zonal structures in tokamaks (Zonca *et al.* 2015; Falessi & Zonca 2019). Reverting to (4.13), we immediately realise that its solution is iterative. Indeed, after introducing the Laplace transform, we find

$$(p + i l \omega_b) \hat{h}_k^{(n)l} = -i k \bar{v}_\psi \hat{h}_k^{(n-1)l} + \mathcal{N}_{k'}^{ll'l''} \int dp' \hat{\Phi}_{k'}^{l''}(p') \hat{h}_{k-k'}^{(n-1)l''}(p-p'), \tag{4.19}$$

where

$$\frac{\hat{h}_k^{(n+1)l}}{\hat{h}_k^{(n)l}} \sim \frac{\omega_{E \times B}}{l\omega_b} \sim \frac{k\bar{v}_\psi}{l\omega_b} \ll 1. \tag{4.20}$$

This solution has a very attractive diagrammatic interpretation. Given an $\hat{h}^{(n-1)}$, the next-order iterative solution is the creation of a new field $\hat{h}_k^{l,(n)} = \hat{h}_{st}^{(n)} + \hat{h}_{nl}^{(n)}$, given by two modifications of the momentum-space propagator $(p + i l \omega_b)^{-1}$, that is,

$$\hat{h}_k^{l,(n)} = \frac{1}{p + i l \omega_b} \{ \text{stellarator} + \text{nonlinearity} \}. \tag{4.21}$$

The first term is obtained via interaction of $\hat{h}_k^{(n-1)}$ at the vertex $-i k \bar{v}_\psi$ (resulting in the emission of the curly propagator emerging from $-i k \bar{v}_\psi$ in figure 1); the other is given by the annihilation of $\hat{h}^{(n-1)}$ and Φ (see figure 1). At each step, the full result is the sum of all the fields produced by the iteration. We analyse in more detail the nonlinearity of the first-order solution of figure 1. This is

$$(p + i l \omega_b) \hat{h}_{nl}^{(n)} = \mathcal{N}_{k'}^{ll'l''} \int dp' \hat{\Phi}_{k'}^{l''}(p') \hat{h}_{k-k'}^{(n-1)l''}(p-p'). \tag{4.22}$$

The right-hand side is a state characterised by two momentum-like quantities (p, k) and an index l . This is the result of the interaction at the circular vertex of figure 1 of two incoming fields: a $\hat{\Phi}^{l''}$ with ‘momentum’ (p', k') and an $\hat{h}^{(n-1)l''}$ with ‘momentum’ $(p-p', k-k')$. At the vertex momenta are conserved. The interaction is integral and its action mixes bounce harmonics via the $\mathcal{N}_{k'}^{ll'l''}$ tensor.




FIGURE 1. First-order iterative solution of (4.13).




FIGURE 2. Second-order iterative solution of (4.13).

Perturbatively, we might write, after introducing the regularised zeroth-order solution

$$\tilde{h}_k^{(0)l} = (p + i\omega_B) \hat{h}_k^{(0)l}, \tag{4.23}$$

with

$$\hat{h}_k^{(0)l} = \frac{1}{p + i\omega_B} \left\{ -\frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} q^{l'} \frac{\partial \Phi_k^{l-l'}}{\partial t} + \frac{\partial F_0}{\partial \psi} H_{l,l'} \Phi_k^{l'} \right\}, \tag{4.24}$$

the first-order solution

$$\tilde{h}_k^{(1)l} = \frac{-ik\bar{v}_\psi}{(p + i\omega_B)^2} \tilde{h}_k^{(0)l} + \frac{\mathcal{N}_{k'}^{l_1 l_2}}{p + i\omega_B} \int dp_1 \frac{\hat{\Phi}_{k_1}^{l_1}(p_1)}{p - p_1 + il_2\omega_b} \tilde{h}_{k-k_1}^{(0)l_2}(p - p_1). \tag{4.25}$$

Then, the second-order solution is

$$\begin{aligned} \tilde{h}_k^{(2)l} = & \frac{-ik\bar{v}_\psi}{(p + i\omega_B)} \left\{ \frac{-ik\bar{v}_\psi}{(p + i\omega_B)^2} \tilde{h}_k^{(0)l} + \frac{\mathcal{N}_{k'}^{l_1 l_2}}{p + i\omega_B} \int dp_1 \frac{\hat{\Phi}_{k_1}^{l_1}(p_1)}{p - p_1 + il_2\omega_b} \tilde{h}_{k-k_1}^{(0)l_2}(p - p_1) \right\} \\ & + \frac{\mathcal{N}_{k'}^{l_1 l_2}}{p + i\omega_B} \int dp_1 \hat{\Phi}_{k_1}^{l_1}(p_1) \left\{ \frac{-ik\bar{v}_\psi}{(p - p_1 + il_2\omega_B)^2} \tilde{h}_{k-k_1}^{(0)l_2}(p - p_1) \right. \\ & \left. + \frac{\mathcal{N}_{k'}^{l_2 l_1 l_3}}{p - p_1 + il_2\omega_B} \int dp_2 \frac{\hat{\Phi}_{k_1}^{l_1}(p_2)}{p - p_1 - p_2 + il_3\omega_b} \tilde{h}_{k-k_1-k_2}^{(0)l_3}(p - p_1 - p_2) \right\}, \end{aligned} \tag{4.26}$$

and so on. Here all the poles are now expressed explicitly. For the n th iteration, we generate 2^n graphs. In the following we show how to exploit the newly found nonlinear solution for the transport studies introduced in § 2.




FIGURE 3. The three types of momentum-space second-order propagators for wave-like perturbations when $-i\omega_b = p_0 = p_1$. Signs are taken into account by arrows, and quantities are conserved at the vertices. At this order, stellarator corrections and nonlinearity interact.

5. Wave-like fluctuations and transport

At this point, the nonlinear theory of energetic particles and drift-wave-like modes is assumed to be given as an input. This provides mode frequencies and amplitudes, as explained in § 2. Then, for all perturbations

$$\Phi_k(\theta, \varphi, t) = \phi^v e^{i\omega_v t}, \tag{5.1}$$

where ω_v is the local eigenvalue (2.17) of the system of equations for the fields (2.14)–(2.16) and ϕ^v solves for the nonlinear Schrödinger-type equation discussed in § 2 (notice that there is no summation for repeated indices). After applying the Laplace transform, one obtains

$$\hat{\Phi}_k^l = \Theta_{mn}^l \frac{\phi_{mn,k}^v}{p + i\omega_v} \equiv \frac{\zeta_{k,v}^l}{p + i\omega_v}, \tag{5.2}$$

and all the convolution p integrals in (4.19) can be performed by applying the Cauchy theorem and evaluating the residuals. To see specific stellarator nonlinear behaviour, we need to evaluate the second-order correction $\hat{h}_k^{(2)l}$. The result is the sum of four different combinations of momentum-space propagators which look like

$$\begin{aligned} & \frac{k^2 \bar{v}_\psi^2}{(p + i\omega_b)^3} \div \frac{k \bar{v}_\psi \mathcal{N} \zeta}{(p + i\omega_b)^2 (p - p_0)} \div \frac{k \bar{v}_\psi \mathcal{N} \zeta}{(p + i\omega_b) (p - p_0)^2} \\ & \div \frac{\mathcal{N}^2 \zeta^2}{(p + i\omega_b) (p - p_0) (p - p_1)}, \end{aligned} \tag{5.3}$$

where $p_0 = -i(l_2\omega_b + \omega_v)$ and $p_1 = -i(\omega_b l_3 + 2\omega_v)$. How such poles generate is highlighted in Appendix B.

What is left is a sum of momentum-space propagators that combine the available vertices of the theory, the stellarator-specific $k\bar{v}_\psi$ and that coming from nonlinear interaction $\mathcal{N}\zeta$. The three types of second-order propagators are shown in figure 3.

Diagrammatically, the first term of (5.3) is a stellarator self-energy loop. The second and the third terms are stellarator radiative corrections to a ‘luftballon’ diagram and the

last is a nonlinear self-energy.

$$\begin{aligned}
 h = & \text{---} + \text{---} \circlearrowleft \text{---} + \text{---} \circlearrowright \text{---} \\
 & + \text{---} \circlearrowleft \text{---} + \text{---} \circlearrowright \text{---} + \text{---} \circlearrowleft \text{---} + \dots \\
 & + \text{---} \circlearrowleft \text{---} + \text{---} \circlearrowright \text{---} + \text{---} \circlearrowleft \text{---} + \dots \\
 & + \text{---} \circlearrowleft \text{---} + \text{---} \circlearrowright \text{---} + \text{---} \circlearrowleft \text{---} + \dots
 \end{aligned}
 \tag{5.4}$$

Keeping this in mind, we are able to construct all possible solutions to all orders. We notice that the sum of all the two-vertex functions in (5.4) gives the analogue of the radiative corrections to the electron mass in quantum electrodynamics (Peskin & Schroeder 1995) (sum à-la-Dyson). The choice of specific resonant conditions allows us to evaluate which diagram is dominant in the sum, at each order.

A physics-based choice of resonances that dominate the sum is the following. Given a pole p_0 , we consider a resonant condition as $p = p_0 + \delta p$, with $\delta p \sim k\bar{v}_\psi \ll 1$, and

$$k\bar{v}_\psi \ll \mathcal{N}\zeta \ll p_0.
 \tag{5.5}$$

That is, for a slowly evolving turbulence, and an optimised machine, we can select which diagrams enter the Dyson summation in the final transport theory.

The perturbative nonlinear approach is a useful technique that helps in gaining insight into the problem. However, an equation for the complete infinite sum is preferable. The fact that the perturbative nonlinear solution can be constructed to all orders allows us to do so.

6. Phase-space zonal structure equation

Let us then focus our attention on the component of the perturbed distribution function that contributes to a ‘redefinition’ of the equilibrium quantities, in the spirit of what was originally proposed by Zonca *et al.* (2015), and then derived for axisymmetric systems by Falessi & Zonca (2019). That is, we need to identify the specific component of the fluctuating distribution function which tends to have structure at the equilibrium scale and, very importantly, is not damped or suppressed by resonantly kinetic effects. Within our treatment, this is the $l = 0$ bounce-harmonic component, as (4.13) shows, since here the wave–particle resonant term (the second term on the left-hand side) is absent, in total

analogy with the more familiar case first studied by Rosenbluth & Hinton (1998) (which we reproduce in Appendix A). Let us then consider the infinite sum

$$p \sum_{n=0}^{\infty} \hat{h}_k^{l_0, (n)} \equiv p \hat{H}_k^{l_0} (p) = p \left(\hat{h}_k^{l_0, (0)} + \hat{h}_k^{l_0, (1)} + \sum_{n \geq 2} \hat{h}_k^{l_0, (n)} \right), \tag{6.1}$$

where we notice that stellarator corrections drive the nonlinearity in the perturbed distribution function, the terms hh , only beyond second order (see the second line of the diagram in figure 2). We then iterate the perturbative n th solution twice, in order to isolate the linear drive and the $\phi h^{(0)}$ ‘nonlinearity’:

$$\begin{aligned} (p + il_0\omega_b)\hat{h}_k^{l_0, (n)} &= \int \int \frac{dp_1 dp_2}{p - p_1 + il_0\omega_b} \left\{ -ik\bar{v}_\psi \delta_{k_1, 0} \delta_{l_2, l_0} \delta(p_1) + \mathcal{N}_{k_1}^{l_0 l_1 l_2} \hat{\Phi}_{k_1}^{l_1}(p_1) \right\} \\ &\times \left\{ -i(k - k_1)\bar{v}_\psi \delta_{k_2, 0} \delta_{l_4, 0} \delta(p_2) + \mathcal{N}_{k_2}^{l_2 l_3 l_4} \hat{\Phi}_{k_2}^{l_3}(p_2) \right\} \hat{h}_{k-(k_1+k_2)}^{l_4, (n-2)} (p - p_1 - p_2). \end{aligned} \tag{6.2}$$

Plugging this result into (6.1), we obtain

$$\begin{aligned} p \hat{H}_k^0 (p) &= p \left(\hat{h}_k^{0, (0)} + \hat{h}_k^{0, (1)} \right) \\ &+ \int \int \frac{dp_1 dp_2}{p - p_1} \left\{ -ik\bar{v}_\psi \delta_{k_1, 0} \delta_{l_2, 0} \delta(p_1) + \mathcal{N}_{k_1}^{0 l_1 l_2} \hat{\Phi}_{k_1}^{l_1}(p_1) \right\} \\ &\times \left\{ -i(k - k_1)\bar{v}_\psi \delta_{k_2, 0} \delta_{l_4, 0} \delta(p_2) + \mathcal{N}_{k_2}^{l_2 l_3 l_4} \hat{\Phi}_{k_2}^{l_3}(p_2) \right\} \sum_{n \geq 2} \hat{h}_{k-(k_1+k_2)}^{l_4, (n-2)} (p - p_1 - p_2), \end{aligned} \tag{6.3}$$

where we set $l_0 = 0$, since we are interested in the zonal part of the resummed perturbed distribution function. On the other hand, $\sum_{n \geq 2} \hat{h}_{k-(k_1+k_2)}^{l_4, (n-2)} = \hat{H}_{k-(k_1+k_2)}^{l_4} (p)$; thus, for $l_0 \equiv 0$,

$$\begin{aligned} p \hat{H}_k^0 (p) &= p \left(\hat{h}_k^{0, (0)} + \hat{h}_k^{0, (1)} \right) \\ &+ \int \int \frac{dp_1 dp_2}{p - p_1} \left\{ -ik\bar{v}_\psi \delta_{k_1, 0} \delta_{l_2, 0} \delta(p_1) + \mathcal{N}_{k_1}^{0 l_1 l_2} \hat{\Phi}_{k_1}^{l_1}(p_1) \right\} \\ &\times \left\{ -i(k - k_1)\bar{v}_\psi \delta_{k_2, 0} \delta_{l_4, 0} \delta(p_2) + \mathcal{N}_{k_2}^{l_2 l_3 l_4} \hat{\Phi}_{k_2}^{l_3}(p_2) \right\} \hat{H}_{k-(k_1+k_2)}^{l_4} (p - p_1 - p_2). \end{aligned} \tag{6.4}$$

This is the equivalent of the Dyson equation within the Al’tshu’l–Karpman treatment (their equation 2.23), which is quadratic in the field amplitude, but linear for the phase-space zonal structure, that is, the collisionlessly undamped ($l_0 = 0$) bounce harmonic. Equation (6.4) determines the nonlinear distortion to the local kinetic equilibrium, and its moments describe the sought-after transport theory which embeds nonlinear wave–particle resonant interactions which dominate energetic particle physics.

7. Discussion and conclusion

In this article we have introduced a new scheme for the study of the nonlinear interaction between drift-wave-like turbulence and energetic particles in stellarator geometry. The theory is fully electromagnetic. The equations for the fluctuating fields were derived by taking moments of the gyrokinetic equation and using an ordering inspired by linear

ballooning theory. We therefore describe fluctuations as strongly varying across the equilibrium magnetic field, slowly varying along it, and characterised by an envelope that encapsulates some of their global features. The solution of the nonlinear distribution function is found by applying an iterative procedure which is based on the smallness of the bounce-averaged radial drift frequency and nonlinear $\mathbf{E} \times \mathbf{B}$ frequency. The resonant wave–particle interaction is treated with a mixed bounce harmonics/Fourier expansion. We introduced a diagrammatic interpretation of the iterative solutions, which can give guidance in the resummation of all perturbative nonlinear solutions derived at each iteration. We present a second-order perturbative nonlinear solution, for the case of wave-like fluctuating perturbations, and discuss the interplay between nonlinearity and specific stellarator features. We find that a finite bounce-averaged radial drift can couple to the nonlinearity in the fields and contribute to the evolution of turbulence. Finally we extended the concept of phase-space zonal structure to stellarator geometry, deriving the equivalent of the Dyson equation as proposed by Al'tshu'l & Karpman (1966) in the theory of nonlinear electrostatic fluctuations in plasmas. The result provides the basis for transport studies in which kinetically undamped nonlinear fluctuations contribute to the redefinition of the equilibrium distribution function. This behaviour is essential for the description of transport dynamics dominated by energetic particle physics.

Acknowledgements

A.Z. is grateful to Dr D. Rosco for illuminating discussions on the resummation of singularities that contributed to the development of the diagrammatic solutions here presented.

Editor A. Schekochihin thanks the referees for their advice in evaluating this article.

Funding

This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (grant agreement nos. 633053 and 101052200 EUROfusion). Views and opinions expressed are, however, those of the authors only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

Declaration of interests

The authors report no conflict of interest.

Appendix A. Linear theory and the Rosenbluth–Hinton response

We show how to recover the seminal result of Rosenbluth & Hinton (1998) from our treatment. We consider the auxiliary functions δg_1 and δg_2 , so that $\delta g = \delta g_1 + \delta g_2$, where δg_1 follows the nonlinear gyrokinetic equation, but it is driven by flux-surface-averaged fluctuations only:

$$\left(\frac{\partial}{\partial t} + \bar{v}_\psi \frac{\partial}{\partial \psi} + v_\parallel \mathbf{b} \cdot \nabla \right) \delta g_1 = - \left\{ \frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} e^{iQ} \frac{\partial}{\partial t} \langle \chi \rangle_\psi + \frac{c}{\sqrt{g} B^2} \langle \chi \rangle'_\psi e^{iQ} \left[G(\psi) \frac{\partial}{\partial \theta} - I(\psi) \frac{\partial}{\partial \varphi} \right] e^{-iQ} \delta g_1 \right\}. \quad (\text{A1})$$

Plugging back the result for δG_1 into the full gyrokinetic equation, one obtains the equation for δG_2 :

$$\begin{aligned} & \left(\frac{\partial}{\partial t} + \bar{v}_\psi \frac{\partial}{\partial \psi} + v_{\parallel} \mathbf{b} \cdot \nabla + \frac{c}{B} \mathbf{b} \cdot \nabla \chi \times \nabla \right) \delta g_2 \\ &= -e^{iQ} \left\{ \frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} \frac{\partial}{\partial t} (\chi - \langle \chi \rangle_\psi) + \frac{c}{B} \mathbf{b} \cdot \nabla \chi \times \nabla F_0 \right. \\ & \quad \left. + \frac{c}{B} \mathbf{b} \cdot \nabla (\chi - \langle \chi \rangle_\psi) \times \nabla e^{-iQ} \delta g_1 \right\}. \end{aligned} \tag{A2}$$

So, the non-adiabatic response is $\delta G = \exp(-iQ)(\delta g_1 + \delta g_2)$, where δg_1 and δg_2 solve (A1)–(A2).

Let us consider the local, linearised, electrostatic versions of (A1)–(A2):

$$\frac{\partial \delta g_1}{\partial t} + ik_\psi \bar{v}_\psi \delta g_1 + v_{\parallel} \nabla_{\parallel} \delta g_1 = \frac{eF_0}{T} e^{iQ} \frac{\partial}{\partial t} \langle J_0 \phi \rangle_\psi, \tag{A3}$$

$$\frac{\partial \delta g_2}{\partial t} + ik_\psi \bar{v}_\psi \delta g_2 + v_{\parallel} \nabla_{\parallel} \delta g_2 = \frac{eF_0}{T} e^{iQ} \left\{ \frac{\partial}{\partial t} (J_0 \phi - \langle J_0 \phi \rangle_\psi) - \frac{c}{B} \mathbf{b} \cdot \nabla J_0 \phi \times \nabla F_0 \right\}, \tag{A4}$$

where we are setting $\partial_\psi \equiv ik_\psi$, so $Q = k_\psi H$, and we are using a hybrid notation for the diamagnetic term, in the sense that the explicit J_0 is meant to remind the reader that the gyroaverage operation is taken at constant particle position. The electrostatic potential, ϕ , will be determined with a $k_{\perp}^2 \rho_i^2 \sim Q^2 \ll 1$ expansion of the quasi-neutrality condition:

$$-\frac{e}{T_i} \phi + \frac{1}{n} \int d^3 \mathbf{v} J_0 \delta G_i = \frac{e}{T_e} \phi + \frac{1}{n} \int d^3 \mathbf{v} \delta G_e, \tag{A5}$$

where δG_s are found for $\partial/\partial t \ll k_{\parallel} v_{thi}$ (Rosenbluth & Hinton 1998; Xiao & Catto 2006).

We anticipate an electrostatic potential which is a flux function to leading order in $k_{\perp} \rho_i \sim Q_i^2 \ll 1$, and introduce the Laplace transform of (A3)–(A4):

$$(p + ik_\psi \bar{v}_\psi) \delta \hat{g}_1 + v_{\parallel} \nabla_{\parallel} \delta \hat{g}_1 = \frac{eF_0}{T} e^{iQ} p \langle J_0 \hat{\phi} \rangle_\psi + \delta \hat{g}_1(0), \tag{A6}$$

$$(p + k_\psi \bar{v}_\psi) \delta g_2 + v_{\parallel} \nabla_{\parallel} \delta g_2 = \frac{eF_{0e}}{T_{0e}} e^{iQ} \partial \left(J_0 \hat{\phi} - \langle J_0 \hat{\phi} \rangle \right). \tag{A7}$$

We then expand for $p/(k_{\parallel} v_{thi}) \ll 1$ and obtain, to leading order (Rosenbluth & Hinton 1998; Xiao & Catto 2006),

$$v_{\parallel} \nabla_{\parallel} \delta \hat{g}_j^{(0)} = 0, \tag{A8}$$

which implies $\delta \hat{g}_j^{(0)} = \delta \hat{g}_j^{(0)}(p, \psi, \mu, \mathcal{E})$, for $j = 1, 2$. The actual form of $\delta \hat{g}_j^{(0)}$ is determined by bounce-/transit-averaging (A6)–(A7):

$$\delta \hat{g}_1^{(0)} = \frac{eF_0}{T} \frac{p}{p + ik_\psi \bar{v}_\psi} \overline{e^{iQ} \langle J_0 \hat{\phi} \rangle_\psi} + \frac{\delta \hat{g}_1(0)}{p + ik_\psi \bar{v}_\psi}, \tag{A9}$$

$$\delta \hat{g}_2^{(0)} = \frac{eF_0}{T} \frac{p}{p + ik_\psi \bar{v}_\psi} \overline{e^{iQ} \left(J_0 \hat{\phi} - \langle J_0 \hat{\phi} \rangle \right)}. \tag{A10}$$

For electrons, one simply sets $Q = 0$, $J_0 = 1$, $e \rightarrow -e$ and $\bar{v}_\psi \rightarrow -\bar{v}_\psi$.

Let us consider the $k_{\perp}^2 \rho_j^2 \sim Q^2 \ll 1$ limit in (A10), and insert the result in (A7), for passing electrons, to see that

$$\delta g_{1,e}^{(0)} = -\frac{eF_{0e}}{T_{0e}} \frac{p}{p + k_{\psi} \bar{v}_{\psi,e}} \langle \hat{\phi} \rangle_{\psi}. \tag{A11}$$

Similarly, to leading order, $\delta g_{2,e}^{(0)} = 0$. Then, in the tokamak (or helical symmetric case), for $k_{\psi} \bar{v}_{\psi} \rightarrow 0$, we obtain the modified form of quasi-neutrality (Dorland & Hammett 1993):

$$(\tau + 1) \frac{e\hat{\phi}}{T_i} = \frac{1}{n} \int d^3 \mathbf{v} \delta G_i + \frac{e}{T_e} \langle \hat{\phi} \rangle_{\psi}, \tag{A12}$$

otherwise known as (usually) proper electron response (Hammett *et al.* 1993). We now replace in (A5) the ion solution written in terms of (A9)–(A10):

$$\begin{aligned} \left(\frac{1}{\tau} + 1\right) \hat{\phi} &= \frac{1}{\tau} \int d^3 \mathbf{v} \frac{F_{0i}}{n} \frac{p}{p + ik_{\psi} \bar{v}_{\psi}} e^{-iQJ_0} \left(\overline{e^{iQJ_0} \hat{\phi}} + \frac{\delta \hat{g}_1(0)}{p} \right) \\ &+ \int d^3 \mathbf{v} \frac{F_{0e}}{n} \frac{p}{p + ik_{\psi} \bar{v}_{\psi,e}} \langle \hat{\phi} \rangle_{\psi}. \end{aligned} \tag{A13}$$

Hence, up to second order,

$$\begin{aligned} \left(\frac{1}{\tau} + 1\right) \hat{\phi} &= \frac{1}{\tau} \int d^3 \mathbf{v} \frac{F_{0i}}{n} \frac{p}{p + ik_{\psi} \bar{v}_{\psi}} \left(1 - iQ - \frac{Q^2}{2}\right) \left(1 - \frac{k_{\perp}^2 \rho_i^2 \hat{v}_{\perp}^2}{4}\right) \\ &\times \left[\left(1 - i\bar{Q} - \frac{\bar{Q}^2}{2}\right) \left(1 - \frac{\overline{k_{\perp}^2 \rho_i^2 \hat{v}_{\perp}^2}}{4}\right) \langle \hat{\phi} \rangle_{\psi} + \frac{\delta \hat{g}_1(0)}{p} \right] \\ &+ \int d^3 \mathbf{v} \frac{F_{0e}}{n} \frac{p}{p + ik_{\psi} \bar{v}_{\psi,e}} \langle \hat{\phi} \rangle_{\psi}, \end{aligned} \tag{A14}$$

where we now point out that $\delta \hat{g}_1(0) = \langle k_{\perp}^2 \rho_i^2 \rangle_{\psi} e\hat{\phi}(0)/T_i \sim k_{\perp}^2 \rho_i^2 e\phi/T_i$. Notice that the velocity-space integrands of the tokamak result of Rosenbluth and Hinton are now simply multiplied by the factor $p/(p + ik_{\psi} \bar{v}_{\psi})$, and $p/(p + ik_{\psi} \bar{v}_{\psi}) \rightarrow 1$, for $\bar{v}_{\psi} \rightarrow 0$. Taking the flux surface average, we are left with

$$\begin{aligned} &\left\{ \left[(\tau + 1) - \sum_s \frac{T_i}{T_s} \left\langle \int d^3 \mathbf{v} \frac{F_{0s}}{n} \frac{p}{p + ik_{\psi} \bar{v}_{\psi,s}} \right\rangle_{\psi} \right] + \left\langle \frac{1}{4} \rho_i^2 k_{\perp}^2 \int d^3 \mathbf{v} \frac{F_{0i}}{n} \frac{p \left(\hat{v}_{\perp}^2 + \overline{\hat{v}_{\perp}^2} \right)}{p + ik_{\psi} \bar{v}_{\psi}} \right\rangle_{\psi} \right. \\ &+ \left. \left\langle \frac{1}{2} \int d^3 \mathbf{v} \frac{F_{0i}}{n} \frac{p}{p + ik_{\psi} \bar{v}_{\psi}} \left(Q^2 + \overline{Q^2} - 2Q\bar{Q} \right) \right\rangle_{\psi} \right\} \langle \hat{\phi} \rangle_{\psi} \\ &= \langle k_{\perp}^2 \rho_i^2 \rangle_{\psi} \hat{\phi}(0) \left\langle \int d^3 \mathbf{v} \frac{F_{0i}}{n} \frac{1}{p + ik_{\psi} \bar{v}_{\psi}} \right\rangle_{\psi}. \end{aligned} \tag{A15}$$

Thus, for a vanishing bounce-averaged radial drift (tokamak case), we find the celebrated result of Rosenbluth and Hinton:

$$\left\{ \frac{1}{2} \langle k_{\perp}^2 \rho_i^2 \rangle_{\psi} + \left\langle \int d^3 \mathbf{v} \frac{F_{0i}}{n} \mathcal{Q} (\mathcal{Q} - \bar{\mathcal{Q}}) \right\rangle_{\psi} \right\} p \langle \hat{\phi} \rangle_{\psi} = \langle k_{\perp}^2 \rho_i^2 \rangle_{\psi} \hat{\phi}(0), \tag{A16}$$

where we notice the exact cancellation of the adiabatic terms on the left-hand side of (A15). In a device with a finite bounce-averaged radial drift, the particles are not allowed to fully experience a Boltzmann response (nor non-adiabatic), since their motion along the field line, necessary to achieve an isothermal state, is interrupted by the radial particle excursion. This physical effect mathematically manifests in the lack of cancellation of the first term on the left-hand side of (A15), when $\bar{v}_{\psi} \neq 0$. This is the fundamental reason why, at long wavelengths, zonal flows in stellarators are smaller than in tokamak geometry, since now

$$\langle \hat{\phi} \rangle_{\infty} = \lim_{t \rightarrow \infty} \langle \hat{\phi}(t) \rangle \approx \langle k_{\perp}^2 \rho_i^2 \rangle_{\psi} \hat{\phi}(0), \tag{A17}$$

for $\bar{v}_{\psi} \neq 0$. Equation (A15) can also be obtained by calculating the neoclassical radial particle flux and using a long-wavelength approximation of Poisson’s equation for the guiding centre density (Mishchenko *et al.* 2008):

$$\langle k_{\psi}^2 \rho_i^2 \rangle_{\psi} \phi = \sum_s e_s N_s \equiv \sum_s e_s \int d^3 \mathbf{v} \delta G_s. \tag{A18}$$

Obviously, in this case, the second term in (A16) gets replaced in the following way:

$$\left\langle \frac{1}{4} \rho_i^2 k_{\perp}^2 \int d^3 \mathbf{v} \frac{F_{0i}}{n} \frac{p (\hat{v}_{\perp}^2 + \overline{\hat{v}_{\perp}^2})}{p + ik_{\psi} \bar{v}_{\psi}} \right\rangle_{\psi} \rightarrow \frac{1}{2} \langle k_{\perp}^2 \rho_i^2 \rangle_{\psi}, \tag{A19}$$

since Larmor radius corrections, in this case, are not evaluated by solving a gyrokinetic equation that retains the $p + ik_{\psi} \bar{v}_{\psi}$ term.

Appendix B. Second-order momentum-space propagators

For wave-like perturbations, if we consider $\Phi_k(t) = \phi^v \exp(i\omega_v t)$, then

$$\hat{\Phi}_k^l \equiv \frac{\zeta_{k,v}^l}{p + i\omega_v}, \tag{B1}$$

where $\zeta_{k,v}^l$ are now constants. After inserting this in the second-order kinetic solution (4.26), recalling that

$$\tilde{h}_k^{(0)l} = (p + il\omega_B) h_k^{(0)l}, \tag{B2}$$

with

$$(p + il\omega_B) h_k^{(0)l} = -\frac{e}{m} \frac{\partial F_0}{\partial \mathcal{E}} q_l \frac{\partial \Phi_k^{l-l'}}{\partial t} + \frac{\partial F_0}{\partial \psi} H_{l,l'} \Phi_k^{l'}, \tag{B3}$$

we need to evaluate the integrals

$$\mathcal{I}_1 = \frac{1}{p + il\omega_b} \int dp_1 \frac{\zeta_{k_1,v}^{l_1}}{p_1 + i\omega_v} \frac{\tilde{h}_{k-k_1}^{(0)l_2} (p - p_1)}{p - p_1 + il_2 \omega_b}, \tag{B4}$$

$$\mathcal{I}_2 = \frac{1}{p + i\omega_b} \int dp_1 \frac{\zeta_{k_1, v}^{l_1}}{p_1 + i\omega_v} \frac{-i(k - k_1) \bar{v}_\psi}{(p - p_1 + i l_2 \omega_b)^2} \tilde{h}_{k-k_1}^{(0)l_2}(p - p_1) \quad (\text{B5})$$

and

$$\mathcal{I}_3 = \frac{1}{p + i\omega_b} \int dp_1 \frac{\zeta_{k_1, v}^{l_1}}{p_1 + i\omega_v} \frac{\mathcal{N}_{k-k_1, k_2}^{l_2, l_1, l_3}}{p - p_1 + i l_2 \omega_b} \int dp_2 \frac{\zeta_{k_2, v}^{l_1}}{p_2 + i\omega_v} \frac{\tilde{h}_{k-k_1-k_2}^{(0)l_3}(p - p_1 - p_2)}{p - p_1 - p_2 + i l_3 \omega_b}, \quad (\text{B6})$$

which produce the singularities found in the text.

REFERENCES

- ABEL, I.G. & COWLEY, S.C. 2013 Multiscale gyrokinetics for rotating tokamak plasmas. II. Reduced models for electron dynamics. *New J. Phys.* **15** (2), 023041.
- AL'TSHU'L, L.M. & KARPMAN, V.I. 1966 Theory of nonlinear oscillations in a collisionless plasmas. *Sov. Phys. JETP* **22**, 361.
- ALEJNIKOVA, K. & ZOCCO, A. 2017 Quantitative study of kinetic ballooning mode theory in simple geometry. *Phys. Plasmas* **24** (9), 092106.
- ANTONSEN, T.M. & LANE, B. 1980 Kinetic equations for low frequency instabilities in inhomogeneous plasmas. *Phys. Fluids* **23** (6), 1205–1214.
- BIANCALANI, A., BOTTINO, A., SIENA, A.D., GÜRCAN, Ö., HAYWARD-SCHNEIDER, T., JENKO, F., LAUBER, P., MISHCHENKO, A., MOREL, P., NOVIKAU, I., *et al* 2021 Gyrokinetic investigation of Alfvén instabilities in the presence of turbulence. *Plasma Phys. Control. Fusion* **63** (6), 065009.
- BOOZER, A.H. 1981 Plasma equilibrium with rational magnetic surfaces. *Phys. Fluids* **24** (11), 1999–2003.
- BOTTINO, A., FALESSI, M.V., HAYWARD-SCHNEIDER, T., BIANCALANI, A., BRIGUGLIO, S., HATZKY, R., LAUBER, P., MISHCHENKO, A., POLI, E., RETTINO, B., *et al*. 2022 Time evolution and finite element representation of Phase Space Zonal Structures in ORB5. *J. Phys. Conf. Ser.* **2397** (1), 012019.
- CARLEVARO, N., MENG, G., MONTANI, G., ZONCA, F., HAYWARD-SCHNEIDER, T., LAUBER, P., LU, Z. & WANG, X. 2022 One dimensional reduced model for ITER relevant energetic particle transport. *Plasma Phys. Control. Fusion* **64** (3), 035010.
- CHEN, L. & ZONCA, F. 2016 Physics of Alfvén waves and energetic particles in burning plasmas. *Rev. Mod. Phys.* **88**, 015008.
- CHEN, L. & ZONCA, F. 2019 Self-consistent kinetic theory with nonlinear wave-particle resonances. *Plasma Sci. Technol.* **21** (12), 125101.
- CONNOR, J.W., HASTIE, R.J. & TAYLOR, J.B. 1978 Shear, periodicity, and plasma ballooning modes. *Phys. Rev. Lett.* **40** (6), 396.
- CONNOR, J.W., HASTIE, R.J. & TAYLOR, J.B. 1979 High mode number stability of an axisymmetric toroidal plasma. *Proc. R. Soc. Lond A* **365** (1720), 1–17.
- DEWAR, R.L. & GLASSER, A.H. 1983 Ballooning mode spectrum in general toroidal systems. *Phys. Fluids* **26** (10), 3038–3052.
- DI SIENA, A., GÖRLER, T., POLI, E., BAÑÓN NAVARRO, A., BIANCALANI, A., BILATO, R., BONANOMI, N., NOVIKAU, I., VANNINI, F., JENKO, F., *et al* 2021 Nonlinear electromagnetic interplay between fast ions and ion-temperature-gradient plasma turbulence. *J. Plasma Phys.* **87** (2), 555870201.
- DIAMOND, P.H., ITOH, S.-I., ITOH, K. & HAHM, T.S. 2005 Zonal flows in plasma—a review. *Plasma Phys. Control. Fusion* **47** (5), R35.
- DORLAND, W. & HAMMETT, G.W. 1993 Gyrofluid turbulence models with kinetic effects. *Phys. Fluids B* **5** (3), 812–835.
- FALESSI, M.V. & ZONCA, F. 2019 Transport theory of phase space zonal structures. *Phys. Plasmas* **26** (2), 022305.

- FRIEMAN, E.A. & CHEN, L. 1982 Nonlinear gyrokinetic equations for low-frequency electromagnetic waves in general plasma equilibria. *Phys. Fluids* **25** (3), 502.
- FRIEMAN, E.A., REWOLDT, G., TANG, W.M. & GLASSER, A.H. 1980 General theory of kinetic ballooning modes. *Phys. Fluids* **23** (9), 1750–1769.
- HAMMETT, G.W., BEER, M.A., DORLAND, W., COWLEY, S.C. & SMITH, S.A. 1993 Developments in the gyrofluid approach to tokamak turbulence simulations. *Plasma Phys. Control. Fusion* **35** (8), 973.
- HASEGAWA, A. & SATO, T. 1981 *Space Plasma Physics Stationary Processes*, vol. 1. Springer.
- HAYWARD-SCHNEIDER, T., LAUBER, P., BOTTINO, A. & MISHCHENKO, A. 2022 Multi-scale analysis of global electromagnetic instabilities in iter pre-fusion-power operation plasmas. *Nucl. Fusion* **62** (11), 112007.
- LU, Z., MENG, G., HATZKY, R., HOELZL, M. & LAUBER, P. 2023 Full f and gyrokinetic particle simulations of alfvén waves and energetic particle physics. *Plasma Phys. Control. Fusion* **65** (3), 034004.
- MISHCHENKO, A., HELANDER, P. & KÖNIES, A. 2008 Collisionless dynamics of zonal flows in stellarator geometry. *Phys. Plasmas* **15** (7), 072309.
- MISHCHENKO, A. & ZOCCO, A. 2012 Global gyrokinetic particle-in-cell simulations of internal kink instabilities. *Phys. Plasmas* **19** (12), 122104.
- PESKIN, M.E. & SCHROEDER, D.V. 1995 *An Introduction to Quantum Field Theory*. Perseus Books.
- QIN, H., TANG, W.M. & REWOLDT, G. 1998 Gyrokinetic theory for arbitrary wavelength electromagnetic modes in tokamaks. *Phys. Plasmas* **5** (4), 1035–1049.
- ROGERS, B.N., DORLAND, W. & KOTSCHENREUTHER, M. 2000 Generation and stability of zonal flows in ion-temperature-gradient mode turbulence. *Phys. Rev. Lett.* **85**, 5336–5339.
- ROMANELLI, F. & ZONCA, F. 1993 The radial structure of the ion-temperature-gradient-driven mode. *Phys. Fluids B* **5** (11), 4081–4089.
- ROSENBLUTH, M.N. & HINTON, F.L. 1998 Poloidal flow driven by ion-temperature-gradient turbulence in tokamaks. *Phys. Rev. Lett.* **80**, 724–727.
- SUGAMA, H. & WATANABE, T.-H. 2005 Dynamics of zonal flows in helical systems. *Phys. Rev. Lett.* **94**, 115001.
- TANG, W.M., CONNOR, J.W. & HASTIE, R.J. 1980 Kinetic-ballooning-mode theory in general geometry. *Nucl. Fusion* **20** (11), 1439.
- WEINBERG, S. 1962 Eikonal method in magnetohydrodynamics. *Phys. Rev.* **126** (6), 1899–1909.
- XIAO, Y. & CATTO, P.J. 2006 Short wavelength effects on the collisionless neoclassical polarization and residual zonal flow level. *Phys. Plasmas* **13** (10), 102311.
- ZOCCO, A., ALEYNIKOVA, K. & XANTHOPOULOS, P. 2018a Strongly driven surface-global kinetic ballooning modes in general toroidal geometry. *J. Plasma Phys.* **84** (3), 745840303.
- ZOCCO, A., HELANDER, P. & CONNOR, J.W. 2015 Magnetic compressibility and ion-temperature-gradient-driven microinstabilities in magnetically confined plasmas. *Plasma Phys. Control. Fusion* **57** (8), 085003.
- ZOCCO, A., PLUNK, G.G. & XANTHOPOULOS, P. 2020 Geometric stabilization of the electrostatic ion-temperature-gradient driven instability. II. Non-axisymmetric systems. *Phys. Plasmas* **27** (2), 022507.
- ZOCCO, A., PLUNK, G.G., XANTHOPOULOS, P. & HELANDER, P. 2016 Geometric stabilization of the electrostatic ion-temperature-gradient driven instability. I. Nearly axisymmetric systems. *Phys. Plasmas* **23** (8), 082516.
- ZOCCO, A., XANTHOPOULOS, P., DOERK, H., CONNOR, J.W. & HELANDER, P. 2018b Threshold for the destabilisation of the ion-temperature-gradient mode in magnetically confined toroidal plasmas. *J. Plasma Phys.* **84** (1), 715840101.
- ZONCA, F. & CHEN, L. 1992 Resonant damping of toroidicity-induced shear-alfvén eigenmodes in tokamaks. *Phys. Rev. Lett.* **68**, 592–595.
- ZONCA, F. & CHEN, L. 1993 Theory of continuum damping of toroidal alfvén eigenmodes in finite- β tokamaks. *Phys. Fluids B* **5** (10), 3668–3690.
- ZONCA, F. & CHEN, L. 2014 Theory on excitations of drift Alfvén waves by energetic particles. I. Variational formulation. *Phys. Plasmas* **21** (7), 072120.

- ZONCA, F., CHEN, L., BRIGUGLIO, S., FOGACCIA, G., VLAD, G. & WANG, X. 2015 Nonlinear dynamics of phase space zonal structures and energetic particle physics in fusion plasmas. *New J. Phys.* **17** (1), 013052.
- ZONCA, F., CHEN, L., FALESSI, M.V. & QIU, Z. 2021a Nonlinear radial envelope evolution equations and energetic particle transport in tokamak plasmas. *J. Phys.: Conf. Ser.* **1785** (1), 012005.
- ZONCA, F., TAO, X. & CHEN, L. 2021b Nonlinear dynamics and phase space transport by chorus emission. *Rev. Mod. Plasma Phys.* **5**, 8.