

This is a “preproof” accepted article for *Invasive Plant Science and Management*. This version may be subject to change in the production process, *and does not include access to supplementary material*. DOI: 10.1017/inp.2025.11

Two decades of progress targeting boneseed (*Chrysanthemoides monilifera* subsp. *monilifera*): a global review to inform eradication in Western Australia

Kathryn L. Batchelor¹, John K. Scott² and Bruce L. Webber^{3,4}

¹Experimental Scientist (ORCID 0000-0001-6852-5784), CSIRO Health and Biosecurity, Floreat, Western Australia 6014, Australia; ²Honorary Research Fellow (ORCID 0000-0002-8303-9706), CSIRO Health and Biosecurity, Floreat, Western Australia 6014, Australia; ³Principal Research Scientist (ORCID 0000-0001-5934-6557), CSIRO Health and Biosecurity, Floreat, Western Australia 6014, Australia; ⁴Adjunct Professor, School of Biological Sciences, University of Western Australia, Crawley, Western Australia 6009, Australia.

Author for correspondence: Bruce L. Webber, CSIRO Health and Biosecurity, 147 Underwood Ave, Floreat, Western Australia 6014, Australia. (Email: bruce.webber@csiro.au)

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Abstract

Boneseed (*Chrysanthemoides monilifera* subsp. *monilifera*; syn *Osteospermum moniliferum* subsp. *moniliferum*), is a perennial shrub native to the south-western and southern coasts of South Africa. It was introduced to Australia in about 1852 and now represents a significant threat to natural ecosystems. Despite being listed as a Weed of National Significance, momentum on improving boneseed management has dissipated at a national level, beginning in 2008 (when a national research initiative finished) and increasingly after 2013 (when funding for national coordination ceased). A recent synthesis of past management for boneseed and recommendations for guiding future priorities has rekindled interest in Western Australia. To complement this synthesis and to identify improvements for program efficiency or effectiveness, we reviewed research and management findings on this weed with a focus on the past two decades. We collated information across the ecology and biology of boneseed, and the near relative, bitou bush, as well as useful insight from boneseed management programs applied elsewhere. As part of this review we assessed the classical biological control work that has been done on boneseed, focusing on likely explanations for why, despite nine agents and a naturalised fungus, biological control is not an effective management tool. Our synthesis suggests that for the limited populations with low abundance plants in Western Australia, eradication from the state remains a realistic target. This objective, however, needs to build on the collated baseline of past management efforts and deploy a carefully planned management program over the coming two decades. Systematic surveillance using the latest techniques, combined with manual or herbicide removal and controlled burns where possible, remain the most suitable methods to deploy. The long-lived soil seedbank requires detailed monitoring following initial plant removals and long-term funding to ensure the sustained effort required to deliver the goal of eradication for boneseed in Western Australia.

KEYWORDS: Delimitation, environmental weed, eradication, historical context, invasion, seed dispersal, soil seedbank, weed management

Management Implications

Boneseed (*Chrysanthemoides monilifera* subsp. *monilifera*; syn *Osteospermum moniliferum* subsp. *moniliferum*) is a shrub native to the south western and southern coasts of South Africa. In Australia, boneseed was introduced in the 1850's and has spread extensively in the south-eastern states of Victoria, South Australia and Tasmania. The introduction of boneseed to Western Australia is thought to have happened almost a century later and spread remains far more restricted. After being classified as a Weed of National Significance in Australia in 1999, detailed management plans were developed for boneseed, drawing on a synthesis of available data at the time. This coordinated effort resulted in two detailed literature reviews being produced in 2006 and 2008. Since this time, research and management on boneseed has continued to generate new insight. In South Australia, containment lines are being applied and more effective biological control may improve management outcomes. In Western Australia, a recent synthesis of past management has revealed that eradication remains a realistic target. All of these programs would benefit from an updated synthesis of relevant knowledge. Our review has brought together new information from the past two decades of research on the ecology, biology and management of boneseed, with the specific goal of improving the chance of eradicating boneseed in Western Australia. There remains a lack of detailed understanding of key ecological issues relating to management, including whether or not it is an obligate outcrossing taxon and the impacts of seed dispersal vectors in its non-native range. In contrast, understanding for germination requirements, allelopathic effects, and the effectiveness of management options including fire and herbicides, have been progressed. Regarding classical biological control, our review has revealed plausible explanations as to why the existing agents have not resulted in broadly effective management. Management of boneseed in south-eastern Australia and New Zealand's south island remains focused on containment where possible, some localised extirpations and minimising impact. Invasions elsewhere in the world remain largely unmanaged, but at the same time have no documented evidence of rapid invasion. Our synthesis supports retaining eradication as the management goal for boneseed in Western Australia via a mixed approach of manual removal, herbicide and controlled burns where logistics permit.

Introduction

Boneseed, *Chrysanthemoides monilifera* subsp. *monilifera* (L.) T. Norl. (Asteraceae; syn *Osteospermum moniliferum* subsp. *moniliferum*), is a perennial shrub native to the south-western and southern coasts of South Africa (Weiss et al. 2008). In Australia, boneseed was first recorded as an introduced plant in Sydney in 1852 from MacLeay's garden, Melbourne in 1858 (and subsequently grown in Melbourne suburbs as a garden plant), Adelaide in 1892 from the West Terrace Cemetery, and Ulverstone, Tasmania in 1931 (Weiss et al. 1998). At the time this shrub was mostly cultivated as an ornamental garden plant, but there may have been deliberate naturalisation attempts in western Victoria in the You Yang Ranges and to stabilize coastal sand dunes between Nelson and Portland, Victoria (Weiss et al. 1998).

Currently, boneseed is widely distributed in southern Australia (Brougham et al. 2006). The distribution of boneseed in south-eastern Australia covers an area from the Eyre Peninsula in South Australia to the Victorian - New South Wales border, with a distribution that extends from the coast to a significant way inland in some areas. Extensive infestations occur in South Australia's Mt Lofty Ranges and Murraylands, and Victoria's Mornington Peninsula, Bellarine Peninsula and the You Yang Ranges. In New South Wales (NSW), scattered infestations are present along the coast from Newcastle on the Central Coast to Moruya in the south. The majority of infestations are in the Sydney region, extending west into the Blue Mountains (Brougham et al. 2006). In Tasmania, boneseed is concentrated around the fringes of inhabited areas and is mainly restricted to coastal and estuarine areas (Brougham et al. 2006). The disjunct distribution in Western Australia (WA) is more restricted in its range and abundance, found across multiple populations between Albany and Perth (Batchelor et al. 2023a, 2024).

Boneseed prefers winter rainfall regions, where it is found in a wide range of vegetation communities including coastal dunes, estuarine areas, heath, mallee, woodland, and dry and wet sclerophyll forest (Brougham et al. 2006). Boneseed occurs on a range of soil types but does not tolerate water-logged soils (Muyt 2001). Seedlings emerge during winter, reaching reproductive maturity in the second year of growth. Flowering occurs from August to October with fruiting following during September to November. Flowers are protandrous with seeds usually produced by allogamy (Weiss et al. 2008). There is one seed per fruit, up to six seeds per inflorescence and up to 50,000 seeds per plant, every year (Weiss et al. 1998). In Australia, birds are the primary dispersal vectors, although most seeds fall beneath the plant and either enter the seed bank or are consumed by rodents or ants. Seed

longevity is highly variable and likely depends on local context, ranging from at least three years (Weiss 1984), to greater than eight years (French et al. 2024), eight and a half years (Briden and McAlpine 2012) and nine years (Luke McMillan pers comm). Mature plants are estimated to live 10 to 20 years (Muyt 2001).

Boneseed's impact as a weed is most severe in natural ecosystems, and its presence is associated with the decline of flora and fauna in south-eastern Australia. Grassy woodland, valley grassy forest and lowland forest vegetation communities in Victoria are vulnerable to boneseed invasion, where dense infestations eliminated most native ground flora and prevent virtually all overstory regeneration (Muyt 2001). Dense, continuous canopy cover for boneseed has been recorded in areas of the You Yang Ranges (Adair and Holtkamp 1999), directly threatening the endangered orchid *Pterostylis truncata* Fitzg. (Adair et al. 2012; Bramwells 2003). The local extirpation of some 40 indigenous plant species within these Ranges has been largely attributed to the local boneseed infestation (Blood 1987; Thomas et al. 2005). Moreover, removal of boneseed in a defined 109 ha area has coincided with an overall increase in koalas (*Phascolarctos cinereus*) and an expansion of their use of this habitat (Duffy 2020).

At the national level, boneseed (as well as bitou bush; *Chrysanthemoides monilifera* subsp. *rotunda* (DC.) J.C.Manning & Goldblatt; syn *Osteospermum moniliferum* subsp. *rotundatum* (DC.)) was classified as a Weed of National Significance (WoNS) in 2000, due to the significant impacts it has on natural environments (ARMCANZ 2000). A national strategy was drafted for its management attempting to “arrest the spread and minimise the impact of bitou bush and boneseed in natural ecosystems” (ARMCANZ 2000). The National Bitou and Boneseed Management Group was formed to implement the goals in the strategy. The Strategy was revised in 2012 by the Australian Weeds Committee to progress the legacy of achievements by stakeholders under the previous strategy.

Unlike the boneseed invasion in the south-east of Australia, populations in Western Australia are often reasonably discrete and appear slow to spread. Some populations are also in areas where local conditions are well suited to detailed monitoring and large-scale management interventions. Boneseed in Western Australia became a declared plant in 1979 under the Agriculture and Related Resources Protection Act, 1976-1978 (Government Gazette of Western Australia, No. 4, 1980). In 2006, boneseed in Western Australia was reclassified as a plant not to be introduced to the state with existing populations targeted for eradication (initially ‘P1/P2’, now ‘Category C2’). The current category prevents any sale,

trade, or movement. Surveillance and management since that time has varied in effectiveness, particularly since 2013 when funding from the national WoNS program ceased (Batchelor et al. 2022).

There is a strong likelihood that a more coordinated, consistent, and systematic approach to surveillance could deliver greatly improved outcomes for boneseed management in WA, as well as elsewhere where the plant is an introduced invader. To prepare for such management, two deliverables are required. First, an aggregated synthesis of past management and control efforts is required, detailing present and past population demographics for the weed. Until recently, this has not existed for Western Australia because boneseed management has been done by multiple agencies without enduring overall coordination and has not been consistent in either space or time. A synthesis of this data as well as a full assessment of boneseed populations across the state has recently been completed (Batchelor et al. 2024). This insight will transform the ability of land managers in Western Australia to know *where* and *when* to act.

Second, an updated review of the literature covering boneseed management is essential to leverage existing understanding and identify knowledge gaps. This knowledge will help to inform *how* to act, particularly in regard to improving the efficiency and effectiveness of management. The reviews of Brougham et al. (2006) and Weiss et al. (2008) provided a comprehensive guide to the ecology and biology of boneseed and its management options in Australia as was best known at that time. The Brougham et al. (2006) review comprised six sections, providing detailed insight on ecology, biology, control and post-control restoration, and monitoring. It also featured case studies of boneseed control from South Australia, Tasmania and Victoria. While there is strong overlap in content between the two publications, the Weiss et al. (2008) paper provides greater detail on the taxonomy, distribution (actual and potential), plant growth and development, dispersal and population dynamics and legislative status.

Since these reviews, boneseed has remained a focus for much research in Australia and elsewhere. While most of this work has focused on boneseed as an introduced weed, further context from the native range can also provide relevant guidance for refining management. Here we review the past two decades of research on the ecology, biology and management of boneseed to update the earlier work of Brougham et al. (2006) and Weiss et al. (2008) with the specific goal of guiding more efficient and effective management of boneseed in Western Australia specifically, and elsewhere more generally.

Methods

We took two approaches to assembling content for this review. First, we consulted an extensive literature collection, including “grey” literature, that we have acquired through active searching for any publications on *Chrysanthemoides* over the past 20 years (i.e., from when we started working on this genus; Scott 1996). Second, further literature on the biology, ecology and management of boneseed since 2008 was systematically identified through keyword searches of Web of Science (webofscience.com), Google Scholar (scholar.google.com) and the standard Google search engine (google.com) using the following search strings: boneseed, *Chrysanthemoides*, *Chrysanthemoides monilifera*, *Osteospermum moniliferum*. The latter name reflects a more recent taxonomic interpretation of *Chrysanthemoides* (synonymised in *Osteospermum*; Sadler et al. 2022) that has been accepted by the Plants of the World Online database and iNaturalist (POWO 2024) but is yet to be accepted by the Australian Plant Name Index (APNI) published by the Council of Heads of Australasian Herbaria (<https://chah.gov.au/>). For the purposes of this review, we have retained the use of *Chrysanthemoides monilifera* subsp. *monilifera*, based on the currently accepted name in the Australian Plant Census database (<https://www.anbg.gov.au/cpbr/program/hc/hc-APC.html>). Where relevant and to provide specific context, we referenced literature from before 2008. However, we have not sought to make this a thorough overview of all previous work on boneseed, given the earlier reviews covering this literature in detail (Brougham et al. 2006; Weiss et al. 2008). While this review focuses on boneseed, we also include literature on the closely related bitou bush where it aids in the understanding of boneseed management.

Results and Discussion

History of boneseed in Western Australia

The first confirmed record of boneseed in Western Australia comes from a specimen collected by Brother Kissane on the 27th August 1948 and lodged at the Western Australian Herbarium (PERTH 416444; AVH data). It was nearly three decades, however, before this specimen was formally determined as boneseed (Gray 1976). The locality of the collection was recorded simply as “Armadale”, but on the same day the collector also lodged a record for *Leucopogon capitellatus* DC. (PERTH 3004341; AVH data) with a location of “Cooliabberra Spring”, which is either a winter flowing creek within Bungendore Regional Park off the Albany Hwy in Armadale or a private landholding adjacent to the park. It is very likely that given the temporal proximity, the two records were from a similar area, which would place the first confirmed boneseed record for Western Australia near a well-known infestation that follows Neerigen Brook alongside the Albany Hwy.

The number of residential properties found to have boneseed in the Perth hills and Narrogin (Batchelor et al. 2024) could suggest that at some point boneseed was available from plant nurseries. However, the plant has never been listed in any historical Western Australian nursery or seed catalogue (J. Viska, Australian Garden History Society, pers. comm.). Cherry (2010b) noted that boneseed at Dardadine may have been introduced with teachers from South Australia as this was the site of an old school. Given that the school closed in 1935, this would make that location WA’s earliest plantings if true. We could not identify any other sources to further clarify the origins of boneseed in Western Australia and, therefore, the origin remains unknown. Molecular studies would be a logical way to provide greater clarity on introduction pathways and timing, which in turn can inform improved management (Emmett et al. 2023).

Ecology and biology

Genetics

Multiple genetic analyses that have included the genus *Chrysanthemoides* have confirmed that boneseed is a well circumscribed and separate taxon to bitou bush (Barker et al. 2009; Barker et al. 2015; Barker et al. 2005; Byrne et al. 2022; Emmett et al. 2023). The distribution of boneseed in its region of origin is relatively restricted and uncontroversial,

occurring in the Western Cape Province of South Africa. There is evidence for boneseed hybridisation with bitou bush in Australia where the two subspecies distributions overlap in north coastal Victoria (Adair and Butler 2010; also herbarium collections MEL 1553294A, CANB 377051.1). There is no record of hybridisation in Western Australia. In South Africa, these two *Chrysanthemoides monilifera* subspecies (i.e., bitou bush and boneseed) are allopatric (Norlindh 1943).

Bitou bush is an obligate outcrossing taxon (Gross et al. 2017; Scott et al. 2019b). This means that isolated plants do not produce seeds until another individual germinates nearby and flowers (i.e., subject to Allee effects due to pollination limitations). It is not known if boneseed is likewise obligately outcrossing, despite the mention of allogamy in Weiss et al. (2008). This trait needs to be determined as it has significant implications for the management of boneseed introductions.

Seed dispersal by birds and mammals

In South Africa, Knight (1988) recorded that 15 bird species visited *Chrysanthemoides monilifera* plants to feed on fruits, with the most frequent visitors being the sombre bulbul (*Andropodus importunus*), Cape bulbul (*Pycnonotus capensis*) and the Rameron pigeon (*Columba arquatrix*). Knight (1988) did not identify the subspecies of *Chrysanthemoides monilifera*, but we investigated the flora of his study location (Fernkloof Nature Reserve, Western Cape Province, 34°23'14" S, 19°16'38" E) and used its complete flora list with photos (<https://www.fernkloof.org.za/index.php/all-plants/plant-families/item/osteospermum-moniliferum-subsp-moniliferum>), confirming that the reserve only has boneseed.

Gosper (2003) reported on a detailed study of frugivores on bitou bush from coastal NSW. Of the 22 bird species feeding on the fruit the most frequent species were the pied currawong (*Strepera graculina*), Lewin's honeyeater (*Meliphaga lewinii*), silvereye (*Zosterops lateralis*), red-whiskered bulbul (*Pycnonotus jocosus*), yellow-faced honeyeater (*Lichenostomus chrysops*) and olive-backed oriole (*Oriolus sagittatus*; Gosper 2003). Emu (*Dromaius novaehollandiae*) have been reported to consume fruit and may be the vectors responsible for long distance dispersal (Brougham et al. 2006). While the viability of emu-egested seed has never been confirmed through formal experiments, dense clusters of boneseed seedlings have been observed germinating from emu scats (Batchelor et al. 2024).

An understanding of seed dispersal potential is critical to management of a bird dispersed species such as boneseed. This relationship was studied by Mokotjomela et al. (2013a; 2013b; 2013c) as part of a broader study examining seed dispersal by frugivores in South Africa. The target plants in the study were two native (boneseed and *Olea europaea* subsp. *africana*) and two non-native plant species (*Lantana camara* and *Solanum mauritianum*) and the birds were the small species Cape white eye (*Zosterops capensis*), the medium-size Cape bulbul (*Pycnonotus capensis*) and the large speckled mousebird (*Colius striatus*). Flight distances corresponding with predicted seed gut retention times for boneseed were 9.4 km, 17.8 km, and 21.2 km for *Z. capensis*, *P. capensis* and *C. striatus*, respectively. These maximum potential distances for seed dispersal, based on theoretical bird-ring recapture frequency and gut retention times, were much greater than that previously reported, which was in the order of 1 km (Mokotjomela et al. 2013c). While this study may suggest that very long-distance dispersal events are theoretically possible for boneseed, their likelihood of occurrence in field-relevant conditions is another matter entirely.

Seed predation by birds and mammals

Boneseed in its native habitat, South Africa, is harvested from plants and from the ground by Chacma baboons (*Papio ursinus*), but seeds do not survive ingestion (Knight 1988). Most seed is not dispersed but falls under the parent plant onto the ground (Knight 1988; Scott 1996) where seeds are subject to predation by rodents (Scott 1996).

In south-eastern Australia, Gosper (2003) recorded bitou bush seed predation by crimson rosella (*Platycercus elegans*), eastern rosella (*P. eximius*) and the house sparrow (*Passer domesticus*). In contrast, boneseed is not exposed to seed predation by parrots because the two types of organisms are geographically widely separated in Africa. Boneseed is restricted to the south west Western Cape Province whereas parrots are found in the northern parts of southern Africa (eight species in the family Psittacidae; Newman 1983). It is likely that rodent predation is the evolutionary driver producing a “bone” seed (i.e., round and hard).

Seed germination

Recent seed germination research on boneseed has focused on germination microclimates and chemical amendments to stimulate germination. Boneseed seeds germinated faster over a

range of temperatures with the application of karrikins to harvested seeds, relative to control treatments (Reynolds et al. 2014). Their study also determined that seed imbibition was rapid (within 48 hours) and that dormancy was physiological. Interestingly, germination was not inhibited by the hard, woody endocarp and dormancy occurred in the winter months. Germination occurred over relatively low temperatures (10 – 20°C), characteristic of winter in south-west Western Australia, and ceased at 35°C (Reynolds et al. 2014). In contrast, Batchelor et al. (2023a) found that germination of boneseed seeds was not enhanced by smoke water (containing karrikins), while gibberellic acid accelerated germination but did not increase overall germination percentage. Seeds appeared to have no after-ripening phase and seed desiccation did not increase the likelihood of endocarps fracturing (i.e. to initiate imbibition). Seeds were found to be vulnerable to direct flame exposure, even for short periods of 10 seconds, suggesting controlled burns may well lead to high mortality for seeds not protected deeper in the soil seedbank (Batchelor et al. 2023a).

Survival of seeds of boneseed and bitou bush were compared in controlled aging experiments performed at 45°C, and 60% relative humidity (RH; Schoeman et al. 2010). The number of days to lose 50% viability in boneseed and bitou bush was 47 days and 16 days, respectively. The authors predicted that boneseed may have long-lived seed bank and bitou bush a more transient (< one year) seed bank (Schoeman et al. 2010). French et al. (2024) buried seeds of boneseed and bitou bush at two locations, two depths and sampled regularly for eight years. Seed viability showed a rapid decline with time, although in excess of 10% of seeds were still alive at eight years at some sites. It is possible that edaphic factors, in particular duration of soil moisture could explain the variation. Detailed field sampling of bitou bush over more than a decade (Scott et al. 2019b) found that the soil seedbank of a non-native population in Kwinana, Western Australia appeared to have a seed longevity of approximately seven years. As the time since last known seed additions to the seedbank increased it become increasingly difficult to measure seed bank viability as it consisted of very few seeds (Scott et al. 2019b).

Biotic resistance

Grazing using goats, sheep or cattle is good at suppressing boneseed, but plants soon recovered once the grazers were removed and there is the risk of seed dispersal via the grazing animals (Briden 2008). The Tasmanian distribution of boneseed is concentrated in cities and towns and is generally absent from the intervening rural areas. Scurr et al. (2008) hypothesised, and

demonstrated by exclusion experiments, that grazing by wild macropods (especially Tasmanian pademelons, *Thylogale billardierii*) and domesticated herbivores may be sufficient to prevent the spread of boneseed. This hypothesis could also explain the disparate distribution in Western Australia and needs to be further assessed.

Further supporting the role of native herbivores is the study of overabundant wallabies (mostly swamp wallaby; *Wallabia bicolor*; plus other grazing marsupials) in the Booderee National Park in southern New South Wales (Dexter et al. 2013). Grazing by wallabies inhibiting the recruitment of bitou bush following fire-induced recruitment events in 18 unfenced (browsed) plots whereas healthy recruitment occurred in 16 fenced (unbrowsed) plots (Dexter et al. 2013).

Allelopathy

Understanding allelopathy is important as it may have negative impacts on restoration efforts following boneseed removal. Field experience and anecdotal evidence indicate the likelihood that *Chrysanthemoides* species have allelopathic effects on other vegetation (Weiss et al. 2008). Vranjic et al. (2000) found that shoot and root biomass of coast wattle (*Acacia sophorae*) were significantly lower for seedlings grown in bitou bush soil than for those grown in Acacia soil. Our surveys of bitou bush indicate that seeds were most dense under the parent bush, but without germination unless the parent plant was dead for a few years (Scott et al. 2019b). Ens (2007) investigated the allelopathic effects of bitou bush in a PhD thesis at Wollongong University, and subsequently published that allelopathy in bitou bush was a key mechanism driving the recruitment limitation of indigenous flora (Ens and French 2008; Ens et al. 2008; Ens et al. 2010; Ens et al. 2009a; Ens et al. 2009b).

A second PhD on allelopathy in *Chrysanthemoides* generated a series of papers on boneseed at Victoria University (Al Harun *et al.* 2014, 2015a, 2015b, 2015c). An initial experiment of aqueous solutions of ground-up boneseed plant parts (leaves, stem, roots) had no impact on germination of lettuce (*Lactuca sativa* Asteraceae) and little impact on germination of black wattle (*Acacia mearnsii* Fabaceae), but inhibited growth and germination of *Isotoma axillaris* (Campanulaceae). The latter two species are found in the Australian environment where boneseed grows whereas *L. sativa* is the usual bioassay for allelopathy studies (Al Harun et al. 2014).

A test of volatiles coming from boneseed stems, leaves and roots had no impact on lettuce germination. A similar result occurred with *Acacia mearnsii* and *Xerochrysum bracteatum* (Asteraceae) that grow in the same environment as boneseed (Al Harun et al. 2015a). Four phenolic compounds, catechins, p-coumaric acid, phloridzin, ferulic acid, were identified from boneseed tissue out of the 13 tested (Al Harun et al. 2015c). Leachate from boneseed leaf litter inhibited germination in *X. bracteatum* and *I. axillaris*, but unidentified allelochemicals could still be the cause of the allelopathy observed (Al Harun et al. 2015c). Further experiments using litter alone, litter and soil, and soil alone did not reduce lettuce germination above 20% indicating an effect (Al Harun et al. 2015b), albeit marginal in magnitude.

Management options for boneseed

The primary management tools that have been used to control boneseed infestations in Australia are manual weeding, controlled burns (i.e. fire), and herbicides (Table 1). For dense infestations, mechanical control can be deployed but unless stumps are removed from the ground, there is a danger that plants will resprout. Classical biocontrol has also been pursued for boneseed in Australia, including new developments and insight in the last two decades.

Fire

Fire can successfully control boneseed by killing adult plants and near-surface seeds in the soil. When used as part of an integrated approach (e.g., fire followed by herbicide treatment and/or hand-pulling of surviving plants), it is possible to extirpate boneseed via a controlled burn (i.e., lightly-invaded, intact ecosystems; Melland and Preston 2008). Small scale fire has been applied to boneseed in Western Australia in the past with good control outcomes (Peter Hennig pers. comm.). For example, at one site in Manypeaks, a small fire over 20-30 m² in 2015 resulted in the emergence of c. 250 seedlings the following year, which were subsequently controlled and no plants have been observed since (Peter Hennig pers. comm.). Lindenmayer et al. (2013) demonstrated that a too frequent fire regime (< 5 years interval) led to the vegetation being dominated by bracken, likely as a result of overgrazing of bitou bush and native plant species by wallabies (Dexter et al. 2013).

Herbicide

Much of the research on optimising herbicide use on boneseed was done before 2008 and is summarised in Brougham et al. (2006) and Weiss et al. (2008). Little has been published on improving herbicide effectiveness for boneseed in the last 15 years, noting that there is no evidence to show approved herbicides being or becoming ineffective against boneseed. Herbicide remains an effective control solution for boneseed, particularly at higher densities of plants.

Between 2006-2008 when large populations of boneseed were controlled in WA, herbicide was applied to foliage or by cut-and-paint, with follow-up treatment the following year. Herbicides registered for use on boneseed in Western Australia include: 2,4-D amine, bromoxynil, glyphosate, metsulfuron, picloram plus 2,4-D and metsulfuron plus glyphosate (Moore and Moore 2021).

Aerial spraying of herbicide has been used in eastern Australia against bitou bush since 2005 and has been shown to be effective for control of large dense infestations (boom spraying) or individual plants (cone application; Toth and Winkler 2008; Department of Planning and Environment NSW 2022). New application mechanisms that allow for targeted application via drones would be worth exploring for boneseed, as is being practiced for bitou bush (Department of Planning and Environment NSW 2022), particularly if plants are identified in open yet inaccessible areas.

Biological control

In South Africa, 113 phytophagous arthropods, three fungi and a mycoplasma have been found associated with *C. monilifera*, with 46 taxa (mostly insects) having potential as biological control agents (Scott and Adair 1995; Weiss et al. 1998).

Nine potential agents (arthropod species) have been released in Australia, three potential agents (two insect and one fungus) were studied but not released, and one potential agent was found already established as part of a biological control program for boneseed and bitou bush (Adair et al. 2012). Six of these species were approved for released to manage boneseed: three leaf beetles, *Chrysolina* spp; seed fly *Mesoclanis magnipalpis*; leaf-roller moth *Tortrix* sp. and leaf buckle mite *Aceria* sp. (Table 2). All except one species (buckle mite) failed to establish on boneseed, despite multiple releases (Downey et al. 2007; Morley 2010). One agent, the leaf buckle mite, is possibly established in Tasmania (Morley et al.

2012) and Victoria. It was recently observed in 2021 at Mt Eliza, Victoria at a 2008 release site, but had only spread 25 metres (Atlas of Living Australia, Biocontrol Hub; observation ID TM210214_01; <https://biocollect.ala.org.au/biocontrolhub>).

The release of *Tortrix* in New Zealand to target introduced boneseed was initially thought to be unsuccessful with only “patchy” establishment (Paynter et al. 2012) on the North Island and failure to establish on the South Island (Bownes 2022). By 2022, 11 years post release, establishment was reported for the South Island with indications of damage caused by the biological control agent (Bownes 2022).

An application to release boneseed rust (*Endophyllum osteospermi*) was not made due to the two-year lifecycle making it extremely difficult to do testing and get the necessary supporting data (L. Morin pers. comm.). Another pathogen, *Austropleospora osteospermi* (syn. *Hendersonia osteospermi*), is a leaf spot that somehow naturalised in Australia from southern Africa and is now widespread on bitou bush in coastal New South Wales (Morin et al. 2010) but with no (or limited) demonstrated impact. Its host range is limited to *Chrysanthemoides* and close relatives so it could be considered for use as a biological control agent against boneseed in Western Australia (Morin et al. 2010), but its absence from the state needs to be first confirmed.

As of 2012, there were no agents being actively researched in Australia (Adair et al. 2012). A lack of resourcing has resulted in no further boneseed biocontrol agent development since that time. The obvious question is why have so many biocontrol agents been worked up to a stage where they were approved for release, but have subsequently failed to establish on boneseed? Predation by native invertebrates appears to have hampered establishment or dispersal of *Chrysolina* spp. and *Tortrix* sp. (Adair et al. 2012). Other reasons could be a poor climate match or poor genetic match as part of the host matching process.

We now have a comprehensive knowledge of the genetics of the closely related bitou bush, based on nuclear and chloroplast genomes, which has enabled us to identify the source population in South Africa for the single introduction(s) to Australia (Emmett et al. 2023). This work has identified a previous unsearched region in South Africa where potentially more suitable or specific biological control agents could be found. In contrast boneseed has not been studied in relation to genetic variation or pollination syndromes, with only some glimpses of the genome where boneseed has been used as the outlier group in the study of bitou bush.

An improvement in host-agent matching would be a possible outcome from such molecular work. All past agents were reared on Australian boneseed plants while in quarantine. However, the nutrient and physical status of plants in the field may have been very different to those used for agent rearing and testing in quarantine (e.g., not fertilised, greater leaf toughness). It seems unlikely that climate mismatch between invaded areas and regions where agent searching was conducted has played a role in the lack of success, at least in south-eastern Australia. However, current climate matching models (Adair et al. 2012) are inadequate as projections extend implausibly into desert areas that are unlikely to be suitable for boneseed. If further agent searching was considered, it would be a priority to develop more robust process-based models to inform the agent development pipeline (Kriticos et al 2021).

Integrated control

Recent work on delivering more effective management outcomes for bitou bush control has coalesced around a management approach involving spraying with herbicide, burning, then re-spraying (Lindenmayer et al. 2015; O'Loughlin et al. 2019). A similar approach is recommended to eliminate boneseed (Melland et al. 1999; Melland and Preston 2008; Melland 2007). Six to twelve months before a fire, the infestation should be prepared by hand-pulling and cutting boneseed, so as to provide fuel at ground level, otherwise a fire might be too patchy (Melland et al. 1999).

Fire in autumn at 250 to 300°C will kill boneseed plants and deplete the seed bank (Brougham et al. 2006). Management of boneseed is achieved when followed in 18 months by herbicide treatment or hand-pulling of surviving plants or new germinants. This approach works best at a density of under 1000 seeds m² (i.e., a light infestation). For heavier seed bank densities (<1000 m²) fire treatments are only likely to produce lower population abundance rather than localised extirpation (Melland et al. 1999; Melland and Preston 2008; Melland 2007).

Detection

For management programs that are targeting localised extirpation or containment, improving detection of infestations and single plants for subsequent control is a priority. The role of citizen science in documenting new infestations, particularly via online reporting platforms

(Howard et al. 2022) has increased significantly in the last two decades (e.g. Landscape South Australia, 2020). Given that much of the control of boneseed across Australia is undertaken by community groups, online apps can be very effective tools.

New technologies have been applied to boneseed eradication programs to address the challenges of detecting rare plants in landscapes where operation is difficult due to access or terrain. Honey bee hives were tested as potential aggregation tool for environmental DNA (eDNA; pollen in this case) to detect boneseed plants in an urban landscape (Batchelor et al. 2023b). A species-specific assay was developed and a proof-of-concept trial was successful using pollen collected from bees foraging in dense boneseed populations in South Australia. However, no boneseed DNA was detected when the method was applied to material from hive pollen traps situated near isolated boneseed plants in Western Australia (Batchelor et al. 2023b). It may be that other eDNA substrates (Bell et al. 2024) could be developed for boneseed using the same PCR assay.

Detection of isolated plants in heterogeneous landscapes is also being widely addressed using drone-based remote sensing methods. These methods were applied to detecting low density boneseed plants in Western Australia with mixed success (Batchelor et al. 2023a). Both visible and multispectral band imagery stitched into an orthomosaic for subsequent image classification based on pixel reflectance values. The models ended up with high performance metrics but unacceptably high Type II error, a likely artefact of insufficient training imagery from Western Australia to develop the model (images from high density boneseed populations in South Australia were used, which likely created model transferability issues; Batchelor et al. 2023a). Similar drone-based methods for plant detection have recently been developed for bitou bush (Amarasingam et al. 2024). There may be learning opportunities to adapt this model for boneseed. However, habitat differences mean that boneseed is frequently found in environments with a tree overstorey (as opposed to the often-open dune environments occupied by introduced bitou bush), making the issue of image processing and analysis (i.e. orthomosaic vs single image workflows) an important consideration.

Lessons learned from management elsewhere

Management in south-eastern Australia

While Western Australia has been in the enviable position of being able to maintain an eradication goal for boneseed across the entire state, Victoria, South Australia and Tasmania (where boneseed is abundant) are mostly focusing on containment or asset protection. In 2014, The Eastern Australia Boneseed Eradication Project was working across southern and western New South Wales and eastern Victoria to eradicate all known infestations of boneseed from the former state and seeking to establish a new national containment line on the New South Wales border in a strategic, cross-jurisdictional effort (Martin 2013). This effort, combined with The South Coast Bitou Bush Task Force and relevant stakeholders, implemented the national southern bitou bush containment line at Tuross Head, New South Wales to prevent bitou bush from spreading south and hybridising with boneseed spreading up from Victoria (Cherry 2010; Cherry et al. 2006).

South Australia has a declared plant policy for boneseed under the Landscape South Australia Act 2019, which as of 2021 aimed to *eradicate* boneseed from the Northern, Yorke and Eyre Peninsula and South Australian Arid Lands, *contain* boneseed in the Murraylands and Riverland and Limestone Coast (via destruction of outlier populations and the Murray-Coorong Boneseed Containment Zone) and *asset protect* in the Hills and Fleurieu and Green Adelaide areas. More recently it has been accepted that it will not be feasible to eradicate boneseed from the Northern and Yorke region, which has moved to a *contain* approach (David Hughes, Landscape South Australia, pers. comm.). Kangaroo Island and Alintjara Wiluranara regions are still boneseed free (Government of South Australia 2021; Jason Walter, Landscape South Australia, pers. comm.).

In Tasmania, boneseed is a declared weed under the Biosecurity Act 2019. Eradication remains the stated aim in the northwest, where all boneseed has been controlled and local land managers are engaged (Cherry 2010, Department of Natural Resources and Environment Tasmania, 2011). In the south and northeast, all outlier populations are listed as control priorities and containment lines are maintained around core infestations to prevent further spread to areas known to be free (or in the process of becoming free) of boneseed (Cherry 2010, Department of Natural Resources and Environment Tasmania, 2011). We note that these plans are now more than a decade old and we were not able to confirm that an active coordinated program is still in place.

Boneseed is listed as a noxious weed (Schedule 2) in Victoria under the Catchment and Land Protections Act 1994 and was identified as an environmental weed with typically significant impacts and a 'high' risk rating by White et al. (2018). In Victoria there is regional level variation as to whether the weed is regionally prohibited or regionally controlled. What action is taken is often undertaken by community Landcare groups. Currently boneseed is listed for local eradication in East Gippsland, North Central and North East regions, whereas the weed is managed by prevention of growth and spread, particularly around high value conservation areas, in Corangamite, Glenelg Hopkins, Goulburn, Port Phillip and Westernport, and West Gippsland regions. Control in the Wimmera and Mallee regions was initially focused on eradication but has now transitioned to regional control only. At Port Phillip and Westernport there appears to be ongoing eradication efforts and asset protection targeting a hybrid bitou bush boneseed population.

Across all south-eastern states in Australia, the methods being deployed for boneseed management remain consistent with those outlined in previous guidelines (e.g. Brougham et al. 2006). The failure of classical biological control agents to establish in Australia has meant that manual control, herbicide treatments, managed burns and occasionally mechanical control are the dominant approaches deployed, often in combination.

We were unable to confirm any records of boneseed from Norfolk Island, a small Australian territory in the Pacific Ocean, as reported by Mariotti and Zappa (2022). Rather, this report appears to be a mis-identification of bitou bush, which was recently noted as a new record for the remote island (Martoni et al. 2023). The infestation was first documented in 2011 under a large Norfolk Island pine tree and spread from a small area (c. 2m²) to cover over 100m² at its peak in around 2021 (T. Patel, Norfolk Island Regional Council, pers. comm.). The infestation has been actively managed in recent years with the main infestation mechanically mulched and individual plants that had established on the nearby coastal cliffs being hand weeded (T. Patel, Norfolk Island Regional Council, pers. comm.). Learnings from bitou bush eradications in Western Australia (Scott et al. 2019a) and Queensland (Cherry et al. 2008) are of clear relevance for a successful eradication of bitou bush on Norfolk Island.

Management in New Zealand

Boneseed is also a non-native invasive weed in New Zealand. It was first recorded in 1870 and after a very long lag phase became prominent in the 1990s (Briden 2008). Invasion initially was localised to urban areas, but now occurs in a wide range of native vegetation and

situations, from dunes to islands (Briden 2008). The strategy in New Zealand is to have surveillance and weed-led control to find and eradicate new infestations, and a site-led approach to manage large infestations in valuable ecosystems.

The range of control methods, manual, herbicide, mechanical, and biological control are used in New Zealand, similarly to the controls outlined in the Boneseed Management Manual (Brougham et al. 2006; Table 1). An additional control method used in New Zealand is the “mechanical shredder”, a mechanical mulcher mounted in an all-terrain vehicle with rubber tracks (Briden 2008). Biological control activities have recently been revived with the redistribution of *Tortrix* sp. encouraged now that it is established in the South Island (Bownes 2022).

The New Zealand experience is that the effort required for control decreases over time. Most effort is required in the first 1 to 3 years with the removal of large plants and control of seedlings. After 5-6 years the ongoing seed bank is much reduced and ongoing maintenance takes little effort (Briden 2008). However, this approach does not consider the impact of fire. A boneseed infestation cleared 8.5 years previous had a massive germination of seeds from the soil seedbank following a fire (Briden and McAlpine 2012). There is also evidence that soil disturbance and canopy removal stimulates germination of boneseed seeds over native plants, indicating that any form of disturbance to an ecosystem will favour germination of boneseed (McAlpine et al. 2009). Howell (2012) assessed ten years of progress towards environmental weed eradication in New Zealand. His sample of 90 eradication programs included two on boneseed that had made zero progress. The main conclusion of this work was that to succeed with any weed eradication it pays to start few programs and to focus on those most likely to succeed.

Management elsewhere (Chile, California, and Europe)

Atala et al. (2023) describe the impact on local species diversity of *C. monilifera* in Valparaíso, Chile, without identifying the subspecies involved. Numerous photos are available online of the study area showing flowers and fruits (Fundación RA Philippi de Estudios Naturales 2024) which enables us to conclude that the subspecies is most likely *C. m. monilifera*. The invasion is attributed to the introduction as an ornamental plant to the Quinta Vergara Park in Viña del Mar, Chile, from which it escaped cultivation and spread to inland dunes of the Valparaíso region (Atala et al. 2023). We could not find any evidence of management for this invasion and would suggest that further work to confirm identification would be prudent.

Similarly, there doesn't appear to be active management of boneseed in California where it is cultivated as a garden plant and naturalised, but reportedly infrequently escapes and/or rarely persists in the flora (Brusati et al. 2014; Strother 2006).

Boneseed is recorded from multiple areas across Europe, including France (Channel Islands, Saint Raphaël, Théoule sur Mer) Monaco, Italy (Sicily, Ventimiglia), Gibraltar and Andorra (Bock 2024; Greuter 2006; Mariotti and Zappa 2022). We were not able to obtain independent verification of the report from Andorra. This location, given its high altitude and winter climate, seems an unlikely spot for boneseed to establish. In Italy, Mariotti and Zappa (2022) report that the Sicilian population of boneseed has been eradicated, and that the Ventimiglia introduction took place in 1869 via seeds planted at the Hanbury Botanic Gardens. The Ventimiglia population was reported as naturalised in 1996 as part of the first survey to assess the status of non-native plants of the area.

Occasional reports of 'boneseed' being introduced to Saint Helena, a remote island in the South Atlantic Ocean, are likely a misreported occurrence based on common name confusions. The closely related *Osteospermum sanctae-helenae*, an endemic to Saint Helena, is also known as boneseed (Cronk 1987).

Updated insight to guide boneseed management

In the near four decades since boneseed was first targeted for control in Western Australia, management outcomes have had mixed success. Existing ecological knowledge has generally been sufficient to guide effective management choices, particularly with sparsely distributed plants. However, if eradication is going to be achieved in the state, a step change in management will be required over a considerable duration. The recent assessment of past management and current distribution (Batchelor et al. 2024) provides a robust platform for launching a management program in Western Australia. This review has delivered additional complementary insight across four areas that could help to further improve the program in Western Australia, as well as boneseed management programs elsewhere.

First, the origins and introduction pathway for boneseed in Western Australia remain unclear. Improving understanding in this regard, particularly if combined with insight across the full Australian and South African boneseed distribution, would help to guide genetically-informed native range surveys for classical biological control agents. It is important to note that classical biological control remains an unfeasible management option for Western Australia, particularly given the lack of any notable effectiveness elsewhere and the difficulty

of maintaining agent presence with the very low ongoing abundance of mature plants. More generally, however, the recently reported positive impacts of *Tortrix* on New Zealand boneseed suggest there may be benefit for re-evaluating this agent failure elsewhere where containment or minimising impact is a management goal (e.g. south-eastern Australia).

Second, improved knowledge on the ecology and biology of boneseed has reinforced that currently deployed control methods remain the most effective for using in an eradication program for Western Australia. Important knowledge gaps remain with respect to determining whether boneseed is capable of self-fertilization (autogamy), with determining the full cohort of dispersal agents across the introduced range. Both emus and the currently controlled starlings could challenge the working assumption that a 500 m buffer zone (based on dispersal kernels) is adequate to inform population delimitation in Western Australia (Batchelor et al. 2024). Some ecological knowledge gaps have also been addressed to provide management insight. Completely extinguishing the soil seedbank remains the biggest threat to a successful eradication campaign (Panetta 2004). Ten years would be a minimum timeframe to actively manage a boneseed seedbank with 15 years as a more conservative duration likely necessary for eradication programs. Evidence for using chemical amendments to accelerate seedbank depletion is inconclusive, yet fire appears to cause significant seed mortality.

Third, insight from active programs suggests that a combination of manual, chemical and fire-based control, with mechanical removal for large infestations, is still optimal for boneseed management in Western Australia. These learnings could also be applied to overseas introductions where active management seems absent. An opportunity to eradicate boneseed before it has spread widely, particularly in areas with Mediterranean climates, appears to be worth prioritising, even if confirmed identifications found the introduction to be another taxon of non-native *Chrysanthemoides*.

Last, novel detection techniques for isolated plants in heterogenous or hard to access landscapes could transform the likelihood of achieving successful eradication for boneseed in Western Australia. Such techniques are equally applicable to maintaining containment lines for introduced boneseed elsewhere. While initial attempts to develop eDNA and drone-based remote sensing methods to detect boneseed were not successful, they generated promising insight that could be further refined in future work.

This synthesis from invasions in Australia's south-eastern states and elsewhere overseas, combined with updated ecological knowledge of the local weed context, suggests that boneseed continues to represent a significant weed threat to natural ecosystems in Western Australia. Integrating knowledge obtained from this literature review with a collated baseline of past management efforts (Batchelor et al. 2024) to produce robust and enduring management programs will give land managers the best chance of achieving their eradication objectives.

Acknowledgments

This work was funded by CSIRO and DPIRD's Boosting Biosecurity Defences program and made possible by the Royalties for Regions program.

Competing Interests

No competing interests have been declared.

References

- Adair RJ and Butler KL (2010) Hybridisation within *Chrysanthemoides monilifera* and implications for the eradication of bitou bush from Victoria, Australia. Pages 378-381 in Zydenbos SM, ed. Proceedings of the 17th Australasian Weeds Conference. 26-30 September, 2010, Christchurch, New Zealand
- Adair RJ and Holtkamp RH (1999) Development of a pesticide exclusion technique for assessing the impact of biological control agents for *Chrysanthemoides monilifera*. *Biocontrol Sci Tech* 9: 383-390
- Adair RJ, Morley T, and Morin L (2012) *Chrysanthemoides monilifera* (L.) T.Norl. - bitou bush and boneseed. Pages 170-183 in Julien M, McFadyen R, Cullen J, eds. *Biological control of weeds in Australia*. Collingwood, Victoria, Australia: CSIRO Publishing
- Al Harun MAY, Johnson J, and Robinson RW (2015a) The contribution of volatilization and exudation to the allelopathic phytotoxicity of invasive *Chrysanthemoides monilifera* subsp. *monilifera* (boneseed). *Biol Invasions* 17: 3609-3624
- Al Harun MAY, Johnson J, Uddin MN, and Robinson RW (2015b) The effects of temperature on decomposition and allelopathic phytotoxicity of boneseed litter. *J Environ Sci-China* 33: 1-11

- Al Harun MAY, Johnson J, Uddin MN, and Robinson RW (2015c) Identification and phytotoxicity assessment of phenolic compounds in *Chrysanthemoides monilifera* subsp. *monilifera* (Boneseed). PLoS One 10: e0139992
- Al Harun MAY, Robinson RW, Johnson J, and Uddin MN (2014) Allelopathic potential of *Chrysanthemoides monilifera* subsp. *monilifera* (boneseed): A novel weapon in the invasion processes. S Afr J Bot 93: 157-166
- Amarasingam N, Kelly JE, Sandino J, Hamilton M, Gonzalez F, Dehaan RL, Zheng L, and Cherry H (2024) Bitou bush detection and mapping using UAV-based multispectral and hyperspectral imagery and artificial intelligence. Remote Sensing Applications: Society and Environment 34: 101151
- Atala C, Reyes SA, Osses J, Jeldes-Cajas O, and Vargas R (2023) The invader shrub *Chrysanthemoides monilifera* (boneseed) negatively impacts native plant communities in a Mediterranean zone in Central Chile. Brazil J Bot 46: 695-703
- Barker NP, Howis S, Nordenstam B, Källersjö M, Eldenäs P, Griffioen C, and Linder HP (2009) Nuclear and chloroplast DNA-based phylogenies of *Chrysanthemoides* Tourn. ex Medik. (Calenduleae; Asteraceae) reveal extensive incongruence and generic paraphyly, but support the recognition of infraspecific taxa in *C. monilifera*. S Afr J Bot 75: 560-572
- Barker NP, Paterson I, and Howis S (2015) "Barcoding" and ISSR data illuminate a problematic infraspecific taxonomy for *Chrysanthemoides monilifera* (Calenduleae; Asteraceae): Lessons for biocontrol of a noxious weed. Biochem Syst Ecol 61: 541-551
- Barker NP, von Senger I, Howis S, Zachariades C, and Ripley BS (2005) Plant phylogeography based on rDNA ITS sequence data: two examples from the Asteraceae. Pages 217-244 in Bakker FT, Chatrou LW, Gravendeel B, Pelsers PB, eds. Plant Species-Level Systematics: New Perspectives on Pattern & Process. Ruggell, Liechtenstein: A.R.G. Gantner Verlag

- Batchelor, K.L., Scott, J.K. and Webber, B.L. (2022) Boneseed and bitou bush in Western Australia: the tale of two *Chrysanthemoides*. Proceedings of the 22nd Australasian Weeds Conference, 25-29 September 2022, Adelaide, South Australia. Ed. Rachel Melland, R., Brodie, C., Emms, J., Feuerherdt, L., Ivory, S. and Potter, S. Published by Weed Management Society of South Australia Inc. pp 7-10
- Batchelor KL, Yeoh PB, Bell KL, Campos M, Ota N, Richetti J, and Webber BL (2023a) Boneseed management in Western Australia. Improved methods for population depletion and detection to enhance the effectiveness of control programs. CSIRO, Perth, Australia. 89 p
- Batchelor KL, Bell KL, Campos M, and Webber BL (2023b) Can honey bees be used to detect rare plants? Taking an eDNA approach to find the last plants in a weed eradication program. *Environmental DNA* 5: 1516-1526
- Batchelor KL, Scott JK, and Webber BL (2024) Informing eradication feasibility: ecological context and delimitation for boneseed (*Chrysanthemoides monilifera* subsp. *monilifera*) in Western Australia. *Invasive Plant Sci Manag.* doi: 10.1017/inp.2024.18 (Accepted 17 Jun 2024)
- Bell KL, Campos M, Hoffmann BD, Encinas-Viso F, Hunter GC, and Webber BL (2024) Environmental DNA methods for biosecurity and invasion biology in terrestrial ecosystems: progress, pitfalls, and prospects. *Science of the Total Environment* 926: 171810
- Blood K (1987) Flora of the You Yangs Forest Parks. Victorian College of Agriculture and Horticulture, Burnley. 135 p
- Bock B (2024) Photoflora <http://www.photoflora.fr/FiTax.php?NumTaxon=9356>. Accessed 28 Oct 2024
- Bownes A (2022) It established after all. <https://www.landcareresearch.co.nz/publications/weed-biocontrol/weed-biocontrol-articles/it-established-after-all/>. Accessed 8 May 2024
- Bramwells H (2003) Flora and Fauna Guarantee Action Statement #63: Brittle Greenhood *Pterostylis truncata* <http://www.dse.vic.gov.au>. Accessed 22 Dec 2021

- Briden K (2008) Current status and management of boneseed in New Zealand. *Plant Prot Quart* 23: 20-22
- Briden K and McAlpine KG (2012) Wildfire stimulates boneseed germination 8.5 years after adult plants cleared. Pages 81-81 *in* Eldershaw V, ed. Proceedings of the 18th Australasian Weeds Conference. Melbourne, Victoria, Australia, 8-11 October 2012
- Brougham KJ, Cherry H, and Downey PO (2006) Boneseed management manual: current management and control options for boneseed (*Chrysanthemoides monilifera* ssp. *monilifera*) in Australia. Sydney South NSW: Department of Environment and Conservation NSW. 82 p
- Brusati ED, Johnson DW, and DiTomaso JM (2014) Predicting invasive plants in California. *California Agriculture* 68: 89-95
- Byrne D, Scheben A, Scott JK, Batchelor KL, Severn-Ellis AA, Webber BL, Gooden B, and Bell KL (2022) Genomics reveals the history of a complex plant invasion and improves the management of a biological invasion from the South African-Australian biotic exchange. *Ecol Evol* 12: e9179
- Cherry H (2010) Boneseed in Australia: research, management and coordinated action. Pages 81-84 *in* Zydenbos SM, ed. Proceedings of the 17th Australasian Weeds Conference. 26-30 September, 2010, Christchurch, New Zealand
- Cherry H, Leighton S, Clark A, and Austin P (2006) National weed containment and eradication zones: challenges, successes, and lessons learned. Pages 44-47 *in* Preston C, Watts JH, Crossman ND, eds. Proceedings of the 15th Australian Weeds Conference. 24-28 September 2006, Adelaide, South Australia
- Cherry, H, Willsher, L, and Whyte, B (2008) Bitou bush and boneseed eradication and containment in Australia. *Plant Prot Quart* 23: 38-40.
- Cronk QCB (1987) The history of endemic flora of St Helena: a relictual series. *New Phyt* 105: 509-520

- Department of Natural Resources and Environment Tasmania (2011) Boneseed – statutory weed management plan. Department of Natural Resources and Environment, Hobart, Tasmania. 17p
- Department of Planning and Environment NSW (2022) Aerial spraying guidelines. Environment and Heritage, Department of Planning and Environment, Paramatta, NSW. 63 p
- Dexter N, Hudson M, James S, MacGregor C, and Lindenmayer DB (2013) Unintended consequences of invasive predator control in an Australian forest: Overabundant wallabies and vegetation change. PLoS One 8: e69087
- Downey PO, Holtkamp RH, Ireson JE, Kwong RM, and Swirepik AE (2007) A review of the *Chrysanthemoides monilifera* biological control program in Australia: 1987-2005. Plant Prot Quart 22: 24-32
- Duffy J (2020) How boneseed weed affects koalas
<https://www.echidnawalkabout.com.au/how-boneseed-weed-affects-koalas/>. Accessed 18 Dec 2021
- Emmett AM, Scott JK, Webber BL, Severn-Ellis AA, and Bell KL (2023) Combining multiple lines of evidence to elucidate the origin and introduction pathway of bitou bush (*Chrysanthemoides monilifera* subsp. *rotundata*) in Australia. Biol Invasions 25: 1881-1905
- Ens E-J (2007) Indigenous plant recruitment limitation by bitou bush (*Chrysanthemoides monilifera* spp. *rotundata*): effect on life history stages and allelopathic mechanisms. PhD Thesis: University of Wollongong. 154 p
- Ens E-J and French K (2008) Exotic woody invader limits the recruitment of three indigenous plant species. Biol Conserv 141: 590-595
- Ens E-J, French K, and Bremner JB (2008) Bitou bush invasion is facilitated by soil chemistry changes which inhibit the growth of native plants. Pages 104-106 in van Klinken RD, Osten VA, Panetta FD, Scanlan JC, eds. Proceedings of the 16th Australian Weeds Conference. Cairns, North Queensland, Australia, 18-22 May, 2008

- Ens E-J, French K, Bremner JB, and Korth J (2010) Novel technique shows different hydrophobic chemical signatures of exotic and indigenous plant soils with similar effects of extracts on indigenous species seedling growth. *Plant and Soil* 326: 403-414
- Ens EJ, Bremner JB, French K, and Korth J (2009a) Identification of volatile compounds released by roots of an invasive plant, bitou bush (*Chrysanthemoides monilifera* spp. *rotundata*), and their inhibition of native seedling growth. *Biol Invasions* 11: 275-287
- Ens EJ, French K, and Bremner JB (2009b) Evidence for allelopathy as a mechanism of community composition change by an invasive exotic shrub, *Chrysanthemoides monilifera* spp. *rotundata*. *Plant and Soil* 316: 125-137
- French K, Ashcroft MB, Panetta FD, Raghu S, and Cherry H (2024) Differences in soil seed longevity for two subspecies of the invasive weed, *Chrysanthemoides monilifera* (Asteraceae). *Weed BiolManag* 24: 104-112
- Fundación RA Philippi de Estudios Naturales (2024) *Chrysanthemoides monilifera* <https://fundacionphilippi.cl/catalogo/chrysanthemoides-monilifera/>. Accessed 21 May 2024
- Gosper CR (2003) Consequences of weed invasion and control of plant-bird interactions and bird communities. PhD Thesis: University of Wollongong. 190 p
- Government of South Australia (2021) Declared Plant Policy: Boneseed (*Chrysanthemoides monilifera*) https://pir.sa.gov.au/__data/assets/pdf_file/0019/137314/boneseed_policy.pdf Accessed 8 Dec 2021
- Gray M (1976) Miscellaneous notes on Australian plants. 2. *Chrysanthemoides* (Compositae). *Contributions from Herbarium Australiense* 16: 1-5
- Greuter W (2006) Compositae (pro parte majore) in Greuter W, von Raab-Straube E, eds. *Compositae Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity*

- Gross CL, Whitehead JD, de Souza CS, and Mackay D (2017) Unsuccessful introduced biocontrol agents can act as pollinators of invasive weeds: Bitou bush (*Chrysanthemoides monilifera* ssp. *rotundata*) as an example. *Ecol Evol* 7: 8643-8656
- Howard L, van Rees CB, Dahlquist Z, Luikart G, and Hand BK (2022) A review of invasive species reporting apps for citizen science and opportunities for innovation. *NeoBiota* 71: 165-188.
- Howell CJ (2012) Progress toward environmental weed eradication in New Zealand. *Invasive Plant Sci Manag* 5: 249-258
- Knight RS (1988) Aspects of plant dispersal in the southwestern Cape with particular reference to the roles of birds as dispersal agents. PhD Thesis: University of Cape Town. 320 p
- Kriticos DJ, Ireland KB, Morin L, Kumaran N, Rafter MA, Ota N, and Raghu S (2021) Integrating ecoclimatic niche modelling methods into classical biological control programmes. *Biol Control* 160: 104667
- Landscape South Australia (2020) Battle waged on boneseed. <https://www.landscape.sa.gov.au/mr/news/201116-battle-waged-on-boneseed-nws>. Accessed 19 November 2024
- Lindenmayer DB, MacGregor C, Dexter N, Fortescue M, and Cochrane P (2013) Booderee National Park Management: Connecting science and management. *Ecol Manag Rest* 14: 2-10
- Lindenmayer DB, Wood J, MacGregor C, Buckley YM, Dexter N, Fortescue M, Hobbs RJ, and Catford JA (2015) A long-term experimental case study of the ecological effectiveness and cost effectiveness of invasive plant management in achieving conservation goals: Bitou bush control in Booderee National Park in eastern Australia. *PLoS One* 10: e0128482
- Mariotti M and Zappa E (2022) Remarks on the exotic flora of Capo Mortola (Ventimiglia, northern Italy) and its changes over time. *Italian Botanist* 14: 1-43

- Martin P (2013) Eastern Australia Boneseed Eradication Project. A state level partnership with local outcomes: eradicating outlier boneseed infestations from Sydney to the Victorian border 17th NSW Weeds Conference Proceedings. RSL Club, Corowa, New South Wales
- Martoni F, Achari S, Blacket M, Brohier N, Constable F, Dugdale T, Ekanayake P, Kant P, Kelly G, Kinoti C, Li T, Lovelock D, Mann R, Nogarotto E, Sawbridge T, Smith R, Wainer J, and Rodoni B (2023) Plant pest and disease survey on Norfolk Island Final Project Report: The State of Victoria Department of Energy, Environment and Climate Change. 140 p
- McAlpine KG, Timmins SM, and Westbrooke I (2009) Bone-seed (*Chrysanthemoides monilifera* ssp. *monilifera*) invasion effects on native regeneration in New Zealand coastal plant communities. *N Z J Ecol* 33: 72-82
- Melland R, Ainsworth N, and Roush RT (1999) Reducing the fitness of the fittest: how to control a *Chrysanthemoides monilifera* ssp. *monilifera* (Norlich) invasion. Pages 294-297 in Bishop AC, Boersma M, Barnes CD, eds. Proceedings of the 12th Australian Weeds Conference. 12-16 September 1999, Hobart, Tasmania, Australia
- Melland R and Preston C (2008) The role of fire in integrated management of boneseed (*Chrysanthemoides monilifera* subsp. *monilifera*). *Plant Prot Quart* 23: 32-33
- Melland RL (2007) Management of boneseed (*Chrysanthemoides monilifera* ssp. *monilifera* (L.) T.Norl. using fire, herbicides and other techniques in Australian woodlands. PhD Thesis: University of Adelaide. 219 p
- Mokotjomela TM, Musil CF, and Esler KJ (2013a) Do frugivorous birds concentrate their foraging activities on those alien plants with the most abundant and nutritious fruits in the South African Mediterranean-climate region? *Plant Ecol* 214: 49-59
- Mokotjomela TM, Musil CF, and Esler KJ (2013b) Frugivorous birds visit fruits of emerging alien shrub species more frequently than those of native shrub species in the South African Mediterranean climate region. *S Afr J Bot* 86: 73-78

- Mokotjomela TM, Musil CF, and Esler KJ (2013c) Potential seed dispersal distances of native and non-native fleshy fruiting shrubs in the South African Mediterranean climate region. *Plant Ecol* 214: 1127-1137
- Moore CB and Moore JH (2021) HerbiGuide - The Pesticide Expert on a Disk www.herbiguide.com.au.
- Morin L, Shivas RG, Piper MC, and Tan YP (2010) *Austropleospora osteospermi* gen. et sp. nov. and its host specificity and distribution on *Chrysanthemoides monilifera* ssp. *rotundata* in Australia. *Fungal Diversity* 40: 65-74
- Morley T (2010) A new attempt to introduce the lacy-winged seed fly *Mesoclanis magnipalpis* to Australia for biological control of boneseed *Chrysanthemoides monilifera* subsp. *monilifera*. Pages 243-244 in Zydenbos SM, ed. Proceedings of the 17th Australasian Weeds Conference. Christchurch, New Zealand, 26-30 September, 2010
- Morley T, Ireson J, and Ivory S (2012) Attempts to establish the boneseed leaf buckle mite in Australia for the biological control of boneseed. Pages 353-355 in Eldershaw V, ed. Proceedings of the 18th Australasian Weeds Conference. Melbourne, Victoria, Australia, 8-11 October 2012
- Muyt A (2001) Bush invaders of South-East Australia : a guide to the identification and control of environmental weeds found in South-East Australia. Meredith, Vic.: R.G. & F.J. Richardson. Pp. 304
- Newman K (1983) Newman's birds of southern Africa. Johannesburg: MacMillan South Africa Publishers. Pp. 461
- Norlindh T (1943) Studies in the Calenduleae. 1. Monograph of the genera *Dimorphotheca*, *Castalis*, *Osteospermum*, *Gibbaria* and *Chrysanthemoides*. Lund: CWK Gleerup. Pp. 367- 423
- O'Loughlin LS, Gooden B, Foster CN, MacGregor CI, Catford JA, and Lindenmayer DB (2019) Invasive shrub re-establishment following management has contrasting effects on biodiversity. *Sci Rep* 9: 1-11

- Panetta FD (2004) Seed banks: the bane of the weed eradicator. Pages 523-526 in Sindel BM, Johnson SB, eds. Proceedings of the 14th Australian Weeds Conference. Wagga Wagga, New South Wales, 6-9 September 2004
- Paynter Q, Forgie SA, Winks CJ, Peterson PG, Ward DF, Nicholson L, and Van Zoelen R (2012) Biotic resistance: Facilitation between invasive Homoptera and invasive ants limits the establishment of an introduced weed biocontrol agent in New Zealand. *Biol Control* 63: 188-194
- POWO (2024) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet <http://www.plantsoftheworldonline.org/>. Accessed 06 Jun 2024
- Reynolds CJ, Long RL, Flematti GR, Cherry H, and Turner SR (2014) Karrikins promote germination of physiologically dormant seeds of *Chrysanthemoides monilifera* ssp. *monilifera* (boneseed). *Weed Res* 54: 48-57
- Sadler R, Parker T, Verboom GA, Ellis AG, Jackson M, van Zyl J, Manning J, and Bergh NG (2022) A molecular phylogeny of Calenduleae (Asteraceae) supports the transfer of *Dimorphotheca polyptera* DC. to *Osteospermum* L. *S Afr J Bot* 151: 234-245
- Schoeman J, Buckley YM, Cherry H, Long RL, and Steadman KJ (2010) Inter-population variation in seed longevity for two invasive weeds: *Chrysanthemoides monilifera* ssp. *monilifera* (boneseed) and ssp. *rotundata* (bitou bush). *Weed Res* 50: 67-75
- Scott JK (1996) Population ecology of *Chrysanthemoides monilifera* in South Africa: Implications for its control in Australia. *J Appl Ecol* 33: 1496-1508
- Scott JK and Adair RJ (1995) Selection of biological control agents for *Chrysanthemoides monilifera*, a weed of protected natural areas in Australia. Pages 61-67 in Delfosse ES, Scott RR, eds. Proceedings of the VII International Symposium on Biological Control of Weeds. 2-7 February 1992, Christchurch, New Zealand: DSIR/CSIRO
- Scott JK, Batchelor KL, Jucker T, and Webber BL (2019a) Aerial photography and dendrochronology as tools for recreating invasion histories: do they work for bitou bush (*Chrysanthemoides monilifera* subsp. *rotundata*)? *Biol Invasions* 21: 2983-2996

- Scott JK, Batchelor KL, and Webber BL (2019b) Long term monitoring of recruitment dynamics determines eradication feasibility for an introduced coastal weed. *NeoBiota* 50: 31-53
- Scurr G, Kirkpatrick JB, Daniels GD, and McQuillan PB (2008) Biotic resistance to *Chrysanthemoides monilifera* ssp. *monilifera* in Tasmania. *Aust Ecol* 33: 941-950
- Strother JL (2006) *Chrysanthemoides monilifera* (Linnaeus) Norlindh. Pages 380 *Flora of North America*
- Thomas PB, Possingham H, and Roush R (2005) Effects of boneseed (*Chrysanthemoides monilifera* (L.) Norl. ssp. *monilifera*) on the composition of the vegetation and the soil seed bank of an open eucalypt woodland. *Plant Prot Quart* 20: 74-80
- Toth J and Winkler MA (2008) Bitou bush aerial spraying in New South Wales - what have we learned? *Plant Prot Quart* 23: 43-44
- Vranjic JA, Woods MJ, and Barnard J (2000) Soil-mediated effects on germination and seedling growth of coastal wattle (*Acacia sophorae*) by the environmental weed, bitou bush (*Chrysanthemoides monilifera* ssp. *rotundata*). *Aust Ecol* 25: 445-453
- Weiss PW (1984) Seed characteristics and regeneration of the species in invaded coastal communities. *Aust J Ecol* 9: 99-106
- Weiss PW, Adair RJ, and Edwards PB (1998) *Chrysanthemoides monilifera* (L.) T. Norl. Pages 49-61 in Panetta FD, Groves RH, Shepherd RCH, eds. *The biology of Australian weeds Volume 2*. Melbourne: R.G. and F.J. Richardson
- Weiss PW, Adair RJ, Edwards PB, Winkler MA, and Downey PO (2008) *Chrysanthemoides monilifera* subsp. *monilifera* (L.) T.Norl. and subsp. *rotundata* (DC.) T.Norl. *Plant Prot Quart* 23: 3-14
- White, M, Cheal, D, Carr, G W, Adair, R, Blood, K, and Meagher, D (2018) Advisory list of environmental weeds in Victoria. Arthur Rylah Institute for Environmental Research Technical Report Series No. 287. Department of Environment, Land, Water and Planning, Heidelberg, Victoria. 102 p

Table 1. Summary of methods used to control boneseed (*Chrysanthemoides monilifera* spp. *monilifera*) in Australia (adapted from Brougham et al. (2006) with updates).

Type of control	Method	Specific details
Manual control	Hand pulling	Boneseed has a relative shallow root system and most plants can be pulled from the soil intact.
Chemical control	Cut stump	Plants are cut off at the base and herbicide applied to the stump.
	Stem injection	Plants are drilled at the base of the trunk and herbicide added to holes.
	Foliar spraying	Herbicide is applied to leaves and stems as a fine spray.
	Splatter gun	Spraying with large droplet size, suitable for targeting large plants, not often used against boneseed.
Mechanical control	Mechanical pulling	Use heavy machinery to pull plants from the ground, with minimal disturbance to the soil. Suitable for agricultural areas and very large plants.
	Slashing	Suitable for non-natural areas, but stumps will resprout.
Fire	Effect on boneseed	Fire kills plants if entirely scorched. Fire will kill some seeds but stimulates mass germination.
	Control opportunities	If a bushfire burns a boneseed infested area, be prepared for follow-up control of seedlings.
Biological control	Insects, mites and pathogens	Nine biological control agents released between 1989 and 2006, none established on boneseed (Adair et al. 2012; Table 2).

Table 2. Biological control releases on boneseed (*Chrysanthemoides monilifera* spp. *monilifera*) in Australia adapted from Brougham et al. (2006) and Adair et al. (2012) with updates.

Biological control agent	Biological control agent years released	Agent release events	Establishment summary	Current status
Bitou tip moth, <i>Comostolopsis germana</i>	1989 - 1998	37 (Vic, SA, Tas)	Did not establish – possibly due to predation or poor climate match	Established and widespread on bitou bush but not boneseed in eastern Australia
Black boneseed leaf beetle, <i>Chrysolina scotti</i>	1989 - 1996	18 (Vic), 5 (SA), 10 (NSW), 10 (Tas)	1 Did not establish – possibly due to predation	Not present in Australia
Blotched boneseed leaf beetle, <i>Chrysolina picturata</i>	1992 - 1995	4 (Vic), 8 (Tas), 8 (NSW)	2 Did not establish – possibly due to predation	Not present in Australia
Painted boneseed beetle, <i>Chrysolina oberprieleri</i>	1994 - 1995	7 (Vic), 2 (SA)	2 Did not establish – possibly due to predation	Not present in Australia
Lacy-winged seed fly, <i>Mesoclanis magnipalpis</i>	1998 - 2000 & 2005 (onto bitou bush and boneseed)	? (released onto boneseed hybrids in Vic and NSW)	Released onto bitou bush, did not establish – small release numbers	Not present in Australia
Seed fly, <i>Mesoclanis polana</i>	1998	0 (however, sampled on boneseed)	Did not establish – poor host match	Widespread and abundant on bitou bush but not

			where it overlaps with bitou bush)		boneseed in eastern Australia
Boneseed rolling moth, <i>Tortrix</i> sp.	leaf- 2000 - 2004	112 (Vic), >9 (NSW)	Did not establish – possibly due to predation and competition in larval stages	Established on bitou bush but not boneseed in eastern Australia. Established on boneseed in North and South Island NZ	
Leaf buckle mite, <i>Aceria</i> sp.	2008 - 2012	90 (Vic, SA, Tas)	12 months after release, seen at 4 sites in Vic, 1 in Tas. Not established in SA	Last presence survey in SA (2011), Tas (2012), and Vic (2021). A 2012 survey in SA did not find agents	
Boneseed fungus, <i>Endophyllum osteospermi</i>	rust	n/a	n/a	Host-specificity testing abandoned – too difficult	
Leaf spot fungus, <i>Austropleospora osteospermi</i>	Naturalised in eastern Australia	n/a	South African species on bitou bush	Established in New South Wales on bitou bush, not observed on boneseed but (Morin et al. 2010) demonstrates susceptibility in lab conditions	