

SOLID BASES AND FUNCTORIAL CONSTRUCTIONS FOR (P -)BANACH SPACES OF ANALYTIC FUNCTIONS

GUOZHENG CHENG¹, XIANG FANG², CHAO LIU³ AND YUFENG LU¹

¹*School of Mathematical Sciences, Dalian University of Technology, Dalian,
P. R. China*

²*Department of Mathematics, National Central University, Chungli, Taiwan*
³*School of Mathematics and Statistics, Yunnan University, Kunming, P. R. China*

Corresponding author: Chao Liu, email: chaoliu@ynu.edu.cn

(Received 21 July 2023)

Abstract Motivated by new examples of functional Banach spaces over the unit disk, arising as the symbol spaces in the study of random analytic functions, for which the monomials $\{z^n\}_{n \geq 0}$ exhibit features of an unconditional basis yet they often don't even form a Schauder basis, we introduce a notion called *solid basis* for Banach spaces and p -Banach spaces and study its properties. Besides justifying the rich existence of solid bases, we study their relationship with unconditional bases, the weak-star convergence of Taylor polynomials, the problem of a solid span and the curious roles played by c_0 . The two features of this work are as follows: (1) during the process, we are led to revisit the axioms satisfied by a typical Banach space of analytic functions over the unit disk, leading to a notion of \mathcal{X}^{\max} (and \mathcal{X}^{\min}), as well as a number of related functorial constructions, which are of independent interests; (2) the main interests of solid basis lie in the case of non-separable (p -)Banach spaces, such as BMOA and the Bloch space instead of VMOA and the little Bloch space.

Keywords: random analytic functions; norm convergence of Taylor polynomials; solid spaces; BMOA; Bloch spaces; unconditional bases

2020 Mathematics subject classification: 30B20; 47B37

1. Introduction and main results

A number of new Banach spaces of analytic functions over the unit disk arise naturally as the symbol spaces during the study of random analytic functions in Hardy spaces, Bergman spaces, Dirichlet spaces and their multipliers, analytic functions of bounded mean oscillation (BMOA), etc. [11–13, 29, 32, 34]. Let \mathcal{X} be a Banach space of analytic functions over the unit disk \mathbb{D} . Let $\{X_n\}_{n \geq 0}$ be independent, identically distributed symmetric random variables over a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The symbol space \mathcal{X}_* for random \mathcal{X} -functions is another deterministic space, consisting of ‘symbols’ for random elements in \mathcal{X} :

$$\mathcal{X}_* = \{f \in H(\mathbb{D}) : \mathbb{P}(\mathcal{R}f \in \mathcal{X}) = 1\},$$



where $\mathcal{R}f(z) = \mathcal{R}^{\{(X_n)n\}}f(z)$ is given by $\mathcal{R}f(z) \doteq \sum_{n=0}^{\infty} a_n X_n z^n$, where $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D})$. By the Kolmogorov zero-one law, for any $f(z)$, $\mathbb{P}(\mathcal{R}f \in \mathcal{X}) \in \{0, 1\}$ under mild conditions. This justifies the introduction of \mathcal{X}_* .

The first nontrivial example of \mathcal{X}_* is perhaps the Littlewood theorem, reformulated as follows. Let $f(z) = a_0 + a_1 z + a_2 z^2 + \dots \in H^2(\mathbb{D})$ be an element of the Hardy space over the unit disk. Let $\{\epsilon_n\}_{n \geq 0}$ be a sequence of independent, identically distributed Bernoulli random variables, that is, $\mathbb{P}(\epsilon_n = 1) = \mathbb{P}(\epsilon_n = -1) = \frac{1}{2}$ for all $n \geq 0$. Littlewood’s theorem, proved in 1930 [29], states that $(\mathcal{R}f)(z) = \sum_{n=0}^{\infty} a_n \epsilon_n z^n \in H^p(\mathbb{D})$ almost surely for all $p \geq 0$. When $f \notin H^2(\mathbb{D})$, for almost every choice of signs, $\mathcal{R}f$ has a radial limit almost nowhere. This implies, in particular, $H^p(\mathbb{D})_* = H^2(\mathbb{D})$ for any $p > 0$. The same is true for a standard Steinhaus sequence [28, 36] and a standard Gaussian sequence ([22], p. 54). More generally, a Gaussian process version is obtained in [10].

When the random series $\mathcal{R}f$ represents an $H^\infty(\mathbb{D})$ -function almost surely is much harder, where $H^\infty(\mathbb{D})$ denotes the bounded analytic functions. Paley, Zygmund and Salem [35, 38] gave some necessary conditions and sufficient conditions. In [6], Billard showed that the Bernoulli case is equivalent to the Steinhaus case. A remarkable characterization was finally obtained by Marcus and Pisier in 1978 [31] (see also [22, 32]). Their characterization, or, in other words, a description of $H^\infty(\mathbb{D})_*$, builds on the celebrated Dudley–Fernique theorem. The space $H^\infty(\mathbb{D})_*$ can be equipped with a Banach space norm and lies strictly between the standard Dirichlet space and $H^2(\mathbb{D})$. Further understanding of $H^\infty(\mathbb{D})_*$ appears desirable.

In 1974, Anderson, Clunie and Pommerenke [2] studied the case of the Bloch space \mathcal{B} . Among other things, they showed that the condition $\sum_{n=1}^{\infty} |a_n|^2 \log n < \infty$ implies that $\sum_{n=0}^{\infty} e^{2\pi i \alpha_n} a_n z^n \in \mathcal{B}$ a.s. By Paley and Zygmund [35], this condition does not imply that $\sum_{n=0}^{\infty} a_n \epsilon_n z^n \in H^\infty(\mathbb{D})$ a.s. On the other hand, the condition $\sum_{n=1}^{\infty} |a_n|^2 (\log n)^{1+\epsilon} < \infty$ for some $\epsilon > 0$ implies that $\sum_{n=0}^{\infty} a_n \epsilon_n z^n \in \mathcal{A}$ a.s., where \mathcal{A} denotes the disk algebra. A necessary and sufficient condition for a random Taylor series to represent a Bloch function is given by Gao [18]. The space \mathcal{B}_* is less understood so far, and although not explicitly discussed in [18], it differs for the Gaussian and the Rademacher/Steinhaus randomization methods.

In 1981, Sledd [39] showed that the condition $\sum_{n=1}^{\infty} |a_n|^2 \log n < \infty$ implies that $\sum_{n=0}^{\infty} a_n \epsilon_n z^n \in \text{BMOA}$ a.s. Actually, he proved that it is indeed in the class of analytic functions of vanishing mean oscillation (VMOA). Then, some related results were extended by [42]. Sledd and Stegenga [40] showed that $\sum_{n=0}^{\infty} \epsilon_n a_n z^n \notin \text{BMOA}$ a.s. for some sequence $\{a_n\}_{n \geq 0} \in \ell^2$. In [16], Duren explored the difference between $\bigcap_{0 < p < \infty} H^p(\mathbb{D})$ and BMOA for the random Taylor series. Konyagin–Queffélec–Saksman–Seip provide one sufficient condition for a random Dirichlet series belonging to BMOA [24]. It remains an outstanding open problem to characterize BMOA_* and VMOA_* , which are different according to Nishry and Paquette [34].

Another elegant example is a theorem due to Cochran, Shapiro and Ullrich on the Dirichlet space [13]. They proved in 1993 that a Dirichlet function with random signs is a.s. a Dirichlet multiplier. Equivalently, $\mathcal{M}(\mathcal{D})_* = \mathcal{D}$. This result is generalized in [30]. Our preliminary investigation suggests that extending this result from $\mathcal{D} \doteq \mathcal{D}^2$ to \mathcal{D}^p , $p > 0$, leads to a new family of nontrivial Banach spaces.

On the other hand, for essentially every example of \mathcal{X}_* which we encounter, the monomials $\{z^n\}_{n \geq 0}$ exhibit features of an unconditional basis, yet they often don’t even

form a Schauder basis. When this happens, the space under consideration is usually non-separable. There is clearly a lack of proper technical tools in the current literature to describe such a phenomenon, and it prompts us to introduce the following definition, which is the focus of this study. Let $\mathfrak{S} = \{a = (a_1, a_2, \dots, a_n, \dots) : a_n \in \mathbb{C}, n \geq 1\}$ denote the vector space of sequences of complex scalars. Then, \mathfrak{S} is a locally convex vector space under the seminorms $p_n(a) = |a_n|, n \geq 1$. For simplicity, let $e_n \doteq (0, \dots, 0, 1, 0, \dots), n \geq 1$.

Definition 1. Let E be a Banach space, or a p -Banach space with $p \in (0, 1)$, and $\{e_n\}_{n \geq 1}$ a sequence of unit vectors in E . We say that $\{e_n\}_{n \geq 1}$ is a solid basis for E if there exists a linear map $\mathcal{T} : E \rightarrow \mathfrak{S}$, called the coefficient map, with $\mathcal{T}x = ((\mathcal{T}x)_1, \dots, (\mathcal{T}x)_n, \dots)$ and $\mathcal{T}e_n = e_n, n \geq 1$, such that

- (i) $\|\sum_{n=1}^N \lambda_n a_n e_n\|_E \leq \|\sum_{n=1}^N a_n e_n\|_E$ for $\lambda_n \in \mathbb{C}, |\lambda_n| \leq 1, N \in \mathbb{N}$ and $a_n \in \mathbb{C}$; and
- (ii) $\lim_{N \rightarrow \infty} \|\sum_{n=1}^N (\mathcal{T}x)_n e_n\|_E = \|x\|_E, x \in E$.

Remarks. (a) Recall that a functional $\|\cdot\| : E \rightarrow [0, \infty)$ is called a p -norm with $p \in (0, 1)$ if E is a complex vector space and $x, y \in E$, then

- (i) $\|x\| > 0$ if $x \neq 0$, and $\|\lambda x\| = |\lambda| \|x\|, \lambda \in \mathbb{C}$; and
- (ii) $\|x + y\|^p \leq \|x\|^p + \|y\|^p$.

If (E, d) , with $d(x, y) = \|x - y\|^p$, is a complete metric space, then it is called a p -Banach space. Banach spaces are p -Banach spaces for each $p \in (0, 1)$, although we don't need such an identification in this paper.

(b) For convenience, let $P_N(x) = \sum_{n=1}^N (\mathcal{T}x)_n e_n, x \in E$, which we view as an abstract Taylor polynomial.

(c) By Definition 1, the quantity $\|P_N(x) - x\|_E$ is necessarily decreasing in N ; hence, the limit $\text{gap}(x) \doteq \lim_{N \rightarrow \infty} \|P_N(x) - x\|_E$ exists. Those elements such that $\text{gap}(x) = 0$ form a closed subspace of E . The situation when $\text{gap}(x) > 0$ is the more interesting part of our study, and this happens usually when E is non-separable.

In order to quantify the clause ‘essentially every example of \mathcal{X}_* ’ above and to put our study on a proper footing, we are led to revisit the axioms satisfied by canonical Banach spaces of analytic functions over the unit disk. This is a subject with a long history. The first systematic investigation is probably due to Taylor in 1951 [41]. Two subsequent developments, somehow similar to our study in spirit, are due to Brown and Shields [9], in which the authors proposed a set of axioms suitable for their research of cyclic vectors, and due to Blasco and Pavlović [7], in which they presented a framework suitable for the study of coefficient multipliers. In this paper, we present another set of axioms (Definition 2), which are suitable not only for our study of random analytic functions but also of interests for the general study of Banach spaces of analytic functions. Compared with earlier works, a feature of our axioms is that they lead to several functorial constructions.

For any (p -)Banach space \mathcal{X} of analytic functions over the unit disk \mathbb{D} , let δ_z denote the point evaluation functional: $\delta_z(f) = f(z), f \in \mathcal{X}, z \in \mathbb{D}$ and let $f_w(z) = f(wz)$ for

$z \in \mathbb{D}$ and $w \in \overline{\mathbb{D}}$. Our axioms for canonical Banach spaces of analytic functions are the following:

Definition 2. A functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, \mathcal{X} of analytic functions over \mathbb{D} , which contains all the polynomials, is called of homogeneous type if

- (i) the point evaluations are continuous: $\{\delta_z : z \in \mathbb{D}\} \subset \mathcal{X}^*$;
- (ii) the rotations preserve the norm: $\|f_\xi\|_{\mathcal{X}} = \|f\|_{\mathcal{X}}$ for $f \in \mathcal{X}$ and $\xi \in \mathbb{T}$;
- (iii) the dilation is continuous: The function $F = F_f(\cdot) : \mathbb{D} \rightarrow \mathcal{X}$, defined by $F(z) = f_z$ for $f \in \mathcal{X}$, is well-defined and continuous; and
- (iv) the norm is increasingly continuous: $\sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} = \|f\|_{\mathcal{X}}$ for $f \in \mathcal{X}$.

Each of the above four axioms is clearly familiar to a practitioner in the field, and they are inclusive enough to accommodate every example of interests to us. The main point, which we stress again, is that they turn out to suit our study of \mathcal{X}_* well. Note that the function $r \mapsto \|f_r\|_{\mathcal{X}}$ is necessarily continuous and increasing over $(0, 1)$ which follows from (iii) and (iv) in Definition 2. In this paper, we usually use E to denote an abstract (p -)Banach space and \mathcal{X} a functional (p -)Banach space. Moreover, let $S_N f$ denote the degree- N Taylor polynomial for $f \in H(\mathbb{D})$ ($N \geq 0$).

Measurability assumption. Several results in this paper involve not only \mathcal{X} above, but also \mathcal{X}_* , its symbol space. In order to make a statement about \mathcal{X}_* , we still need to impose another very weak but necessary assumption: For any $f \in H(\mathbb{D})$, the event $\{\mathcal{R}f \in \mathcal{X}\}$ is Borel measurable, so that $\mathbb{P}(\mathcal{R}f \in \mathcal{X})$ is defined. The Kolmogorov zero-one law implies, then, $\mathbb{P}(\mathcal{R}f \in \mathcal{X}) \in \{0, 1\}$, and hence, \mathcal{X}_* is well defined. Subsequently, in this paper, whenever we mention \mathcal{X}_* , we assume that this measurability assumption is in force. On the other hand, once \mathcal{X}_* is defined, this assumption will no longer enter any argument.

The following is our principal example of solid bases.

Theorem A. Let $\{X_n\}_{n \geq 0}$ be a standard random sequence (Definition 6).

- (i) Let \mathcal{X} be a functional Banach space of homogeneous type. Then
 - (i.1) \mathcal{X}_* is a functional Banach space of homogeneous type under the norm $\|f\|_{\mathcal{X}_*} = \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}$, which is necessarily finite.
 - (i.2) If $\{X_n\}_{n \geq 0}$ is a standard Steinhaus or complex Gaussian sequence, then $\left\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\right\}_{n \geq 0}$ is a solid basis for \mathcal{X}_* .
 - (i.3) If $\{X_n\}_{n \geq 0}$ is a standard Rademacher or real Gaussian sequence, then $\left\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\right\}_{n \geq 0}$ is a solid frame for \mathcal{X}_* . Moreover, $\|f\|_{\mathcal{X}_*} = \lim_{N \rightarrow \infty} \|S_N f\|_{\mathcal{X}_*}$ for $f \in \mathcal{X}_*$.
- (ii) Let \mathcal{X} be a functional p -Banach space of homogeneous type with $p \in (0, 1)$. Assume that $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. Then
 - (ii.1) \mathcal{X}_* is a functional p -Banach space of homogeneous type under the p -norm $\|f\|_{\mathcal{X}_*} = (\mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}^p)^{1/p}$, which is necessarily finite.
 - (ii.2) $\left\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\right\}_{n \geq 0}$ is a solid frame for \mathcal{X}_* .

Definition 3. A sequence of unit vectors $\{e_n\}_{n \geq 1}$ is said to be a solid frame for E , a Banach or p -Banach space, if there exists a linear coefficient map $\mathcal{T} : E \rightarrow \mathfrak{S}$ with $\mathcal{T}x = ((\mathcal{T}x)_1, \dots, (\mathcal{T}x)_n, \dots)$ and $\mathcal{T}e_n = e_n, n \geq 1$, such that

- (i) there exists a constant C such that $\|\sum_{n=1}^N \lambda_n a_n e_n\|_E \leq C \|\sum_{n=1}^N a_n e_n\|_E$ for any $\lambda_n \in \mathbb{C}, |\lambda_n| \leq 1$, and $N \in \mathbb{N}, a_n \in \mathbb{C}$; and
- (ii) there exist C' and C'' such that $C'\|x\|_E \leq \sup_{N \geq 1} \|P_N(x)\|_E \leq C''\|x\|_E, x \in E$.

Theorem E below explains how to transform solid frames into solid bases.

Remarks. (a) By the contraction principle ([21, Theorem 6.1.13, p. 9]) and Lemma 8, we can choose $C = \frac{\pi}{2}, C' = C'' = 1$ and $C' = 1, C'' = 2^{\frac{1}{p}-1}$, respectively, for the constants of solid frames in (i.3) and (ii.2), Theorem A.

(b) From the assumption $\{\delta_z : z \in \mathbb{D}\} \subset \mathcal{X}^*$, one can show that $\liminf_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} \geq 1$.

Functorial constructions. Next, we summarize various functorial constructions for \mathcal{X} and E , which are features following from our choice of axioms for a typical (p-)Banach space of analytic functions. Although they are motivated by the study of random analytic functions, these constructions are clearly of independent interests for the general study of functional (p-)Banach spaces.

- (a) \mathcal{X}^{\min} and \mathcal{X}^{\max} : We shall introduce a notion called *homogeneous extension* (Definition 4), and in this sense, for any homogeneous \mathcal{X} , we show that there exists a unique \mathcal{X}^{\min} and a unique \mathcal{X}^{\max} such that

$$\mathcal{X}^{\min} \subset \mathcal{X} \subset \mathcal{X}^{\max}.$$

Examples of the pair $(\mathcal{X}^{\min}, \mathcal{X}^{\max})$ include

- $(A(\mathbb{D}), H^\infty(\mathbb{D}))$, the disk algebra and bounded analytic functions
- $(\mathcal{B}_0, \mathcal{B})$, the little Bloch and Bloch spaces, and
- $(VMOA, BMOA)$.

These notions appear in Theorem B, Theorem F and Lemma 10.

- (b) $E_{\mathcal{P}}$ and $E^{s-\max}$: We shall introduce a notion called *solid extension* (Definition 5). This allows us to introduce sequential analogs of \mathcal{X}^{\min} and \mathcal{X}^{\max} for E , denoted by $E_{\mathcal{P}}$ and $E^{s-\max}$, respectively. Then, for any E with a solid basis $\{e_n\}_{n \geq 1}$, $E_{\mathcal{P}}$ is simply given by $\overline{\text{span}}\{e_n : n \geq 1\}$ and $E^{s-\max}$ by (7). Examples of the pair $(E_{\mathcal{P}}, E^{s-\max})$ include
 - (c_0, ℓ^∞) , and
 - $((H^\infty(\mathbb{D}))_*, (H^\infty(\mathbb{D}))_*)$.

These notions appear in Theorem H, Theorem I, Proposition 14, Lemma 16 and Proposition 17.

- (c) $(\mathcal{X}, \|\cdot\|_{\#})$: The construction

$$\|f\|_{\#} = \sup_{N \geq 0} \sup_{\|\lambda\|_{\ell^\infty} \leq 1} \left\| \sum_{k=0}^N \lambda_k a_k z^k \right\|_{\mathcal{X}}, \tag{1}$$

where $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}$, allows us to transform a solid frame into a solid basis; see Theorem E.

- (d) A pseudo-canonical embedding $\mathfrak{J} : E \rightarrow E^{**}$: Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. Let $J : E \rightarrow E^{**}$ be the canonical embedding map into its second dual. We shall establish the weak-star convergence of $J(P_N x)$ as $N \rightarrow \infty$, leading to what we call a pseudo-canonical map $\mathfrak{J} : E \rightarrow E^{**}$, which stands in interesting contrast to the map J ; see Theorem D. In particular, $Jx = \mathfrak{J}x$ if and only if $x \in E_{\mathcal{P}}$. A related map $\Phi : E^{\text{s-max}} \rightarrow (E_{\mathcal{P}})^{**}$ is constructed in order to see how large $E^{\text{s-max}}$ usually is. Two examples of $E^{\text{s-max}}$ are $E = c_0$, $E^{\text{s-max}} = E^{**} = \ell^{\infty}$ and

$$E = \ell^{\infty}, E^{\text{s-max}} = \ell^{\infty} \subset (\ell^{\infty})^{**} = E^{**}.$$

To get a grasp of $E^{\text{s-max}}$ in general, we construct an embedding map $\Phi : E^{\text{s-max}} \rightarrow (E_{\mathcal{P}})^{**}$, which fixes $E_{\mathcal{P}}$ (under the natural identification).

Precise statements of results related to the above constructions are presented next.

Definition 4. Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type. We say that \mathcal{Y} is a homogeneous extension of \mathcal{X} if

- (i) \mathcal{Y} is a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type;
- (ii) $\mathcal{X} \subset \mathcal{Y}$, and $f_r \in \mathcal{X}$ if $0 \leq r < 1$, $f \in \mathcal{Y}$; and
- (iii) $\|f\|_{\mathcal{Y}} = \|f\|_{\mathcal{X}}$ for any $f \in \mathcal{X}$.

\mathcal{X}^{min} and \mathcal{X}^{max} . For any \mathcal{X} as above, we set $\mathcal{X}^{\text{min}} = \mathcal{X}_{\mathcal{P}}$, the closure of polynomials in \mathcal{X} . Two facts need to be verified and both are straightforward:

- $\mathcal{X}_{\mathcal{P}}$ is of homogeneous type, and
- \mathcal{X} is a homogeneous extension of $\mathcal{X}_{\mathcal{P}}$.

For \mathcal{X}^{max} , we prove

Theorem B. Let $\{X_n\}_{n \geq 0}$ be a standard random sequence (Definition 6). Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous

type; for the latter, we also assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. Then,

- (i) there exists a unique maximal homogeneous extension, denoted by \mathcal{X}^{max} ; and
- (ii) both $(\mathcal{X}^{\text{max}})_{\star}$ and $(\mathcal{X}_{\mathcal{P}})_{\star}$ are solid spaces.

A solid space, introduced by Anderson and Shields [3], is a sequence space E such that $\{b_n\}_{n \geq 1} \in E$ whenever $\{a_n\}_{n \geq 1} \in E$ and $|b_n| \leq |a_n|$. Although a solid space is not a space with a solid basis, our Definition 1 is clearly motivated by it. Note that, in this paper, a function $f \in H(\mathbb{D})$ is often identified with its sequence of Taylor coefficients. In general, \mathcal{X}_{\star} may fail to be solid; see the example after the proof of Theorem B in Section 4.

Fortunately, Theorem B is quite applicable, since, indeed, most natural examples of \mathcal{X} are of the forms: \mathcal{X}^{\max} or $\mathcal{X}_{\mathcal{P}}$.

Remark. When $\{X_n\}_{n \geq 0}$ is a standard random sequence (Definition 6), it is known that $(H^\infty(\mathbb{D}))_*$ is solid. This follows from the celebrated characterization of $(H^\infty(\mathbb{D}))_*$ in terms of the modulus of the Taylor coefficients of its elements [32]. Theorem B implies that

$$\text{BMOA}_*, \text{VMOA}_*, \mathcal{B}_*, \text{ and } (\mathcal{B}_0)_*$$

are solid as well, and this is not obvious. It is of value to characterize when \mathcal{X}_* is solid, and we present it as Problem A in order to spur more interests.

Unconditional bases. Next, we discuss the relationship between solid bases and unconditional bases. A Schauder basis $\{e_n\}_{n \geq 1}$ in a Banach space or a p -Banach space E is called an *unconditional basis* if for every $x \in E$, there exists a unique sequence of scalars $\{a_n(x)\}_{n \geq 1}$ such that $x = \sum_{n=1}^\infty a_n(x)e_n$, and the series $\sum_{n=1}^\infty a_n(x)e_n$ converges unconditionally, that is, $\sum_{n=1}^\infty a_{\pi(n)}(x)e_{\pi(n)}$ converges for every permutation π of positive integers. Equivalently, for any choice of phases $\theta = (\theta_k)_{k \geq 1} \in \mathbb{T}^{\mathbb{N}}$, the symmetries $M_\theta : \sum_{n=1}^\infty a_n(x)e_n \mapsto \sum_{n=1}^\infty \theta_n a_n(x)e_n$ are continuous on E and $\sup_\theta \|M_\theta\| < \infty$; see [25, Proposition III.3, p. 91], [27, p. 18] or [1, Theorem 1.10]. The quantity $K = \sup_\theta \|M_\theta\|$ is called the unconditional constant of the basis $\{e_n\}_{n \geq 1}$.

Theorem C.

(i) Let \mathcal{X} be a functional Banach space of homogeneous type. Then, the following are equivalent:

- (i.1) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid basis for \mathcal{X} ;
- (i.2) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid basis for $\mathcal{X}_{\mathcal{P}}$;
- (i.3) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is an unconditional basis for $\mathcal{X}_{\mathcal{P}}$ with constant 1;
- (i.4) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid basis for \mathcal{X}^{\max} .

(ii) Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter we assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. Then, the following are equivalent:

- (ii.1) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid frame for \mathcal{X} ;
- (ii.2) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid frame for $\mathcal{X}_{\mathcal{P}}$;
- (ii.3) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is an unconditional basis for $\mathcal{X}_{\mathcal{P}}$;
- (ii.4) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid frame for \mathcal{X}^{\max} .

For an abstract Banach space E , the relationship between solid bases/frames and unconditional bases is presented in Lemma 16.

Weak-star convergence of Taylor polynomials. Next, we claim that, in the presence of a solid basis, partial summations, viewed as abstract Taylor polynomials, are weak-star sequentially convergent in the double dual. This stands in interesting contrast to a typical existing result, say, Proposition 1 in [43], which concerns the norm

convergence and where the Banach space is separable; hence, the interest in Theorem D below lies in the non-separable case. Also note that, under mild conditions, weak convergence and norm convergence of Taylor polynomials are equivalent – see the discussion at the beginning of Section 6 – but weak-star sequential convergence results are indeed rare. Let $J : E \rightarrow E^{**}$ denote the canonical imbedding of a Banach space into its second dual.

Theorem D. *Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. For each $x \in E$, $J(P_N x)$ converges in w^* -topology as $N \rightarrow \infty$.*

The Banach–Alaoglu theorem implies that $\{P_N x\}_{N \geq 1}$ has a w^* -convergent subnet since $\sup_{N \geq 1} \|P_N x\|_E \leq \|x\|_E < \infty$. A subtle point here is that the subnet is not necessarily a subsequence. Our contribution, hence, is to boost to the sequential convergence. Let $\mathfrak{J}(x)$ denote the limit in Theorem D. It is interesting to gain more insight, in comparison with the canonical embedding J , about the map $\mathfrak{J} : E \rightarrow E^{**}$. In particular, we shall show that ‘ $\mathfrak{J}(x) = J(x)$ if and only if $x \in E_{\mathcal{P}}$ ’; hence, the two maps appear to be considerably different when E is non-separable. The existence of $\mathfrak{J}x$ depends only on (8), and it can be modified to yield an embedding $\Phi : E^{s\text{-max}} \rightarrow (E_{\mathcal{P}})^{**}$. This Φ may be viewed as an abstract extension of some classical embedding maps.

Remark. The motivation for Theorem D comes from the so-called w.u.C. condition in Banach space theory; see, in particular, [15, p. 44] or [25, Proposition II.5, p. 87]. Given a Banach space E , a (formal) series $\sum_{n=1}^{\infty} x_n$ is called *weakly unconditionally Cauchy* if $\sum_{n=1}^{\infty} |\phi(x_n)| < \infty$ for every $\phi \in E^*$.

From frames to bases. Next, we show how to transform a solid frame into a solid basis, with the help of the norm $\|\cdot\|_{\sharp}$, which is motivated by [33, Definitions 4.1.13, 4.2.15].

Theorem E. *Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter, we also assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$.*

For each $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}$, let $\|f\|_{\sharp} = \sup_{N \geq 0} \sup_{\|\lambda\|_{\ell^{\infty}} \leq 1} \left\| \sum_{k=0}^N \lambda_k a_k z^k \right\|_{\mathcal{X}}$. Then, the following are equivalent:

- (i) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid frame for \mathcal{X} .
- (ii) $(\mathcal{X}, \|\cdot\|_{\sharp})$ is a Banach space (or p -Banach space), and $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid basis for $(\mathcal{X}, \|\cdot\|_{\sharp})$.

A large class of examples. Next, we present a characterization of solid frames in a case, which covers a large class of natural examples.

Theorem F. *Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter, we also assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$.*

If $\mathcal{X} = \mathcal{X}^{\text{max}}$ or $\mathcal{X} = \mathcal{X}_p$, then $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid frame for \mathcal{X} if and only if $\sup_{N \geq 0} \|S_N\| < \infty$ and \mathcal{X} is a solid space. Here, $\|S_N\|$ is the operator norm.

Remark. Since $(\mathcal{X}_*)_{\mathcal{P}} = (\mathcal{X}_{\mathcal{P}})_*$ and $(\mathcal{X}_*)^{\max} = (\mathcal{X}^{\max})_*$ (Lemma 10), the above result should be compared with (ii) in Theorem B.

Solid span. A sequence $\{e_n\}_{n \geq 1}$ in a Banach space E is said to be a *basic sequence* if it is a Schauder basis for the closure of its span, i.e., $\overline{\text{span}\{e_n : n \geq 1\}}$. For a new notion of Banach space basis, such as the one we propose in this paper, a natural question to ask is: How to construct the solid span of a basic sequence? This does not have an easy answer since c_0 and ℓ^∞ show that we can find two Banach spaces $E \subsetneq F$ such that $\{e_n\}_{n \geq 1}$ is a solid basis for both. This discrepancy can be nicely resolved by introducing the following:

Definition 5. Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. A Banach space F is called a *solid extension* of E if

- (i) there exists an isometric embedding $\iota : E \hookrightarrow F$; and
- (ii) $\{\iota e_n\}_{n \geq 1}$ is also a solid basis for F .

Theorem G. Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$.

- (i) There exists a unique maximal solid extension under the isometric isomorphism, denoted by $E^{\text{s-max}}$ (which is necessarily given by (7)).
- (ii) Let W be a Banach space. Assume that $\iota : E_{\mathcal{P}} \hookrightarrow W$ is an isometric embedding. Then $\{\iota e_n\}_{n \geq 1}$ is a solid basis for W if and only if W can be embedded into $(E_{\mathcal{P}})^{\text{s-max}}$ isometrically.

Remark. This theorem may be viewed as an abstract extension of Theorem C for \mathcal{X} .

The role of c_0 . At last, we highlight the curious roles played by c_0 in our study. This part is motivated by the corresponding action of c_0 in the theory of unconditional bases; see, for example, [25, Chapter 3]. Recall that c_0 is the subspace of \mathfrak{S} consisting of sequences convergent to 0. Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. Let $c_0 \cdot E \doteq \left\{ \sum_{n=1}^{\infty} t_n (\mathcal{T}x)_n e_n : (t_n)_{n \geq 1} \in c_0, x \in E \right\}$. Similarly, for $a = (a_1, a_2, \dots) \in \mathfrak{S}$, let $c_0 \cdot a = \{(a_n t_n)_{n \geq 1} : (t_n)_{n \geq 1} \in c_0\}$. The proofs of the following two theorems involve the dual space E^* , hence confined to Banach spaces.

Theorem H. Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. Then,

- (i) $c_0 \cdot E = E_{\mathcal{P}}$.
- (ii) For each $a \in \mathfrak{S}$, $c_0 \cdot a \in E_{\mathcal{P}}$ if and only if $a \in E^{\text{s-max}}$.

Theorem I. Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. Then, $\{e_n\}_{n \geq 1}$ is an unconditional basis for $E^{\text{s-max}}$ if and only if $E^{\text{s-max}}$ does not contain c_0 .

The proof of the last result is a quick application of one of the famous James' theorems [25, Theorem V.4, p. 98].

2. Preliminary issues

For the reader’s convenience, in this section, we collect standard definitions and technical tools which are used repeatedly in this paper. Let $H(\mathbb{D})$ denote the collection of all analytic functions over the unit disk \mathbb{D} in the complex plane, and we often identify $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D})$ with its sequence of Taylor coefficients $\{a_n\}_{n \geq 0}$. For $0 < p < \infty$, $H^p(\mathbb{D})$ [17] denotes the Hardy space, consisting of $f \in H(\mathbb{D})$ such that

$$\|f\|_{H^p(\mathbb{D})} = \sup_{0 \leq r < 1} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p} < \infty.$$

The Bergman space $L_a^p(\mathbb{D})$ [19] consists of $f \in H(\mathbb{D})$ such that

$$\|f\|_{L_a^p(\mathbb{D})} = \left(\int_{\mathbb{D}} |f(z)|^p dA(z) \right)^{1/p} < \infty,$$

where $dA = \frac{1}{\pi} dx dy$. To be convenient, we shall write H^p for $H^p(\mathbb{D})$, H^∞ for $H^\infty(\mathbb{D})$, the algebra of bounded analytic functions on \mathbb{D} , and L_a^p for $L_a^p(\mathbb{D})$. Let \mathcal{D}^p denote the Dirichlet space, consisting of $f \in H(\mathbb{D})$ such that $f' \in L_a^p$. Let $\mathcal{M}(\mathcal{D}^p)$ denote its multiplier algebra, that is, $h \in \mathcal{M}(\mathcal{D}^p)$ if and only if $hf \in \mathcal{D}^p$ for each $f \in \mathcal{D}^p$. For simplicity, \mathcal{D}^2 is shorthand as \mathcal{D} . The Bloch space \mathcal{B} [2] consists of $f \in H(\mathbb{D})$ such that $\sup_{z \in \mathbb{D}} (1 - |z|^2)|f'(z)| < \infty$ and the little Bloch space \mathcal{B}_0 the closure of polynomials in \mathcal{B} .

Definition 6. A random variable X is called Rademacher if $\mathbb{P}(X = 1) = \mathbb{P}(X = -1) = \frac{1}{2}$, Steinhaus if it is uniformly distributed on the unit circle, standard real Gaussian if its density is $\frac{1}{\sqrt{2\pi}} \exp(-t^2/2)$ with $t \in \mathbb{R}$ and standard complex Gaussian if its density is $\frac{1}{\pi} \exp(-|z|^2)$ with $z \in \mathbb{C}$. Moreover, let

$$X \in \{\text{Rademacher, Steinhaus, standard real Gaussian or complex Gaussian}\}.$$

Then, by a standard X -sequence, we mean a sequence of independent, identically distributed X variables. Lastly, a standard random sequence $\{X_n\}_{n \geq 0}$ refers to either a Rademacher, Steinhaus, standard real or complex Gaussian sequence, denoted by

$$\{\epsilon_n\}_{n \geq 0}, \quad \{e^{2\pi i \alpha_n}\}_{n \geq 0}, \quad \{\xi_n\}_{n \geq 0} \quad \text{and} \quad \{\gamma_n\}_{n \geq 0},$$

respectively.

We assume that all random variables are defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with expectation denoted by $\mathbb{E}(\cdot)$, where \mathcal{F} is a σ -algebra on Ω and \mathbb{P} is a probability measure defined on \mathcal{F} . The probability of an event $\{\omega \in \Omega : X(\omega) \in A\}$, with $A \in \mathcal{F}$, is $\mathbb{P}(X \in A)$. The abbreviation ‘a.s.’ stands for ‘almost surely’. As usual, ‘ $A \approx B$ ’ implies that there exists a positive constant $C < \infty$ such that $\frac{A}{C} \leq B \leq CA$, ‘ $A \lesssim B$ ’ (resp. ‘ $A \gtrsim B$ ’) means that there exists a positive constant $C < \infty$ such that $A \leq CB$ (resp. $A \geq CB$), and ‘ \Leftrightarrow ’ stands for ‘if and only if’.

3. The symbol space \mathcal{X}_\star

In this section, we prove Theorem A.

Proof of Theorem A. The proof of (i.1) is divided into three steps.

Step 1. $f \in \mathcal{X}_\star \implies \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}} < \infty$.

By (iv) of Definition 2, $\sup_{0 \leq r < 1} \|(\mathcal{R}f)_r\|_{\mathcal{X}} = \|\mathcal{R}f\|_{\mathcal{X}} < \infty$ a.s. Then, the Marcinkiewicz–Zygmund–Kahane theorem [26, Theorem II.4, p. 240] implies that $\sup_{N \geq 0} \|S_N(\mathcal{R}f)\|_{\mathcal{X}} < \infty$ a.s. since $\mathcal{R}f$ is A -bounded in \mathcal{X} with $A = (r_n^m)_{n,m \in \mathbb{N}}$ and $r_n \rightarrow 1^-$ as $n \rightarrow \infty$. Let $M = \sup_{N \geq 0} \|S_N(\mathcal{R}f)\|_{\mathcal{X}}$. Then, by Kahane’s inequality [25, Theorem V.1, p. 139] (for Rademacher or Steinhaus sequences) and Fernique’s theorem [25, Theorem V.26, p. 255] (for Gaussian sequences), we have $\mathbb{E}(\exp(\lambda M)) < \infty$ for some small enough $\lambda > 0$. Now, we need the following lemma, which is a consequence of [4, Theorem 3.1]. \square

Lemma 7. *Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter, we also assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. If $f \in \mathcal{X}$, then for each fixed $0 \leq r < 1$, $\lim_{N \rightarrow \infty} \|S_N f_r - f_r\|_{\mathcal{X}} = 0$.*

By Lemma 7 and (iv) of Definition 2, $\|\mathcal{R}f\|_{\mathcal{X}} \leq M$ a.s. Now, Step 1 follows from Kahane’s inequality [25, Theorem V.1, p. 139] and Fernique’s theorem [25, Theorem V.26, p. 255].

Step 2. $(\mathcal{X}_\star, \|\cdot\|_{\mathcal{X}_\star})$ is a Banach space.

Let $\{f_n\}_{n \geq 1}$ be a Cauchy sequence in \mathcal{X}_\star with $f_n(z) = \sum_{k=0}^{\infty} a_k^{(n)} z^k$. Then, there exists $F(z, \omega) = \sum_{k=0}^{\infty} c_k(\omega) z^k \in L^1(\Omega; \mathcal{X})$ such that $\lim_{n \rightarrow \infty} \mathbb{E}\|\mathcal{R}f_n - F\|_{\mathcal{X}} \rightarrow 0$. By the Riesz lemma, there exists a subsequence $\{f_{n_i}\}_{i \geq 1}$ and $\Omega_0 \subset \Omega$ with $\mathbb{P}(\Omega_0) = 1$, such that for $\omega \in \Omega_0$,

$$\lim_{i \rightarrow \infty} \|\mathcal{R}f_{n_i}(z, \omega) - F(z, \omega)\|_{\mathcal{X}} = \lim_{i \rightarrow \infty} \left\| \sum_{k=0}^{\infty} a_k^{(n_i)} X_k(\omega) z^k - \sum_{k=0}^{\infty} c_k(\omega) z^k \right\|_{\mathcal{X}} = 0. \tag{2}$$

Since $\{\delta_z : z \in \mathbb{D}\} \subset \mathcal{X}^*$, for each fixed k and $\omega \in \Omega_0$, we have $\lim_{i \rightarrow \infty} a_k^{(n_i)} X_k(\omega) = c_k(\omega)$. Consequently, there exists a constant a_k such that $c_k(\omega) = a_k X_k(\omega)$, $\omega \in \Omega_0$. Now, (2) implies that $\mathcal{R}f_{n_i} \rightarrow \mathcal{R}f$ a.s. with $\mathcal{R}f(z) = \sum_{k=0}^{\infty} a_k X_k z^k \in \mathcal{X}_\star$. Then, for any $\epsilon > 0$,

$$\begin{aligned} \mathbb{E}\|\mathcal{R}f_{n_i} - \mathcal{R}f\|_{\mathcal{X}} &\leq \lim_{r \rightarrow 1^-} \liminf_{j \rightarrow \infty} \mathbb{E}\|(\mathcal{R}f_{n_i})_r - (\mathcal{R}f_{n_j})_r\|_{\mathcal{X}} \\ &\leq \sup_{j \geq N} \mathbb{E}\|\mathcal{R}f_{n_i} - \mathcal{R}f_{n_j}\|_{\mathcal{X}} \\ &< \epsilon \end{aligned}$$

when i and N are large enough. Now, $\mathbb{E}\|\mathcal{R}f_n - \mathcal{R}f\|_{\mathcal{X}} \rightarrow 0$ as $n \rightarrow \infty$ by the triangle inequality.

Step 3. $(\mathcal{X}_*, \|\cdot\|_{\mathcal{X}_*})$ is of homogeneous type.

Firstly, let $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}_*$. Then, there exists constant $C > 0$ such that

$$|f(z)| \leq \sum_{n=0}^{\infty} |a_n| |z|^n \leq C \sum_{n=0}^{\infty} \frac{\|f\|_{\mathcal{X}_*}}{\|z^n\|_{\mathcal{X}}} |z|^n = C \left(\sum_{n=0}^{\infty} \frac{|z|^n}{\|z^n\|_{\mathcal{X}}} \right) \|f\|_{\mathcal{X}_*},$$

where the second inequality follows from the contraction principle ([21, Theorem 6.1.13, p. 9]) and Lemma 8 below, i.e., $\|a_n z^n\|_{\mathcal{X}_*} \leq C \|S_N f\|_{\mathcal{X}_*} \leq C \|f\|_{\mathcal{X}_*}$ if $N > n$. Since $\{\delta_z : z \in \mathbb{D}\} \subset \mathcal{X}^*$, one has $\liminf_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{1/n} \geq 1$, which yields $\sum_{n=0}^{\infty} \frac{|z|^n}{\|z^n\|_{\mathcal{X}}} < \infty$ for fixed $z \in \mathbb{D}$. So $\{\delta_z : z \in \mathbb{D}\} \subset (\mathcal{X}_*)^*$. Secondly,

$$\|f_{\xi}\|_{\mathcal{X}_*} = \mathbb{E}\|(\mathcal{R}f)_{\xi}\|_{\mathcal{X}} = \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}} = \|f\|_{\mathcal{X}_*}, \quad \xi \in \mathbb{T}.$$

Thirdly, the continuity of F follows from [4, Theorem 3.1]. Lastly, for $f \in \mathcal{X}_*$,

$$\|f\|_{\mathcal{X}_*} = \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}} = \mathbb{E}\left(\sup_{0 \leq r < 1} \|\mathcal{R}f_r\|_{\mathcal{X}}\right) = \sup_{0 \leq r < 1} \mathbb{E}\|\mathcal{R}f_r\|_{\mathcal{X}} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}_*}.$$

Now, we turn to (i.2). If $\{X_n\}_{n \geq 0}$ is a standard Steinhaus or complex Gaussian sequence, then by the contraction principle ([21, Theorem 6.1.13, p. 9]),

$$\left\| \sum_{n=0}^N \lambda_n a_n \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\|_{\mathcal{X}^*} \leq \mathbb{E} \left\| \sum_{n=0}^N a_n X_n \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\|_{\mathcal{X}}$$

for $|\lambda_n| \leq 1, N \in \mathbb{N}$ and $a_n \in \mathbb{C}$. Take $\mathcal{T} : \mathcal{X}_* \rightarrow \mathfrak{S}$ as $f(z) = \sum_{n=0}^{\infty} a_n z^n \mapsto (a_n \|z^n\|_{\mathcal{X}})_{n \geq 0}$.

Then $P_N f(z) = \sum_{n=0}^N (\mathcal{T}f)_n \frac{z^n}{\|z^n\|_{\mathcal{X}}} = S_N f(z)$. It suffices to show that $\|S_N f\|_{\mathcal{X}_*} \rightarrow \|f\|_{\mathcal{X}_*}$, as $N \rightarrow \infty$, for $f \in \mathcal{X}_*$, which follows from (3) in Lemma 8 below.

Now, (i.3) for a standard Rademacher or real Gaussian sequence, which are real symmetric, follows from the contraction principle ([21, Theorem 6.1.13, p. 9]) just as case (i.2). The details are skipped.

The proof of (ii) is similar to that of (i); hence, we only indicate the differences briefly. For $f \in \mathcal{X}_* \implies \mathbb{E}(\|\mathcal{R}f\|_{\mathcal{X}}^p) < \infty$, the Kahane’s inequality and Fernique’s theorem used in the proof of Theorem A (Step 1) still hold for p -Banach spaces; see [11, Theorem 28 and Theorem 31]. The other parts of the proof can be modified accordingly. That $(\mathcal{X}_*, \|\cdot\|_{\mathcal{X}_*})$ is a functional p -Banach space can be proved by replacing $\mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}$ with $\mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}^p$. To check that $(\mathcal{X}_*, \|\cdot\|_{\mathcal{X}_*})$ is of homogeneous type, the only difference is that the continuity of $F : \mathbb{D} \rightarrow \mathcal{X}_*$, defined by $F(z) = f_z$, is a consequence of the assumption $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. Then, (ii.2) follows from the contraction principle for p -Banach spaces (see [23, Proposition 2.5]) and Lemma 8.

Remark. If $\mathcal{X} = \mathcal{X}_p$ for a p -Banach space, the assumption $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$ in Theorem A (ii) can be dropped. In detail, by the claim in the proof of Lemma 10,

$f \in (\mathcal{X}_p)_*$ implies that $(\mathcal{R}f)_r$ converges to $\mathcal{R}f$ a.s. as $r \rightarrow 1^-$. Then, by the Marcinkiewicz–Zygmund–Kahane theorem [26, Theorem II.4, p. 240], $S_N(\mathcal{R}f)$ converges to $\mathcal{R}f$ a.s. as $N \rightarrow \infty$. $S_N f$ converges to f in $(\mathcal{X}_p)_*$ as $N \rightarrow \infty$ by Kahane’s inequality [25, Theorem V.1, p. 139] and Fernique’s theorem [25, Theorem V.26, p. 255]. The rest of the proof is similar to the proof of part (i) of Theorem A. The same remark applies to part (i) of Lemma 9.

Remark. The role of M , appearing in Step 1 of the proof of (i), deserves further attention in the study of random analytic functions. We introduce a deterministic version

$$\mathcal{X}_T = \left\{ f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D}) : \|f\|_{\mathcal{X}_T} = \sup_{N \geq 0} \left\| \sum_{n=0}^N a_n z^n \right\|_{\mathcal{X}} < \infty \right\}.$$

Then, $(\mathcal{X}_T)_*$ is a solid space by Proposition 11 and \mathcal{X}^{\max} is identified as \mathcal{X}_T under certain conditions in Proposition 15.

We end this section with a supplemental property about the norm $\|f\|_{\mathcal{X}_*}$ in Theorem A.

Lemma 8. *Let $\{X_n\}_{n \geq 0}$ be a standard random sequence.*

(i) *If \mathcal{X} is a functional Banach space of homogeneous type and $f \in \mathcal{X}_*$, then*

$$\|f\|_{\mathcal{X}_*} = \sup_{N \geq 0} \|S_N f\|_{\mathcal{X}_*} = \lim_{N \rightarrow \infty} \|S_N f\|_{\mathcal{X}_*}. \tag{3}$$

(ii) *If \mathcal{X} is a functional p -Banach space of homogeneous type with $p \in (0, 1)$, and*

$$\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1, \text{ then for each } f \in \mathcal{X}_*, \|f\|_{\mathcal{X}_*} \leq \sup_{N \geq 0} \|S_N f\|_{\mathcal{X}_*} \leq 2^{\frac{1}{p}-1} \|f\|_{\mathcal{X}_*}.$$

Proof. Let $g \in \mathcal{X}$. Observe that, by (iv) of Definition 2, we have $\|g_{r_2}\|_{\mathcal{X}} \geq \|g_{r_1}\|_{\mathcal{X}}$ when $0 < r_1 < r_2 < 1$. If \mathcal{X} is a functional Banach space, then $\|S_N f\|_{\mathcal{X}_*}$ is increasing in N by [21, Proposition 6.1.5, p. 5]. So it suffices to show that $\|f\|_{\mathcal{X}_*} = \sup_{N \geq 0} \|S_N f\|_{\mathcal{X}_*}$.

Note that

$$\|f\|_{\mathcal{X}_*} = \mathbb{E} \|\mathcal{R}f\|_{\mathcal{X}} \leq \sup_{N \geq 0} \lim_{r \rightarrow 1^-} \mathbb{E} \|S_N(\mathcal{R}f_r)\|_{\mathcal{X}} = \sup_{N \geq 0} \|S_N f\|_{\mathcal{X}_*}.$$

Conversely, by [21, Proposition 6.1.5, p. 5],

$$\|S_N f\|_{\mathcal{X}_*} = \mathbb{E} \|S_N(\mathcal{R}f)\|_{\mathcal{X}} \leq \mathbb{E} \|S_N(\mathcal{R}f) + (\mathcal{R}f - S_N(\mathcal{R}f))\|_{\mathcal{X}} = \|f\|_{\mathcal{X}_*}.$$

If \mathcal{X} is a functional p -Banach space, the proof of $\|f\|_{\mathcal{X}_*} \leq \sup_{N \geq 0} \|S_N f\|_{\mathcal{X}_*}$ is a modification of that for Banach spaces. For the other direction, arguing as in [21, Proposition 6.1.5, p. 5],

$$\|S_N f\|_{\mathcal{X}_*} \leq 2^{\frac{1}{p}-1} (\mathbb{E} \|S_N(\mathcal{R}f) + (\mathcal{R}f - S_N(\mathcal{R}f))\|_{\mathcal{X}}^p)^{1/p} = 2^{\frac{1}{p}-1} \|f\|_{\mathcal{X}_*},$$

as desired. □

4. Maximal homogeneous extension

In this section, we prove Theorem B, ending with three nuggets supplementary to Theorem B. The following lemma is not only needed in the proof of Theorem B but also of independent interest, whose proof can be referred in the Appendix in [11].

Lemma 9. *Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter, we also assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. Let $\{X_n\}_{n \geq 0}$ be a standard random sequence, and $s = \min\{1, p\}$.*

- (i) *If $f \in \mathcal{X}_*$, then $\mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}^s < \infty$. Moreover, $f \in (\mathcal{X}_T)_*$ and $\mathbb{E}(\exp(\lambda\|\mathcal{R}f\|_{\mathcal{X}}^s)) < \infty$ for some small enough $\lambda > 0$. This implies, in particular, $\mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}^t < \infty$ for all $t > 0$.*
- (ii) *If $\mathcal{X} = \mathcal{X}^{\max}$ and now we also assume that $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{1/n} = 1$ for the Banach space case, then, for $f \in H(\mathbb{D})$, $f \in \mathcal{X}_*$ if and only if $\mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}}^s < \infty$.*

Proof of Theorem B. (i) Let

$$\mathcal{X}^{\max} = \left\{ f \in H(\mathbb{D}) : f_r \in \mathcal{X} \text{ for } 0 \leq r < 1, \|f\|_{\max} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} < \infty \right\}.$$

We will show that \mathcal{X}^{\max} is the unique maximal homogeneous extension of \mathcal{X} . The proof for the case of p -Banach spaces is similar to that of Banach spaces and will be skipped. The proof is divided into three steps.

Step 1. $(\mathcal{X}^{\max}, \|\cdot\|_{\max})$ is a Banach space.

Let $\{f_n\}_{n \geq 1}$ be a Cauchy sequence in \mathcal{X}^{\max} with $f_n(z) = \sum_{k=0}^{\infty} a_k^{(n)} z^k$. Then, $\{(f_n)_r\}_{n \geq 1}$ is also Cauchy in \mathcal{X} for each $r \in (0, 1)$. Since \mathcal{X} is a Banach space, we assume that $(f_n)_r(z) = \sum_{k=0}^{\infty} a_k^{(n)} r^k z^k$ converges to $\sum_{k=0}^{\infty} a_{k,r} z^k$ in \mathcal{X} as $n \rightarrow \infty$. By (i) in Definition 2, for each fixed $k \geq 0$, $a_k^{(n)} r^k \rightarrow a_{k,r}$ as $n \rightarrow \infty$. Then, $a_k^{(n)} \rightarrow \frac{a_{k,r}}{r^k} \doteq a_k$ as $n \rightarrow \infty$. Let $g(z) = \sum_{k=0}^{\infty} a_k z^k$. Then, for each $r \in (0, 1)$, $g_r(z) = \sum_{k=0}^{\infty} a_k r^k z^k = \sum_{k=0}^{\infty} a_{k,r} z^k \in \mathcal{X}$. By (i) in Definition 2 again, the point evaluation functionals are uniformly bounded on each compact subset in \mathbb{D} . So $\{f_n\}_{n \geq 1}$ converges uniformly to g on each compact subset in \mathbb{D} . This implies that $g \in H(\mathbb{D})$. Then,

$$\|f_n - g\|_{\max} = \sup_{0 \leq r < 1} \|(f_n)_r - g_r\|_{\mathcal{X}} \leq \sup_{0 \leq r < 1} \sup_{m \geq N} \|(f_n)_r - (f_m)_r\|_{\mathcal{X}} = \sup_{m \geq N} \|f_n - f_m\|_{\max},$$

which is arbitrarily small. Hence, $g \in \mathcal{X}^{\max}$, as desired.

Step 2. \mathcal{X}^{\max} is of homogeneous type.

Let $z \in \mathbb{D}$ and $f \in \mathcal{X}^{\max}$. Then, $f_r \in \mathcal{X}$ with $0 \leq r < 1$, and

$$|f(rz)| = |\delta_z f_r| \leq \|\delta_z\|_{\mathcal{X}^*} \|f_r\|_{\mathcal{X}} \leq \|\delta_z\|_{\mathcal{X}^*} \|f\|_{\max}.$$

Letting $r \rightarrow 1^-$, we see that $\{\delta_z : z \in \mathbb{D}\} \subset (\mathcal{X}^{\max})^*$. Now, let $f \in \mathcal{X}^{\max}$ and $\xi \in \mathbb{T}$. Then, $\|f\|_{\max} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} = \sup_{0 \leq r < 1} \|(f_r)_{\xi}\|_{\mathcal{X}} = \|f_{\xi}\|_{\max}$, as desired. For each $f \in \mathcal{X}^{\max}$, define $F : \mathbb{D} \rightarrow \mathcal{X}^{\max}$ with $F(z) = f_z, z \in \mathbb{D}$. The proof of the continuity of F follows from similar arguments as those in the proof of Theorem A. Lastly, for $f \in \mathcal{X}^{\max}$, $\|f\|_{\max} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} = \sup_{0 \leq r < 1} \|f_r\|_{\max}$.

Step 3. \mathcal{X}^{\max} is the maximal homogeneous extension of \mathcal{X} .

Clearly, $\mathcal{X} \subset \mathcal{X}^{\max}$, and $f_r \in \mathcal{X}$ if $f \in \mathcal{X}^{\max}, 0 \leq r < 1$. If $f \in \mathcal{X}$, then $\|f\|_{\max} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} = \|f\|_{\mathcal{X}}$. If \mathcal{Y} is a homogeneous extension of \mathcal{X} , and $f \in \mathcal{Y}$, then $\|f\|_{\mathcal{Y}} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{Y}} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} < \infty$, which implies that $f \in \mathcal{X}^{\max}$.

We now turn to (ii). Only the proof for Banach spaces will be given. $(\mathcal{X}_{\mathcal{P}})_{\star}$ is a solid space since Proposition 14 and Lemma 10 imply that $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is an unconditional basis for $(\mathcal{X}_{\mathcal{P}})_{\star}$, which yields the solidity of $(\mathcal{X}_{\mathcal{P}})_{\star}$ by [25, Theorem II.7, p. 89]. Now, for $(\mathcal{X}^{\max})_{\star}$, we would like to prove $(\mathcal{R}f)^{\lambda}(z) \doteq \sum_{n=0}^{\infty} a_n \lambda_n X_n z^n \in \mathcal{X}^{\max}$ a.s. if $f(z) = \sum_{n=0}^{\infty} a_n z^n \in (\mathcal{X}^{\max})_{\star}$ and $\lambda = \{\lambda_n\}_{n \geq 0} \in \ell^{\infty}$. By Lemma 7, for $0 \leq r < 1$, $\lim_{N \rightarrow \infty} \|S_N(\mathcal{R}f)_r - (\mathcal{R}f)_r\|_{\mathcal{X}} = 0$ a.s. Lemma 9 and the dominated convergence theorem imply that $\lim_{N \rightarrow \infty} \mathbb{E}(\|S_N(\mathcal{R}f)_r - (\mathcal{R}f)_r\|_{\mathcal{X}}) = 0$. By the contraction principle ([21, Theorem 6.1.13, p. 9]),

$$\mathbb{E}\|S_M(\mathcal{R}f)_r^{\lambda} - S_N(\mathcal{R}f)_r^{\lambda}\|_{\mathcal{X}} \lesssim \|\lambda\|_{\ell^{\infty}} \mathbb{E}\|S_M(\mathcal{R}f)_r - S_N(\mathcal{R}f)_r\|_{\mathcal{X}}, \tag{4}$$

which implies that $\{S_N(\mathcal{R}f)_r^{\lambda}\}_{N \geq 0}$ converges in $L^1(\Omega; \mathcal{X})$, leading to a convergent subsequence. The subsequence principle [25, Theorem III.5, p. 132] implies that $S_N(\mathcal{R}f)_r^{\lambda}$ converges in \mathcal{X} a.s. So $(\mathcal{R}f)_r^{\lambda}(z) = \sum_{n=0}^{\infty} \lambda_n a_n r^n X_n z^n \in \mathcal{X}$ a.s. Moreover, by (4) and the contraction principle ([21, Theorem 6.1.13, p. 9]),

$$\mathbb{E}\|(\mathcal{R}f)_r^{\lambda}\|_{\mathcal{X}} \lesssim \|\lambda\|_{\ell^{\infty}} \lim_{N \rightarrow \infty} \mathbb{E}\|S_N(\mathcal{R}f)_r\|_{\mathcal{X}} \leq \|\lambda\|_{\ell^{\infty}} \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}} < \infty.$$

Then,

$$\mathbb{E}\|(\mathcal{R}f)^{\lambda}\|_{\max} = \sup_{0 \leq r < 1} \mathbb{E}\|(\mathcal{R}f)_r^{\lambda}\|_{\mathcal{X}} \lesssim \|\lambda\|_{\ell^{\infty}} \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}} = \|\lambda\|_{\ell^{\infty}} \mathbb{E}\|\mathcal{R}f\|_{\max} < \infty.$$

It follows that $(\mathcal{R}f)^{\lambda} \in \mathcal{X}^{\max}$ a.s., as desired. □

Remark. In Theorem B (ii), the condition $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$ can be dropped for the case $(\mathcal{X}_{\mathcal{P}})_{\star}$ if \mathcal{X} is a functional p -Banach space, since $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ form an unconditional basis for $(\mathcal{X}_{\mathcal{P}})_{\star}$. The same remark applies to Lemma 10.

The rest of this section contains three nuggets supplementary to Theorem B: a counterexample to the solidity of \mathcal{X}_* , the commutativity of $\mathcal{X}_{\mathcal{P}}$ and \mathcal{X}^{\max} with \mathcal{X}_* and the solidity of $(\mathcal{X}_T)_*$.

Example. We show that the symbol space \mathcal{X}_* is not always a solid space. Let \mathcal{X}_*^S denote the symbol space under the randomization by the standard Steinhaus sequence. Let \mathcal{B} denote the Bloch space and

$$\mathcal{X} = \left\{ f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{B} : \lim_{k \rightarrow \infty} |a_{2^k}| \text{ exists} \right\}.$$

Recall that $\|f\|_{\mathcal{B}} = |f(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)|$ for $f \in \mathcal{X}$ and one can check that $(\mathcal{X}, \|\cdot\|_{\mathcal{B}})$ is of homogeneous type. For $(\mathcal{X}, \|\cdot\|_{\mathcal{B}})$, the hypotheses of Definition 2 are satisfied since \mathcal{B} is of homogeneous type. So, it is sufficient to prove that $(\mathcal{X}, \|\cdot\|_{\mathcal{B}})$ is complete. Let $\{f_m\}_{m \geq 1}$ be a Cauchy sequence in \mathcal{X} with $f_m(z) = \sum_{n=0}^{\infty} a_n^{(m)} z^n$. Since \mathcal{B} is complete, there exists $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{B}$ such that $\lim_{m \rightarrow \infty} \|f_m - f\|_{\mathcal{B}} = 0$. By [19, p. 16], for any $g(z) = \sum_{n=0}^{\infty} b_n z^n \in \mathcal{B}$, $|b_n| \leq 2\|g\|_{\mathcal{B}}$, $\forall n \geq 1$. Then,

$$\begin{aligned} \left| |a_{2^k}| - |a_{2^j}| \right| &\leq \left| |a_{2^k}| - |a_{2^k}^{(m)}| \right| + \left| |a_{2^k}^{(m)}| - |a_{2^j}^{(m)}| \right| + \left| |a_{2^j}^{(m)}| - |a_{2^j}| \right| \\ &\leq 4\|f_m - f\|_{\mathcal{B}} + \left| |a_{2^k}^{(m)}| - |a_{2^j}^{(m)}| \right|. \end{aligned}$$

So $\{|a_{2^k}|\}_{k \geq 1}$ is a Cauchy sequence in \mathbb{R} , hence $f \in \mathcal{X}$. That is, $(\mathcal{X}, \|\cdot\|_{\mathcal{B}})$ is of homogeneous type. Lastly, we point out that $\sum_{k=1}^{\infty} z^{2^k} \in \mathcal{X}_*^S$. By [19, Theorem 1.14, p. 15], for any lacunary sequence $\{n_k\}_{k \geq 1}$, that is, $\inf_{k \geq 1} \frac{n_{k+1}}{n_k} > 1$, the series $\sum_{k=1}^{\infty} a_k z^{n_k}$ belongs to \mathcal{B} if and only if $\sup_{k \geq 1} |a_k| < \infty$. So the Steinhaus randomization $\sum_{k=1}^{\infty} e^{2\pi i \alpha_k} z^{2^k} \in \mathcal{X}$ a.s., that is, $\sum_{k=1}^{\infty} z^{2^k} \in \mathcal{X}_*^S$. But $\left\{ \sum_{k=1}^{\infty} \lambda_k z^{2^k} : \{\lambda_k\}_{k \geq 1} \in \ell^\infty \right\} \not\subset \mathcal{X}_*^S$. For example, taking $\lambda_{2^k} = 1$ and $\lambda_{2^k-1} = 0$ for $k \geq 1$, one has $\sum_{k=1}^{\infty} \lambda_k z^{2^k} \notin \mathcal{X}_*^S$.

The above example shows that \mathcal{X}_*^S is not always a solid space. The same is true for \mathcal{X}_*^R obtained through the Rademacher sequence. For the standard real (or complex) Gaussian randomization, we now show that \mathcal{X}_*^G is a solid space. Here, we consider only the real Gaussian case, and the complex Gaussian case can be treated in a similar fashion. If $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}_*^G$, then $\lim_{k \rightarrow \infty} |a_{2^k} \xi_{2^k}|$ exists a.s., where $\{\xi_n\}_{n \geq 0}$ is a standard real Gaussian sequence. By the Kolmogorov zero-one law, $\lim_{k \rightarrow \infty} |a_{2^k} \xi_{2^k}| = c$ a.s. for some constant $c \geq 0$. Since $\limsup_{n \rightarrow \infty} \frac{|\xi_n|}{\sqrt{2 \log n}} = 1$ a.s. [25, p. 43], one has $a_{2^k} \rightarrow 0$ as $k \rightarrow \infty$.

Furthermore, by a direct verification, $a_{2k}\xi_{2k}$ converges to 0 in distribution as $k \rightarrow \infty$. It follows that $c=0$. Let $\{\lambda_n\}_{n \geq 0} \in \ell^\infty$. Then, $\sum_{n=0}^\infty a_n \lambda_n z^n \in \mathcal{B}_*^G$ by the remark after Theorem B, and $\lim_{k \rightarrow \infty} a_{2k} \lambda_{2k} \xi_{2k} = 0$ a.s. That is, $\sum_{n=0}^\infty a_n \lambda_n z^n \in \mathcal{X}_*^G$.

Problem A. To characterize homogeneous \mathcal{X} such that \mathcal{X}_* is a solid space.

The second nugget is the following commutativity:

Lemma 10. Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter, we also assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. Let $\{X_n\}_{n \geq 0}$ be a standard random sequence. Then,

$$(\mathcal{X}_*)_{\mathcal{P}} = (\mathcal{X}_{\mathcal{P}})_* \quad \text{and} \quad (\mathcal{X}_*)^{\max} = (\mathcal{X}^{\max})_*$$

Proof. We first claim that if $g \in \mathcal{X}_{\mathcal{P}}$, then $\lim_{r \rightarrow 1^-} g_r = g$, where the convergence is in norm. Assume that $p_n \rightarrow g$ as $n \rightarrow \infty$ with $p_n \in \mathcal{P}$. Since $\lim_{r \rightarrow 1^-} \|(p_n)_r - p_n\|_{\mathcal{X}} = 0$ for each n , the claim follows from

$$\begin{aligned} \|g_r - g\|_{\mathcal{X}}^p &\leq \|g_r - (p_n)_r\|_{\mathcal{X}}^p + \|(p_n)_r - p_n\|_{\mathcal{X}}^p + \|p_n - g\|_{\mathcal{X}}^p \\ &\leq 2\|p_n - g\|_{\mathcal{X}}^p + \|(p_n)_r - p_n\|_{\mathcal{X}}^p \end{aligned}$$

since \mathcal{X} is of homogeneous type. In particular, if \mathcal{X} is a Banach space, we take $p=1$. Now let $f \in (\mathcal{X}_{\mathcal{P}})_*$. The claim implies that $(\mathcal{R}f)_r \rightarrow \mathcal{R}f$ a.s. as $r \rightarrow 1^-$. The Marcinkiewicz–Zygmund–Kahane theorem [26, Theorem II.4, p. 240] implies that $S_N(\mathcal{R}f) \rightarrow \mathcal{R}f$ a.s. as $N \rightarrow \infty$. By the dominated convergence theorem, one has $\lim_{N \rightarrow \infty} \mathbb{E}\|S_N(\mathcal{R}f) - \mathcal{R}f\|_{\mathcal{X}} = 0$, which yields that $f \in (\mathcal{X}_*)_{\mathcal{P}}$. Conversely, let $f \in (\mathcal{X}_*)_{\mathcal{P}}$. By Theorem A, one has $\lim_{N \rightarrow \infty} S_N f = f$ in \mathcal{X}_* . So $f \in (\mathcal{X}_{\mathcal{P}})_*$, as desired.

For the second equality, if $f \in (\mathcal{X}_*)^{\max}$, then $(\mathcal{R}f)_r \in \mathcal{X}$ a.s., $0 \leq r < 1$ and $\sup_{0 \leq r < 1} \mathbb{E}\|(\mathcal{R}f)_r\|_{\mathcal{X}} = \mathbb{E}\left(\sup_{0 \leq r < 1} \|(\mathcal{R}f)_r\|_{\mathcal{X}}\right) < \infty$ since \mathcal{X} is homogeneous. It follows that $\sup_{0 \leq r < 1} \|(\mathcal{R}f)_r\|_{\mathcal{X}} < \infty$ a.s., which implies that $f \in (\mathcal{X}^{\max})_*$. Conversely, let $f \in (\mathcal{X}^{\max})_*$. Then, $(\mathcal{R}f)_r \in \mathcal{X}$ a.s., $0 \leq r < 1$ and $\sup_{0 \leq r < 1} \|(\mathcal{R}f)_r\|_{\mathcal{X}} < \infty$ a.s. The Marcinkiewicz–Zygmund–Kahane theorem [26, Theorem II.4, p. 240] implies $\sup_{N \geq 0} \|S_N(\mathcal{R}f)\|_{\mathcal{X}} < \infty$ a.s. Moreover, $\lim_{N \rightarrow \infty} S_N(\mathcal{R}f)_r = (\mathcal{R}f)_r$ a.s. by Lemma 7. By Kahane’s inequality [25, Theorem V.1, p. 139] and Fernique’s theorem [25, Theorem V.26, p. 255],

$$\sup_{0 \leq r < 1} \mathbb{E}\|(\mathcal{R}f)_r\|_{\mathcal{X}} \leq \mathbb{E}\left(\sup_{0 \leq r < 1} \|(\mathcal{R}f)_r\|_{\mathcal{X}}\right) \leq \mathbb{E}\left(\sup_{N \geq 0} \|S_N(\mathcal{R}f)\|_{\mathcal{X}}\right) < \infty.$$

So $f \in (\mathcal{X}_*)^{\max}$, as desired. □

Lastly, as the third nugget, the following proposition should be compared with Proposition 15, which identifies \mathcal{X}_T with \mathcal{X}^{\max} under certain conditions.

Proposition 11. *Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type. Let $\{X_n\}_{n \geq 0}$ be a standard random sequence. Then, $(\mathcal{X}_T)_*$ is a solid space.*

Proof. We shall only prove the Banach space case since the modification needed for p -Banach spaces is straightforward. Let $f(z) = \sum_{n=0}^\infty a_n z^n \in (\mathcal{X}_T)_*$. By Lemma 9, $(\mathcal{R}f)^\lambda \in \mathcal{X}_T$ a.s. if and only if $\mathbb{E}\|(\mathcal{R}f)^\lambda\|_{\mathcal{X}_T} < \infty$, where $(\mathcal{R}f)^\lambda(z) = \sum_{n=0}^\infty a_n \lambda_n X_n z^n$ and $\lambda = \{\lambda_n\}_{n \geq 0} \in \ell^\infty$. Note that

$$\begin{aligned} \mathbb{E}\|(\mathcal{R}f)^\lambda\|_{\mathcal{X}_T} &\lesssim \lim_{N \rightarrow \infty} \int_0^\infty \mathbb{P}(\|(S_N(\mathcal{R}f))^\lambda\|_{\mathcal{X}} > t) dt \\ &\lesssim \sup_{n \geq 0} |\lambda_n| \lim_{N \rightarrow \infty} \mathbb{E}\|S_N(\mathcal{R}f)\|_{\mathcal{X}} \\ &\leq \sup_{n \geq 0} |\lambda_n| \mathbb{E}\|\mathcal{R}f\|_{\mathcal{X}_T} < \infty, \end{aligned}$$

where the first ‘ \lesssim ’ follows from Lévy’s inequality [11], and the second one holds by the contraction principle ([21, Theorem 6.1.13, p. 9]). □

5. Relationship with unconditional bases

In this section, we first present the (rather short) proof of Theorem C. Then, we present four propositions to further illustrate the connection between solid bases/frames and unconditional bases.

Proof of Theorem C. (i.1) \Rightarrow (i.2) is obvious. (i.2) \Rightarrow (i.3) is by the fact that $\lim_{N \rightarrow \infty} S_N f = f$ if and only if $f \in \mathcal{X}_{\mathcal{P}}$ in the proof of Proposition 14. Next, (i.3) \Rightarrow (i.4). Part (i) of Definition 1 is satisfied since $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is an unconditional basis for $\mathcal{X}_{\mathcal{P}}$ with constant 1. It suffices to show that $\lim_{N \rightarrow \infty} \|S_N f\|_{\max} = \|f\|_{\max}$ for $f \in \mathcal{X}^{\max}$. Since $f_r \in \mathcal{X}_{\mathcal{P}}$ for each $0 \leq r < 1$ by Lemma 7, one has

$$\lim_{N \rightarrow \infty} \|S_N f_r\|_{\mathcal{X}} = \sup_{N \geq 0} \|S_N f_r\|_{\mathcal{X}} = \|f_r\|_{\mathcal{X}} \tag{5}$$

because $\|S_N f_r\|_{\mathcal{X}}$ increases as $N \rightarrow \infty$. Then,

$$\sup_{N \geq 0} \|S_N f\|_{\max} = \sup_{N \geq 0} \sup_{0 \leq r < 1} \|S_N f_r\|_{\mathcal{X}} = \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} = \|f\|_{\max}, \tag{6}$$

as desired. (i.4) \Rightarrow (i.1) follows from the definition of the homogeneous extension. The proof for p -Banach spaces follows from a modification by replacing (5) with $\sup_{N \geq 0} \|S_N f_r\|_{\mathcal{X}} \approx \|f_r\|_{\mathcal{X}}$. □

Remark. In Theorem C (ii), the condition $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$ can be dropped for the case $\mathcal{X} = \mathcal{X}_{\mathcal{P}}$ if \mathcal{X} is a functional p -Banach space. This follows from the fact that $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a Schauder basis for $\mathcal{X}_{\mathcal{P}}$ if and only if $\sup_{N \geq 0} \|S_N\| < \infty$; see [43, Proposition 1]. The same remark applies to Theorem E and Theorem F.

The rest of this section is to prove the following four propositions, which further illustrate the relationship between solid bases/frames and unconditional bases.

Proposition 12. *Let E be a Banach space. Then, $\{e_n\}_{n \geq 1}$ is an unconditional basis for E with constant 1 if and only if it is both a solid basis and a Schauder basis for E .*

Recall that a sequence $\{e_n\}_{n \geq 1}$ in a Banach space E is said to be a *basic sequence* if it is a Schauder basis for the closure of its span. Let $e_n = (0, \dots, 0, 1, 0, \dots)$, $n \geq 1$.

Proposition 13. *Let E be a Banach space and $\{e_n\}_{n \geq 1}$ be a basic sequence with $\|e_n\|_E = 1$, $n \geq 1$. Let W denote the closure of finite linear combinations of $\{e_n\}_{n \geq 1}$. Then, $\{e_n\}_{n \geq 1}$ is an unconditional basis for W if and only if $\{e_n\}_{n \geq 1}$ is a solid frame for $W^{\text{s-max}}$.*

Here, the sequential version of the maximal solid extension $E^{\text{s-max}}$ is defined as

$$E^{\text{s-max}} = \left\{ a = (a_1, a_2, \dots) \in \mathfrak{S} : \|a\|_{\text{s-max}} = \sup_{N \geq 1} \left\| \sum_{n=1}^N a_n e_n \right\|_E < \infty \right\}, \tag{7}$$

which is the unique maximal solid extension of E up to an isometric isomorphism; see Theorem G.

Proposition 14.

- (i) *Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. Then, $\{e_n\}_{n \geq 1}$ is an unconditional basis for $E_{\mathcal{P}}$ with constant 1.*
- (ii) *Let E be a p -Banach space with $p \in (0, 1)$ and $\{e_n\}_{n \geq 1}$ a solid frame. Then, $\{e_n\}_{n \geq 1}$ is an unconditional basis for $E_{\mathcal{P}}$.*

Proposition 15. *Let \mathcal{X} be a functional Banach space, or a functional p -Banach space with $p \in (0, 1)$, of homogeneous type; for the latter, we assume $\lim_{n \rightarrow \infty} \|z^n\|_{\mathcal{X}}^{\frac{1}{n}} = 1$. If $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is an unconditional basis for $\mathcal{X}_{\mathcal{P}}$, then $\mathcal{X}^{\text{max}} = \mathcal{X}_{\mathcal{T}}$.*

Proof of Proposition 12. If $\{e_n\}_{n \geq 1}$ is an unconditional basis with constant 1, then, by [25, Proposition III.5, p.92], part (i) of Definition 1 is satisfied. Condition (ii) holds since it is a Schauder basis. Conversely, if $\{e_n\}_{n \geq 1}$ is a solid basis, then part (i) of Definition 1 ensures that $\{e_n\}_{n \geq 1}$ is an unconditional basis with constant 1. □

Proof of Proposition 13. This proof follows from the following lemma which is less sleek but covers a slightly more general situation. □

Lemma 16. *Let E be a Banach space and $\{e_n\}_{n \geq 1}$ be a sequence of unit vectors. Let W denote the closure of finite linear combinations of $\{e_n\}_{n \geq 1}$.*

- (i) *If $\{e_n\}_{n \geq 1}$ is an unconditional basis for W , then $\{\mathbf{e}_n\}_{n \geq 1}$ is a solid frame for $W^{s\text{-max}}$.*
- (ii) *If $\{e_n\}_{n \geq 1}$ is a Schauder basis for W and $\{\mathbf{e}_n\}_{n \geq 1}$ is a solid frame for $W^{s\text{-max}}$, then $\{e_n\}_{n \geq 1}$ is an unconditional basis for W .*

Proof of Lemma 16. (i) If $\{e_n\}_{n \geq 1}$ is an unconditional basis for W , then $\|\mathbf{e}_n\|_{s\text{-max}} = \|e_n\|_E = 1$. Moreover,

$$\left\| \sum_{k=1}^N \lambda_k a_k \mathbf{e}_k \right\|_{s\text{-max}} = \sup_{1 \leq n \leq N} \left\| \sum_{k=1}^n \lambda_k a_k e_k \right\|_E \lesssim \sup_{1 \leq n \leq N} \left\| \sum_{k=1}^n a_k e_k \right\|_E = \left\| \sum_{k=1}^N a_k \mathbf{e}_k \right\|_{s\text{-max}}$$

for all $|\lambda_k| \leq 1$. For any $a \in W^{s\text{-max}}$, let $\tilde{P}_N a = \sum_{k=1}^N a_k \mathbf{e}_k$, then

$$\|a\|_{s\text{-max}} = \sup_{N \geq 1} \left\| \sum_{k=1}^N a_k e_k \right\|_E = \lim_{N \rightarrow \infty} \sup_{1 \leq n \leq N} \left\| \sum_{k=1}^n a_k e_k \right\|_E = \lim_{N \rightarrow \infty} \left\| \tilde{P}_N a \right\|_{s\text{-max}}.$$

So $\{\mathbf{e}_n\}_{n \geq 1}$ is a solid frame for $W^{s\text{-max}}$.

(ii) Since $\{e_n\}_{n \geq 1}$ is a Schauder basis for W , there exists a coefficient map $\mathcal{T}_1 : W \rightarrow W^{s\text{-max}}$ with $\mathcal{T}_1 x = ((\mathcal{T}x)_1, \dots, (\mathcal{T}x)_n, \dots)$, and $\sum_{n=1}^N (\mathcal{T}_1 x)_n e_n$ converges to x in W . By [25, Theorem II.3, p. 48],

$$\|x\|_E \approx \sup_{N \geq 1} \left\| \sum_{n=1}^N (\mathcal{T}_1 x)_n e_n \right\|_E = \left\| ((\mathcal{T}_1 x)_n)_{n \geq 1} \right\|_{s\text{-max}}.$$

For each $(\theta_n)_{n \geq 1} \in \{-1, 1\}^{\mathbb{N}}$, since $\{\mathbf{e}_n\}_{n \geq 1}$ is a solid frame for $W^{s\text{-max}}$, one has

$$\left\| \sum_{n=N+1}^M \theta_n (\mathcal{T}_1 x)_n e_n \right\|_E \lesssim \left\| \sum_{n=N+1}^M (\mathcal{T}_1 x)_n \mathbf{e}_n \right\|_{s\text{-max}} \approx \left\| \sum_{n=N+1}^M (\mathcal{T}_1 x)_n e_n \right\|_E,$$

where $M > N$. So $\sum_{n=1}^N (\mathcal{T}_1 x)_n e_n$ converges unconditionally in W , as desired. □

Proof of Proposition 14. (i) In order to show that $\{e_n\}_{n \geq 1}$ is a Schauder basis for E_p , we shall prove that for $x \in E$, $\lim_{N \rightarrow \infty} P_N(x) = x \iff x \in E_p$. It suffices to show the sufficiency. Assume that $\lim_{n \rightarrow \infty} x_n = x$ for $x_n = \sum_{j=1}^{k_n} c_j^{(n)} e_j$, where $c_j^{(n)} \in \mathbb{C}$. For $\epsilon > 0$, take n large enough so that $\|x_n - x\|_E < \epsilon$. For any $N > k_n$, $\|P_N x - x\|_E \leq \|P_N x - P_N x_n\|_E + \|P_N x_n - x_n\|_E + \|x_n - x\|_E \leq 2\epsilon$. Now the conclusion follows from [25, Proposition III.3, p. 91] and Definition 1. The proof of part (ii) for p -Banach spaces follows from [1, Theorem 1.10], together with a straightforward modification of the arguments in part (i). □

Proof of Proposition 15. If $f \in \mathcal{X}^{\max}$, then $\|f\|_{\mathcal{X}_T} = \sup_{N \geq 0} \|S_N f\|_{\mathcal{X}} \approx \|f\|_{\max}$, where ‘ \approx ’ can be strengthened to an equality, by (6), if \mathcal{X} is a Banach space. Hence, $f \in \mathcal{X}_T$. Conversely, let $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}_T$. We claim that $f_r \in \mathcal{X}$ for $0 \leq r < 1$. It suffices to show that $\left\{ \sum_{k=0}^n a_k r^k z^k \right\}_{n \geq 0}$ is a Cauchy sequence in \mathcal{X} . Since $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is an unconditional basis for \mathcal{X}_p , by [25, Proposition III.5, p. 92] or [1, Theorem 1.10], for $m > n$,

$$\left\| \sum_{k=n}^m a_k r^k z^k \right\|_{\mathcal{X}} \lesssim r^n \left\| \sum_{k=n}^m a_k z^k \right\|_{\mathcal{X}} \lesssim r^n \sup_{N \geq 0} \left\| \sum_{k=0}^N a_k z^k \right\|_{\mathcal{X}} \lesssim r^n \|f\|_{\mathcal{X}_T} \rightarrow 0$$

as $n \rightarrow \infty$. Then, $f \in \mathcal{X}^{\max}$. \square

Example. For $0 < p < \infty$, recall that H^p , L_a^p , \mathcal{D}^p , \mathcal{B} and $A(\mathbb{D})$ denote the Hardy space, the Bergman space, the Dirichlet space, the Bloch space and the disk algebra, respectively. If $\mathcal{X} = H^p$, L_a^p , or \mathcal{D}^p , with $1 < p < \infty$, then $\mathcal{X}_T = \mathcal{X}^{\max} = \mathcal{X}$. On the other hand, if $\mathcal{X} = H^\infty$, \mathcal{B} , BMOA, H^1 , L_a^1 , or $A(\mathbb{D})$, then $\mathcal{X}_T \subsetneq \mathcal{X}^{\max} = \mathcal{X}$ since there exists $f_0 \in \mathcal{X}$ such that $\|S_N f_0\|_{\mathcal{X}} \rightarrow \infty$ as $N \rightarrow \infty$. See [5, 43] for more details.

Problem B. To characterize homogeneous \mathcal{X} such that $\mathcal{X}^{\max} = \mathcal{X}_T$.

Remark. Let $\phi_n(x) = (\mathcal{T}x)_n$, $x \in E$, the coordinate functional, if $\{e_n\}_{n \geq 1}$ form a solid basis for E , then $\{\phi_n\}_{n \geq 1}$ form an unconditional basic sequence in E^* ; that is, $\{\phi_n\}_{n \geq 1}$ form an unconditional basis for $\overline{\text{span}}\{\phi_n : n \geq 1\}$.

Remark. Let \mathcal{X} be a Banach space of analytic functions over \mathbb{D} with $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ as a solid basis. If $\mathcal{X} = \mathcal{Y}^*$ for some separable Banach space \mathcal{Y} of analytic functions over \mathbb{D} . Then, $\mathcal{X} = \mathcal{X}_T$. It suffices to show that $\mathcal{X}_T \subset \mathcal{X}$. Let $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}_T$. Then, $\sup_{N \geq 1} \left\| \sum_{n=0}^N a_n z^n \right\|_{\mathcal{X}} < \infty$. Since \mathcal{Y} is separable, the w^* -topology on \mathcal{X} is metrizable. The Banach–Alaoglu theorem implies that there exists a subsequence $\{N_k\}_{k \geq 1}$ such that $f_k = \sum_{n=0}^{N_k} a_n z^n$ converges to some $f \in \mathcal{X}$ in w^* -topology. By [9], $f_k(z) \rightarrow f(z)$ as $k \rightarrow \infty$. Then, for each fixed n , it can be checked that the n th Taylor coefficient of f is a_n , as desired.

6. Weak-star convergence of Taylor polynomials

In this section, we prove Theorem D, which states that, in the presence of a solid basis, Taylor polynomials are w^* -convergent in the double dual. When $E = E_{\mathcal{P}}$, Proposition 14 implies that the Taylor polynomials converge in norm. Hence, the strength of Theorem D lies in the non-separable case.

For any homogeneous functional Banach space \mathcal{X} , the Taylor polynomials are certainly pointwisely convergent, hence locally uniformly convergent by the principle of uniform boundedness. When $\mathcal{X} = \mathcal{X}_{\mathcal{P}}$, the norm convergence of Taylor polynomials, which is equivalent to the weak convergence, is well understood; indeed, the following are equivalent:

- (i) $\{S_N f\}_{N \geq 0}$ converges in norm for each $f \in \mathcal{X}$;
- (ii) $\{S_N f\}_{N \geq 0}$ converges in w -topology for each $f \in \mathcal{X}$;
- (iii) $\sup_{N \geq 0} \|S_N\| < \infty$.

We only need to check (ii) \Rightarrow (iii) since the equivalence between (i) and (iii) follows from Proposition 1 in [43]. It suffices to observe that the partial summation operator S_N is bounded for each $N \in \mathbb{N}$, for then the implication follows from the principle of uniform boundedness. Since the coefficient functionals are continuous,

$$\|S_N f\|_{\mathcal{X}} \leq \sum_{n=0}^N |a_n| \|z^n\|_{\mathcal{X}} \leq \sum_{n=0}^N c_n \|f\|_{\mathcal{X}} \|z^n\|_{\mathcal{X}} = \left(\sum_{n=0}^N c_n \|z^n\|_{\mathcal{X}} \right) \|f\|_{\mathcal{X}}$$

for some positive sequence $\{c_n\}_{n \geq 0}$.

If $\mathcal{X} = \mathcal{Y}^*$ for some separable Banach space \mathcal{Y} , and still assuming that $\mathcal{X} = \mathcal{X}_{\mathcal{P}}$, then $\{S_N f\}_{N \geq 0}$ is w^* -convergent if and only if $\{S_N f\}_{N \geq 0}$ converges uniformly on compacta in \mathbb{D} and $\sup_{N \geq 0} \|S_N\| < \infty$; see [9].

If $\mathcal{X} \neq \mathcal{X}_{\mathcal{P}}$, then almost nothing affirmative is known about the convergence of Taylor polynomials, at least from a Banach space viewpoint. For instance, in H^∞ , the Taylor polynomials are not convergent in norm, w -topology or w^* -topology.

Proof of Theorem D. Let $\{e_n\}_{n \geq 1}$ be a solid basis for a Banach space E . Then, for any $x \in E$ and $\phi \in E^*$, we claim that

$$\sum_{n=1}^{\infty} |(\mathcal{T}x)_n| |\phi(e_n)| < \infty. \tag{8}$$

Indeed, by (i) of Definition 1, for every finite subset σ , $\sup_{\theta_n = \pm 1} \left\| \sum_{n \in \sigma} \theta_n (\mathcal{T}x)_n e_n \right\|_E \leq \|x\|_E$.

Let $N \geq 1$. Set

$$\theta_n = \begin{cases} 1, & \text{Re } \phi((\mathcal{T}x)_n e_n) \geq 0; \\ -1, & \text{Re } \phi((\mathcal{T}x)_n e_n) < 0. \end{cases}$$

Similarly, set

$$\theta'_n = \begin{cases} 1, & \text{Im } \phi((\mathcal{T}x)_n e_n) \geq 0; \\ -1, & \text{Im } \phi((\mathcal{T}x)_n e_n) < 0. \end{cases}$$

Then,

$$\begin{aligned} \sum_{n=1}^N |\phi((\mathcal{T}x)_n e_n)| &\leq \sum_{n=1}^N |\text{Re } \phi((\mathcal{T}x)_n e_n)| + \sum_{n=1}^N |\text{Im } \phi((\mathcal{T}x)_n e_n)| \\ &\leq \|\phi\|_{E^*} \left\| \sum_{n=1}^N \theta_n (\mathcal{T}x)_n e_n \right\|_E + \|\phi\|_{E^*} \left\| \sum_{n=1}^N \theta'_n (\mathcal{T}x)_n e_n \right\|_E \\ &\leq 2\|\phi\|_{E^*} \|x\|_E. \end{aligned}$$

So the claim (8) holds. Then, $\sum_{n=1}^{\infty} (\mathcal{T}x)_n(Je_n)(\phi)$ exists. Let

$$\Psi(\phi) \doteq \sum_{n=1}^{\infty} (\mathcal{T}x)_n(Je_n)(\phi) = \sum_{n=1}^{\infty} (\mathcal{T}x)_n(\phi e_n),$$

and then $\sum_{n=1}^N (\mathcal{T}x)_n(Je_n) = J(P_Nx)$ converges to Ψ in w^* -topology. □

We let $\mathfrak{J}(x)$ denote the limit element $\Psi \in E^{**}$, so we have a map $\mathfrak{J} : E \rightarrow E^{**}$ which is easily seen to satisfy: $\|\mathfrak{J}\| = 1$, and \mathfrak{J} is injective. It appears interesting to compare this map with the canonical embedding $J : E \rightarrow E^{**}$.

Proposition 17. *Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$.*

- (i) $J(x) = \mathfrak{J}(x)$ if and only if $x \in E_{\mathcal{P}}$.
- (ii) If $E_{\mathcal{P}} \subsetneq E$, then $\overline{\mathfrak{J}E}^{w^*}$ is a proper subspace in E^{**} .

Remark. For the canonical map J , it is known that $\overline{JE}^{w^*} = E^{**}$.

Proof. (i) Let $x \in E_{\mathcal{P}}$. Then $\lim_{N \rightarrow \infty} P_Nx = x$ by Proposition 14. For $\phi \in E^*$,

$$(Jx)(\phi) = \phi(x) \quad \text{and} \quad (\mathfrak{J}x)(\phi) = \lim_{N \rightarrow \infty} \phi(P_Nx) = \phi(x).$$

So $J(x) = \mathfrak{J}(x)$. Conversely, let $x \in E \setminus E_{\mathcal{P}}$. By the Hahn-Banach theorem there exists a nonzero $\phi \in E^*$ such that $\phi|_{E_{\mathcal{P}}} = 0$, and $\phi(x) \neq 0$. Then, $(\mathfrak{J}x)(\phi) = 0$, and $(Jx)(\phi) = \phi(x) \neq 0$. (ii) Recall that $\overline{\mathfrak{J}E}^{w^*} = (\perp \mathfrak{J}E)^{\perp}$ [37, Theorem 4.7, p. 96], where ‘ \perp ’ denotes the annihilators. We claim that $\perp(\mathfrak{J}E) = \{\phi \in E^* : \phi|_{E_{\mathcal{P}}} = 0\}$. If $\phi \in \{\phi \in E^* : \phi|_{E_{\mathcal{P}}} = 0\}$, then, for each $x \in E$, $(\mathfrak{J}x)\phi = \lim_{N \rightarrow \infty} \phi(P_Nx) = 0$. So $\phi \in \perp(\mathfrak{J}E)$. Conversely, let $\phi \in \perp(\mathfrak{J}E)$. Let $x = \lim_{n \rightarrow \infty} x_n \in E_{\mathcal{P}}$ with x_n being a finite linear combination of $\{e_n\}_{n \geq 1}$. For each n , $\phi(x_n) = \lim_{N \rightarrow \infty} \phi(P_Nx_n) = 0$. So the claim holds. It follows that $\{\phi \in E^* : \phi|_{E_{\mathcal{P}}} = 0\} = \perp(JE_{\mathcal{P}})$. Then,

$$\overline{\mathfrak{J}E}^{w^*} = \overline{JE_{\mathcal{P}}}^{w^*} = (\perp(JE_{\mathcal{P}}))^{\perp} = (\perp(\mathfrak{J}E_{\mathcal{P}}))^{\perp},$$

where the last ‘=’ follows from (i). In other words, $\overline{\mathfrak{J}E}^{w^*} = (E_{\mathcal{P}})^{\perp\perp}$. Since $E_{\mathcal{P}} \subsetneq E$, we can take $x_0 \in E \setminus E_{\mathcal{P}}$ and choose, by Hahn-Banach, a nonzero $\phi_0 \in E^*$ such that $\phi_0|_{E_{\mathcal{P}}} = 0$ and $\phi_0(x_0) = 1$. Then, $Jx_0 \in E^{**}$, but $(Jx_0)\phi_0 = \phi_0(x_0) = 1$, so $Jx_0 \in E^{**} \setminus (E_{\mathcal{P}})^{\perp\perp}$, as desired. □

Example. We give an explicit example of $J \neq \mathfrak{J}$ via the so-called Banach limit. Let $E = \ell^{\infty}$. There exists a bounded linear functional φ on ℓ^{∞} [14, p. 82] (the Banach limit) such that

- (i) $\varphi S(x) = \varphi(x)$; and
- (ii) $x \in c, \varphi(x) = \lim_{n \rightarrow \infty} x_n$,

where $S(x_1, x_2, \dots) = (x_2, x_3, \dots)$, and c is the collection of sequences in \mathfrak{S} having limits. By (i), one has $\varphi(x_1, x_2, \dots, x_n, 0, 0, \dots) = 0$. Let $x \in c \setminus c_0$. Then $(Jx)\varphi = \varphi(x) = \lim_{n \rightarrow \infty} x_n \neq 0$, and, $(\mathfrak{J}x)\varphi = \lim_{n \rightarrow \infty} \varphi(x_1, x_2, \dots, x_n, 0, 0, \dots) = 0$. Hence, $J \neq \mathfrak{J}$ on $c \setminus c_0$. Moreover, by the proof of Proposition 17 (ii), $\overline{\mathfrak{J}E}^{w*} = c_0^{\perp\perp}$, which is isometrically isomorphic to c_0^{**} , which is ℓ^∞ . In other words, $c_0^{\perp\perp} \simeq ((\ell^\infty)^*/c_0^\perp)^* \simeq c_0^{**}$.

Example. Here is another example on functional spaces. Let $E = \mathcal{B}_*^S$, the symbol space of the Bloch space under the Steinhaus randomization. Fix a lacunary sequence $\{n_k\}_{k \geq 1}$, i.e., $\inf_{k \geq 1} \frac{n_{k+1}}{n_k} > 1$, and let

$$A = \left\{ f(z) = \sum_{n=0}^\infty a_n z^n \in \mathcal{B}_*^S : \lim_{k \rightarrow \infty} a_{n_k} \text{ exists} \right\}.$$

Then, A is a subspace of \mathcal{B}_*^S . Define a bounded linear functional φ on A by $\varphi(f) = \lim_{k \rightarrow \infty} a_{n_k}$. Note that $|a_{n_k}| \lesssim \|a_{n_k} z^{n_k}\|_{\mathcal{B}_*^S} \leq \|f\|_{\mathcal{B}_*^S}$. The extension of φ to \mathcal{B}_*^S is still denoted by φ . Now take any $f(z) = \sum_{n=0}^\infty a_n z^n \in A$ with $\lim_{k \rightarrow \infty} a_{n_k} \neq 0$. Then $(Jf)(\varphi) = \varphi(f) = \lim_{k \rightarrow \infty} a_{n_k} \neq 0$ and $(\mathfrak{J}f)(\varphi) = \lim_{N \rightarrow \infty} \varphi\left(\sum_{n=0}^N a_n z^n\right) = 0$.

7. Transforming solid frames into solid bases

Proof of Theorem E. We shall prove that the following are equivalent:

- (i) $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid frame for \mathcal{X} .
- (ii) $\|f\|_{\#} \approx \|f\|_{\mathcal{X}}, f \in \mathcal{X}$.
- (iii) For each $f \in \mathcal{X}, \|f\|_{\#} < \infty$.
- (iv) $(\mathcal{X}, \|\cdot\|_{\#})$ is a Banach space (or p -Banach space), and $\left\{ \frac{z^n}{\|z^n\|_{\mathcal{X}}} \right\}_{n \geq 0}$ is a solid basis for $(\mathcal{X}, \|\cdot\|_{\#})$.

Also we only prove the Banach space case since it takes only a straightforward modification to treat p -Banach spaces. (i) \Rightarrow (iii) is clear. Now, (iii) \Rightarrow (iv). Since \mathcal{X} is of homogeneous type,

$$\|f\|_{\mathcal{X}} = \lim_{r \rightarrow 1^-} \|f_r\|_{\mathcal{X}} \leq \sup_{N \geq 0} \sup_{0 \leq r < 1} \left\| \sum_{k=0}^N a_k r^k z^k \right\|_{\mathcal{X}} \leq \|f\|_{\#}. \tag{9}$$

It suffices to show that $(\mathcal{X}, \|\cdot\|_{\#})$ is a Banach space. Let $\{f_j\}_{j \geq 1}$ be a Cauchy sequence in $(\mathcal{X}, \|\cdot\|_{\#})$ with $f_j(z) = \sum_{n=0}^\infty a_n^{(j)} z^n$. By (9), $f_j \rightarrow f(z) = \sum_{n=1}^\infty a_n z^n$ in $(\mathcal{X}, \|\cdot\|_{\mathcal{X}})$. Then,

$$\|f_j - f\|_{\#} = \sup_{N \geq 0} \sup_{\|\lambda\|_{\ell^\infty} \leq 1} \sup_{\ell > M} \left\| \sum_{k=0}^N \lambda_k (a_k^{(j)} - a_k^{(\ell)}) z^k \right\|_{\mathcal{X}} \leq \sup_{\ell > M} \|f_j - f_\ell\|_{\mathcal{X}} < \epsilon$$

when M is large enough. So $f_j \rightarrow f$ in $(\mathcal{X}, \|\cdot\|_{\#})$. On the other hand, one can check directly that $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is a solid basis for $(\mathcal{X}, \|\cdot\|_{\#})$. (iv) \Rightarrow (ii) is clear. (ii) \Rightarrow (i) is also straightforward now since $\|f\|_{\#} \approx \|f\|_{\mathcal{X}}$, $f \in \mathcal{X}$, and one can verify directly that $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is a solid frame for \mathcal{X} . \square

Remark. An interesting fact from [33, p. 376] allows us to establish a connection between solid frames and Banach algebras. If \mathcal{X} is a functional Banach space of homogeneous type, and $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is a solid frame, then $\mathcal{X}_{\mathcal{P}}$ and \mathcal{X}^{\max} become Banach algebras when equipped with the norm $\|f\|_{\#} = \sup_{N \geq 0} \sup_{\|\lambda\|_{\ell^\infty} \leq 1} \left\| \sum_{n=0}^N \lambda_n a_n z^n \right\|_{\mathcal{X}}$, with the product given by $(f \star g)(z) = \sum_{n=0}^{\infty} a_n b_n \|z^n\|_{\mathcal{X}} z^n$. Then, $(\mathcal{X}_{\mathcal{P}}, \|\cdot\|_{\#}, \star)$ is a commutative Banach algebra without identity. The space $(\mathcal{X}^{\max}, \|\cdot\|_{\#}, \star)$ is also a commutative Banach algebra. Moreover, under the assumption that $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is a solid frame, the followings are equivalent:

- (i) $(\mathcal{X}^{\max}, \|\cdot\|_{\#}, \star)$ has an identity;
- (ii) $\sum_{n=0}^{\infty} \frac{z^n}{\|z^n\|_{\mathcal{X}}} \in \mathcal{X}^{\max}$;
- (iii) \mathcal{X}^{\max} and ℓ^∞ are isomorphic.

8. A characterization for monomials to be solid frames

Proof of Theorem F. We shall only prove the Banach space case. We first consider $\mathcal{X} = \mathcal{X}^{\max}$. If $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 1}$ is a solid frame, then $\sup_{N \geq 0} \|S_N\| < \infty$. Take any $\lambda = \{\lambda_n\}_{n \geq 0} \in \ell^\infty$, and $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}^{\max}$. Let $g(z) = \sum_{n=0}^{\infty} \lambda_n a_n z^n$. For any $0 \leq r < 1$, $\left\| \sum_{k=N}^M \lambda_k a_k r^k z^k \right\|_{\mathcal{X}} \lesssim \|\lambda\|_{\ell^\infty} \left\| \sum_{k=N}^M a_k r^k z^k \right\|_{\mathcal{X}}$. Hence, $g_r \in \mathcal{X}_{\mathcal{P}}$. So

$$\|g\|_{\max} = \sup_{0 \leq r < 1} \|g_r\|_{\mathcal{X}} \lesssim \|\lambda\|_{\ell^\infty} \sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} = \|\lambda\|_{\ell^\infty} \|f\|_{\max}.$$

It follows $g \in \mathcal{X}^{\max}$. For sufficiency, by Theorem E, it suffices to show that

$$\sup_{N \geq 0} \sup_{\|\lambda\|_{\ell^\infty} \leq 1} \left\| \sum_{k=0}^N \lambda_k a_k z^k \right\|_{\mathcal{X}} < \infty \tag{10}$$

for each $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{X}^{\max}$. Let $T_f : \ell^\infty \rightarrow \mathcal{X}^{\max}$ be defined by $T_f(\lambda) = \sum_{n=0}^{\infty} \lambda_n a_n z^n$. The condition that \mathcal{X}^{\max} is a solid space ensures that T_f is well defined. The closed graph theorem and the condition $\sup_{N \geq 0} \|S_N\| < \infty$ imply that (10) holds. Now

we treat the case $\mathcal{X} = \mathcal{X}_p$. By [43, Proposition 1], $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is a Schauder basis for \mathcal{X} if and only if $\sup_{N \geq 0} \|S_N\| < \infty$. The proof of the sufficiency for \mathcal{X}^{\max} above can be modified to show that $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is an unconditional basis for \mathcal{X} . Conversely, if it is a solid frame, then by Proposition 14, it is also an unconditional basis for \mathcal{X} . \square

Remark. For a functional Banach space of homogeneous type, the condition $\sup_{N \geq 0} \|S_N\| < \infty$ does not imply that $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ is a solid frame for \mathcal{X} . An example is provided by $\mathcal{X} = H^p$ ($1 < p < 2$) for which $\{z^n\}_{n \geq 0}$ is a Schauder basis. If it were a solid frame, then it would be an unconditional basis as well. This is known to be false and can be shown quickly. Otherwise, take $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H^p \setminus H^2$. By Littlewood’s theorem [29], there exists a sequence $\{\theta_n\}_{n \geq 0} \in \{-1, 1\}^{\mathbb{N}}$ such that $\sum_{n=0}^{\infty} \theta_n a_n z^n \notin H^p$, contradicting the assumption that $\{z^n\}_{n \geq 0}$ were an unconditional basis.

9. The problem of a solid span

Proof of Theorem G. (i) Let

$$E^{\text{s-max}} = \left\{ a = (a_1, a_2, \dots) : \|a\|_{\text{s-max}} = \sup_{N \geq 1} \left\| \sum_{n=1}^N a_n e_n \right\|_E < \infty \right\}. \tag{11}$$

Then $I(x) = ((\mathcal{T}x)_1, (\mathcal{T}x)_2, \dots)$ is an isometric embedding from E into $E^{\text{s-max}}$, where $\mathcal{T} : E \rightarrow \mathfrak{S}$ is the coefficient map. Note that $Ie_n = \mathbf{e}_n, n \geq 1$. For $|\lambda_n| \leq 1$ and $c_n \in \mathbb{C}$,

$$\left\| \sum_{n=1}^N \lambda_n c_n \mathbf{e}_n \right\|_{\text{s-max}} = \left\| \sum_{n=1}^N \lambda_n c_n e_n \right\|_E \leq \left\| \sum_{n=1}^N c_n e_n \right\|_E = \left\| \sum_{n=1}^N c_n \mathbf{e}_n \right\|_{\text{s-max}}.$$

Also $\lim_{N \rightarrow \infty} \|P_N a\|_{\text{s-max}} = \lim_{N \rightarrow \infty} \left\| \sum_{n=1}^N a_n e_n \right\|_E = \|a\|_{\text{s-max}}$. So $\{\mathbf{e}_n\}_{n \geq 1}$ is a solid basis for $E^{\text{s-max}}$. To complete the proof, it suffices to show that F can be isometrically embedded into $E^{\text{s-max}}$ if F is a solid extension of E (see Definition 5). Let $I' : E \hookrightarrow F$ denote the isometric embedding from E into F , with $\{I'e_n\}_{n \geq 1}$ being a solid basis for F . For convenience, write $e'_n = I'e_n, n \geq 1$, and $\mathcal{T}' : F \rightarrow \mathfrak{S}$ the corresponding coefficient map. We claim that $I'' : F \rightarrow E^{\text{s-max}}$ defined by $I''(x) = ((\mathcal{T}'x)_1, (\mathcal{T}'x)_2, \dots), x \in F$, is an isometric embedding. Note that

$$\left\| \sum_{n=1}^N (\mathcal{T}'x)_n e'_n \right\|_F = \left\| I' \left(\sum_{n=1}^N (\mathcal{T}'x)_n e_n \right) \right\|_F = \left\| \sum_{n=1}^N (\mathcal{T}'x)_n e_n \right\|_E$$

since $I' : E \hookrightarrow F$ is isometric. Then, for $x \in F$,

$$\|x\|_F = \lim_{N \rightarrow \infty} \left\| \sum_{n=1}^N (\mathcal{T}'x)_n e'_n \right\|_F = \lim_{N \rightarrow \infty} \left\| \sum_{n=1}^N (\mathcal{T}'x)_n e_n \right\|_E = \|I''(x)\|_{\text{s-max}}.$$

(ii) If $\{ve_n\}_{n \geq 1}$ is a solid basis for W . Let $\mathcal{T} : W \rightarrow \mathfrak{S}$ be the corresponding coefficient map. Then, we can check that \mathcal{T} is an isometric embedding from W into $(E_{\mathcal{P}})^{s\text{-max}}$. Conversely, $\{e_n\}_{n \geq 1}$ is a solid basis for $(E_{\mathcal{P}})^{s\text{-max}}$, since $\{e_n\}_{n \geq 1}$ is a solid basis for $E_{\mathcal{P}}$. Then, one can check that $\{ve_n\}_{n \geq 1}$ is a solid basis for W , since $\iota : E_{\mathcal{P}} \rightarrow W$ is an isometric embedding and $E_{\mathcal{P}}$ is embedded into $(E_{\mathcal{P}})^{s\text{-max}}$ isometrically. \square

10. The role of c_0

Proof of Theorem H. (i) We first show that $\mathcal{P} \subset c_0 \cdot E \subset E_{\mathcal{P}}$, which follows from Lemma 18 below. Here, \mathcal{P} denotes the collection of finite linear combinations of $\{e_n\}_{n \geq 1}$. \square

Lemma 18. *Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$. If $\{t_n\}_{n \geq 1} \in c_0$ and $x \in E$, then the series $\sum_{n=1}^{\infty} t_n(\mathcal{T}x)_n e_n$ converges in norm in E .*

Proof. By (8), $\sum_{n=1}^{\infty} |(\mathcal{T}x)_n| |\phi(e_n)| < \infty$, $\phi \in E^*$. Then, for $x \in E$, the mapping $T : E^* \rightarrow \ell^1$ defined by $T\phi \doteq \{(\mathcal{T}x)_n \phi(e_n)\}_{n \geq 1}$, $\phi \in E^*$ is continuous by the closed graph theorem. Now, for $m \leq k \leq n$,

$$\left\| \sum_{k=m}^n t_k(\mathcal{T}x)_k e_k \right\|_E \leq \sup_{\phi \in B_{E^*}} \left(\left(\sup_{m \leq k \leq n} |t_k| \right) \sum_{k=m}^n |(\mathcal{T}x)_k \phi(e_k)| \right) \lesssim \sup_{m \leq k \leq n} |t_k| \rightarrow 0$$

as $n > m \rightarrow \infty$, where B_{E^*} denotes the unit ball of E^* . \square

Next, we show that $c_0 \cdot E$ is closed in norm. Let $x_m \rightarrow x$ with $x_m \in c_0 \cdot E$. Then, by Lemma 18, $x_m \in E_{\mathcal{P}}$ and $x \in E_{\mathcal{P}}$. Write $x_m = \sum_{n=1}^{\infty} t_n^{(m)}(\mathcal{T}x_m)_n e_n$ with $\{t_n^{(m)}\}_{n \geq 1} \in c_0$. Since $\{(\mathcal{T}x_m)_n\}_{n \geq 1} \in \ell^{\infty}$, one has $\{t_n^{(m)}(\mathcal{T}x_m)_n\}_{n \geq 1} \in c_0$. Since $\{e_n\}_{n \geq 1}$ is a solid basis, for fixed n , $t_n^{(m)}(\mathcal{T}x_m)_n \rightarrow a_n$ as $m \rightarrow \infty$ for some $a_n \in \mathbb{C}$. This implies that $(\mathcal{T}x)_n = a_n$, $n \geq 1$. Moreover,

$$|a_n| = |a_n - t_n^{(m)}(\mathcal{T}x_m)_n + t_n^{(m)}(\mathcal{T}x_m)_n| \leq \|x_m - x\|_E + |t_n^{(m)}(\mathcal{T}x_m)_n|,$$

so $\{a_n\}_{n \geq 1} \in c_0$ and $x = \sum_{n=1}^{\infty} a_n e_n$ converges in norm. Then, there exists a subsequence of integers $\{n_k\}_{k \geq 1}$ such that $\left\| \sum_{i=n_k}^{n_{k+1}-1} a_i e_i \right\|_E < \frac{1}{k^3}$. Take $t_n = \frac{1}{\sqrt{k}}$ if $n \in [n_k, n_{k+1} - 1]$ and let $b_n = \frac{a_n}{t_n}$. Then, by the contraction principle, $x' = \sum_{n=1}^{\infty} b_n e_n \in E_{\mathcal{P}}$. Hence, $x = \sum_{n=1}^{\infty} t_n b_n e_n \in c_0 \cdot E$, as desired.

(ii) If $a \in E^{s\text{-max}}$, then (8) holds and Lemma 18 applies to $E^{s\text{-max}}$, which implies the sufficiency. Now the necessity. For $\{t_n\}_{n \geq 1} \in c_0$, $\sum_{n=1}^{\infty} t_n a_n e_n \in E_{\mathcal{P}}$ implies that this

series converges, so $\sup_{N \geq 1} \left\| \sum_{n=1}^N t_n a_n e_n \right\|_E < \infty$. This yields that $\sup_{N \geq 1} \left\| \sum_{n=1}^N a_n e_n \right\|_E < \infty$.

Otherwise, $\sup_{N \geq 1} \left\| \sum_{n=1}^N a_n e_n \right\|_E = \infty$, and there exists a subsequence $\{n_k\}_{k \geq 1}$ such that

$$\left\| \sum_{i=1}^{n_{k+1}} a_i e_i \right\|_E - \left\| \sum_{i=1}^{n_k} a_i e_i \right\|_E \geq k^2. \text{ So}$$

$$\left\| \sum_{i=n_k+1}^{n_{k+1}} a_i e_i \right\|_E \geq \left\| \sum_{i=1}^{n_{k+1}} a_i e_i \right\|_E - \left\| \sum_{i=1}^{n_k} a_i e_i \right\|_E \geq k^2.$$

If we take $t_i = \frac{1}{k}$ when $i \in [n_k + 1, n_{k+1}]$, then $t = \{t_n\}_{n \geq 1} \in c_0$, and $\left\| \sum_{i=n_k+1}^{n_{k+1}} t_i a_i e_i \right\|_E \geq$

k . By $\left\| \sum_{i=1}^{n_{k+1}} t_i a_i e_i \right\|_E \geq \left\| \sum_{i=n_k+1}^{n_{k+1}} t_i a_i e_i \right\|_E \geq k$, we conclude that $\sup_{N \geq 1} \left\| \sum_{n=1}^N a_n t_n e_n \right\|_E =$

∞ , a contradiction.

Now, we come to the (very short) proof of Theorem I, which is an application of James' theorem.

Proof of Theorem I. If $\{e_n\}_{n \geq 1}$ is an unconditional basis for $E^{s\text{-max}}$, then, by James' theorem [25, Theorem V.4, p. 98], $E^{s\text{-max}}$ does not contain c_0 . Conversely, we show that $E^{s\text{-max}} = (E^{s\text{-max}})_{\mathcal{P}}$. Otherwise, there exists a nonzero $a \in E^{s\text{-max}}$ such that $P_N a \not\rightarrow a$ as $N \rightarrow \infty$. But $\sup_{N \geq 1} \|P_N a\|_{s\text{-max}} = \|a\|_{s\text{-max}} < \infty$. So the conclusion follows from the Bessaga-Pełczyński theorem ([25, Theorem IV.2, p. 94]). □

Corollary 19. *Let \mathcal{X} be a functional Banach space of homogeneous type with $\{\frac{z^n}{\|z^n\|_{\mathcal{X}}}\}_{n \geq 0}$ as a solid basis. For $f \in H(\mathbb{D})$, we have $c_0 \cdot f \subset \mathcal{X}_{\mathcal{P}} \iff f \in \mathcal{X}^{max}$.*

Remark. By Proposition 15, we have $\mathcal{X}^{max} = \mathcal{X}_T$ in the above corollary.

The rest of this section is devoted to a characterization of $E_{\mathcal{P}}$ via compact operators, which is motivated by [25, Proposition II.6, p. 88].

Theorem 20. *Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$, and $x \in E$. Then $x \in E_{\mathcal{P}}$ if and only if the operator*

$$\begin{aligned} U : c_0 &\longrightarrow E \\ (t_n)_{n \geq 1} &\longmapsto \sum_{n=1}^{\infty} t_n (\mathcal{T}x)_n e_n \end{aligned}$$

is compact.

Combining Theorem H and Theorem 20, we have

Corollary 21. *Let E be a Banach space with a solid basis $\{e_n\}_{n \geq 1}$, and $x \in E$. The following are equivalent:*

- (i) $x \in E_{\mathcal{P}}$;
- (ii) there exists $\{t_n\}_{n \geq 1} \in c_0$ and $x' \in E$ such that $x = \sum_{n=1}^{\infty} t_n (\mathcal{T}x')_n e_n$;
- (iii) there exists $\{t_n\}_{n \geq 1} \in c_0$ and $x' \in E_{\mathcal{P}}$ such that $x = \sum_{n=1}^{\infty} t_n (\mathcal{T}x')_n e_n$;
- (iv) the operator $U : c_0 \rightarrow E$ defined by $(t_n)_{n \geq 1} \mapsto \sum_{n=1}^{\infty} t_n (\mathcal{T}x)_n e_n$ is compact.

Proof of Theorem 20. The operator U is well defined by Lemma 18. Also note that

$$\left\| \sum_{k=1}^n t_k (\mathcal{T}x)_k e_k \right\|_E \leq \sup_{k \geq 1} |t_k| \left\| \sum_{k=1}^n (\mathcal{T}x)_k e_k \right\|_E.$$

Letting $n \rightarrow \infty$, we see that $\|Ut\|_E \leq \|x\|_E \|t\|_{c_0}$; hence U is bounded.

Necessity: Let $x \in E_{\mathcal{P}}$. Note that

$$V = U^* : E^* \longrightarrow \ell^1$$

$$\phi \longmapsto \{(\mathcal{T}x)_n \phi(e_n)\}_{n \geq 1}.$$

We show that V is compact. Given any sequence $\{\phi_k\}_{k \geq 1} \subset B_{E^*}$, we must extract, from $\{V\phi_k\}_{k \geq 1}$, a subsequence convergent in norm in ℓ^1 . Schur’s theorem [25, Theorem III.10, p. 65] implies that it suffices to show that there exists a w -convergent subsequence. We may assume that E is separable by replacing, if necessary, E with $\overline{\text{span}}\{e_n : n \geq 1\}$. Then the compact space (B_{E^*}, w^*) is metrizable. Thus, we can extract a subsequence $\{\phi_{k_j}\}_{j \geq 1}$ w^* -convergent to $\phi \in B_{E^*}$.

Now we show that $V\phi_{k_j}$ converges weakly to $V\phi$. Let $\chi_{\sigma} \in \ell^{\infty}$ with $\sigma \subset \mathbb{N}$. Then,

$$\begin{aligned} \chi_{\sigma}(V\phi_{k_j}) &= \sum_{n \in \sigma} (V\phi_{k_j})(n) = \sum_{n \in \sigma} (\mathcal{T}x)_n \phi_{k_j}(e_n) \\ &= \phi_{k_j} \left(\sum_{n \in \sigma} (\mathcal{T}x)_n e_n \right) \rightarrow \phi \left(\sum_{n \in \sigma} (\mathcal{T}x)_n e_n \right) = \chi_{\sigma}(V\phi) \end{aligned}$$

as $j \rightarrow \infty$. The series $\sum_{n \in \sigma} (\mathcal{T}x)_n e_n$ converges since $x \in E_{\mathcal{P}}$ and $\{e_n\}_{n \geq 1}$ is an unconditional basis for $E_{\mathcal{P}}$. Observe that $\overline{\text{span}}\{\chi_{\sigma} : \sigma \subset \mathbb{N}\} = \ell^{\infty}$ and the sequence $\{V\phi_{k_j}\}_{j \geq 1}$ is bounded. It follows the desired convergence of $V\phi_{k_j}$.

Sufficiency: Let $Q_n : \ell^1 \rightarrow \ell^1$ be a natural projection such that $Q_n(\alpha_1, \alpha_2, \dots) = (0, \dots, 0, \alpha_{n+1}, \dots)$, where $\alpha = (\alpha_1, \alpha_2, \dots) \in \ell^1$. Then $\lim_{n \rightarrow \infty} Q_n(\alpha) = 0$ for $\alpha \in \ell^1$. Since the operator V is compact and the set $K = \overline{V(B_{E^*})}$ is compact, Q_n converges to 0 uniformly in K . It follows that $\{\sum_{n=1}^N (\mathcal{T}x)_n e_n\}_{N \geq 1}$ is a Cauchy sequence in E , since

$$\left\| \sum_{n=p}^q (\mathcal{T}x)_n e_n \right\|_E \leq \sup_{\phi \in B_{E^*}} \sum_{n=p}^q |(\mathcal{T}x)_n| |\phi(e_n)| = \sup_{\phi \in B_{E^*}} \|(Q_q - Q_p)(V\phi)\|_{\ell^1}.$$

This implies that $\sum_{n=1}^{\infty} (\mathcal{T}x)_n e_n$ converges in E , and $x \in E_{\mathcal{P}}$, as desired. □

11. A problem of A.E. Taylor

A common assumption imposed on Banach spaces of analytic functions over the unit disk is to ask

$$\sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} \leq C \|f\|_{\mathcal{X}}, \quad f \in \mathcal{X} \quad (12)$$

for some constant $C > 0$. Here, we essentially repeat a problem A. E. Taylor; see [41, p. 151].

Problem C. *Does there exist a functional Banach or p -Banach space \mathcal{X} such that it satisfies the conditions of Definition 2 except for (iv), and there exists a constant $C > 1$ such that (12) holds but fails when $C = 1$?*

Let \mathcal{X} be a functional Banach space which satisfies the conditions of Definition 2 except for (iv). Then, for each $f \in \mathcal{X}_{\mathcal{P}}$, $e^{i\theta} \mapsto f(\cdot e^{i\theta})$ is Bochner measurable from $\mathbb{T} \rightarrow \mathcal{X}$; see [8, Theorem 2.3]. Combining the Poisson integral expression for f_r and the vectorial Jensen inequality [20, Proposition 1.2.11, p. 19], one can prove the following:

Lemma 22. *Let \mathcal{X} be a functional Banach space which satisfies the conditions of Definition 2 except for (iv). Then, for each $f \in \mathcal{X}_{\mathcal{P}}$, we have*

- (i) $r \mapsto \|f_r\|_{\mathcal{X}}$ is increasing for $r \in (0, 1)$; and
- (ii) $\lim_{r \rightarrow 1^-} \|f_r\|_{\mathcal{X}} \leq \|f\|_{\mathcal{X}}$.

This prompts us to raise the following question:

Problem D. *Does there exist a functional Banach or p -Banach space \mathcal{X} which satisfies the conditions of Definition 2 except for (iv), and there exists some function $f \in \mathcal{X}$ such that $\sup_{0 \leq r < 1} \|f_r\|_{\mathcal{X}} < \|f\|_{\mathcal{X}}$?*

Acknowledgement. G. Cheng is supported by NSFC (12371126). X. Fang is supported by NSTC of Taiwan (NSTC 112-2115-M-008-010-MY2). C. Liu is supported by NSFC (12101103). Y. Lu is supported by NSFC (12031002).

References

- (1) F. Albiac, J. L. Ansorena, P. M. Berná and P. Wojtaszczyk, Greedy approximation for biorthogonal systems in quasi-Banach spaces, *Dissertationes Math.* **560** (2021), 1–88.
- (2) J. M. Anderson, J. Clunie and C. Pommerenke, On Bloch functions and normal functions, *J. Reine Angew. Math.* **270** (1974), 12–37.
- (3) J. M. Anderson and A. L. Shields, Coefficient multipliers of Bloch functions, *Trans. Amer. Math. Soc.* **224**(2) (1976), 255–265.
- (4) W. Arendt and N. Nikolski, Vector-valued holomorphic functions revisited, *Math. Z.* **234**(4) (2000), 777–805.
- (5) G. Bennett, D. A. Stegenga and R. M. Timoney, Coefficients of Bloch and Lipschitz functions, *Illinois J. Math.* **25**(3) (1981), 520–531.

- (6) P. Billard, Séries de Fourier aléatoirement bornées, continues, uniformément convergentes, *Studia Math.* **22** (1963), 309–329.
- (7) O. Blasco and M. Pavlović, Coefficient multipliers on Banach spaces of analytic functions, *Rev. Mat. Iberoam.* **27**(2) (2011), 415–447.
- (8) O. Blasco and G. Stylogiannis, Lipschitz-type conditions on homogeneous Banach spaces of analytic functions, *J. Math. Anal. Appl.* **445**(1) (2017), 612–630.
- (9) L. Brown and A. L. Shields, Cyclic vectors in the Dirichlet space, *Trans. Amer. Math. Soc.* **285**(1) (1984), 269–303.
- (10) G. Cheng, X. Fang, K. Guo, and C. Liu, A Gaussian version of Littlewood’s theorem for random power series, *Proc. Amer. Math. Soc.* **150**(8) (2022), 3525–3536.
- (11) G. Cheng, X. Fang, and C. Liu, A Littlewood-type theorem for random Bergman functions, *Int. Math. Res. Not. IMRN* **2022**(14) (2022), 11056–11091.
- (12) G. Cheng, X. Fang, C. Liu and Y. Lu, Random Dirichlet multipliers and Littlewood-type theorems, (2021), preprint.
- (13) W. G. Cochran, J. H. Shapiro and D. C. Ullrich, Random Dirichlet functions: multipliers and smoothness, *Canad. J. Math.* **45**(2) (1993), 255–268.
- (14) J. B. Conway, *A Course in Functional Analysis. Graduate Texts in Mathematics*, second, Volume 96, (Springer-Verlag, New York, 1990).
- (15) J. Diestel, *Sequences and Series in Banach Spaces. Graduate Texts in Mathematics*, Volume 92, (Springer-Verlag, New York, 1984).
- (16) P. Duren, Random series and bounded mean oscillation, *Michigan Math. J.* **32**(1) (1985), 81–86.
- (17) P. L. Duren, *Theory of H^p spaces. Pure and Applied Mathematics*, Volume 38, (Academic Press, New York-London, 1970).
- (18) F. Gao, A characterization of random Bloch functions, *J. Math. Anal. Appl.* **252**(2) (2000), 959–966.
- (19) H. Hedenmalm, B. Korenblum and K. Zhu, *Theory of Bergman Spaces. Graduate Texts in Mathematics*, Volume 199, (Springer-Verlag, New York, 2000).
- (20) T. Hytönen, J. van Neerven, M. Veraar and L. Weis, *Analysis in Banach Spaces. Vol. I. Martingales and Littlewood-Paley Theory. Ergebnisse der Mathematik und Ihrer Grenzgebiete. 3. Folge*, Volume 63, (Springer, Cham, 2016).
- (21) T. Hytönen, J. van Neerven, M. Veraar and L. Weis, *Analysis in Banach Spaces. Vol. II. Probabilistic Methods and Operator Theory, Ergebnisse der Mathematik und Ihrer Grenzgebiete. 3. Folge*, Volume 67, (Springer, Cham, 2017).
- (22) J. -P. Kahane, *Some Random Series of Functions. Of Cambridge Studies in Advanced Mathematics*, Volume **5**, (Cambridge University Press, Cambridge, second ed, 1985).
- (23) N. J. Kalton, Rademacher series and decoupling, *New York J. Math.* **11** (2005), 563–595.
- (24) S. Konyagin, H. Queffélec, E. Saksman, and K. Seip, Riesz projection and bounded mean oscillation for Dirichlet series, *Studia Math.* **262**(2) (2022), 121–149.
- (25) D. Li and H. Queffélec, *Introduction to Banach Spaces: Analysis and Probability. Vol. 1. Cambridge Studies in Advanced Mathematics*, Volume 1, (Cambridge University Press, Cambridge, 2018) :66.
- (26) D. Li and H. Queffélec, *Introduction to Banach Spaces: Analysis and Probability. Vol. 2*, Cambridge Studies in Advanced Mathematics, Volume 167, (Cambridge University Press, Cambridge, 2018).
- (27) J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces. I, Sequence Spaces. Ergebnisse der Mathematik und Ihrer Grenzgebiete*, Volume 92, (Springer-Verlag, Berlin-New York, 1977).
- (28) J. E. Littlewood, On the mean values of power series, *J. London Math. Soc.* **2**(25) (1926), 328–337.

- (29) J. E. Littlewood, On the mean values of power series (II), *J. London Math. Soc.* **5**(3) (1930), 179–182.
- (30) C. Liu, Multipliers for Dirichlet type spaces by randomization, *Banach J. Math. Anal.* **14**(3) (2020), 935–949.
- (31) M. B. Marcus and G. Pisier, *Necessary and Sufficient Conditions for the Uniform Convergence of Random Trigonometric Series. Of Lecture Notes Series*, Volume **50**, (Aarhus Universitet, Matematisk Institut, Aarhus, 1978).
- (32) M. B. Marcus and G. Pisier, *Random Fourier Series with Applications to Harmonic Analysis. Annals of Mathematics Studies*, Volume **101**, (University of Tokyo Press, Tokyo, 1981).
- (33) R. E. Megginson, *An Introduction to Banach Space Theory. Graduate Texts in Mathematics*, Volume **183**, (Springer-Verlag, New York, 1998).
- (34) A. Nishry and E. Paquette, Gaussian analytic functions of bounded mean oscillation, *Anal. PDE* **16**(1) (2023), 89–117.
- (35) R. E. A. C. Paley and A. Zygmund. On some series of functions (1), *Proc. Camb. Phil. Soc.*, **26** (1930), 337–357.
- (36) R. E. A. C. Paley & A. Zygmund. On some series of functions (2), *Proc. Camb. Phil. Soc.*, **26** (1930), 458–474.
- (37) W. Rudin, *Functional Analysis. International Series in Pure and Applied Mathematics*, second, (McGraw-Hill, Inc, New York, 1991).
- (38) R. Salem and A. Zygmund, *Some properties of trigonometric series whose terms have random signs*, *Acta Math.* **91** (1954), 245–301.
- (39) W. T. Sledd, Random series which are BMO or Bloch, *Michigan Math. J.* **28**(3) (1981), 259–266.
- (40) W. T. Sledd, and D. A. Stegenga, An H^1 multiplier theorem, *Ark. Mat.* **19**(2) (1981), 265–270.
- (41) A. E. Taylor, Banach spaces of functions analytic in the unit circle I, *Studia Math.* **11** (1950), 145–170.
- (42) H. Wulan, Random power series with bounded mean oscillation characteristics, *Acta Math. Sci. (Chinese)* **14**(4) (1994), 451–457.
- (43) K. Zhu, Duality of Bloch spaces and norm convergence of Taylor series, *Michigan Math. J.* **38**(1) (1991), 89–101.