

Curcumin attenuates insulin resistance and hepatic lipid accumulation in a rat model of intra-uterine growth restriction through insulin signalling pathway and sterol regulatory element binding proteins

Yu Niu¹, Jintian He¹, Hussain Ahmad², Chao Wang¹, Xiang Zhong¹, Lili Zhang¹, Tao Cui¹, Jingfei Zhang¹ and Tian Wang^{1*}

¹College of Animal Science and Technology, Nanjing Agricultural University, Nanjing 210095, People's Republic of China

²Animal Nutrition Department, University College of Veterinary and Animal Sciences, The Islamia University Bahawalpur, 63100, Pakistan

(Submitted 16 February 2019 – Final revision received 27 May 2019 – Accepted 14 June 2019; First published online 25 June 2019)

Abstract

The objective of the present study was to investigate the effect of curcumin on insulin resistance (IR) and hepatic lipid accumulation in intra-uterine growth restriction (IUGR). Rats with a normal birth weight (NBW) or IUGR were fed basic diets (NBW and IUGR groups) or basic diets supplemented with curcumin (NBW-C and IUGR-C groups) from 6 to 12 weeks. Rats in the IUGR group showed higher levels of glucose and homeostasis model assessment for insulin resistance index (HOMA-IR) ($P < 0.05$) than in the NBW group. The livers of IUGR rats exhibited higher ($P < 0.05$) concentration of TAG and lower ($P < 0.05$) activities of lipolysis enzymes compared with the normal rats. In response to dietary curcumin supplementation, concentrations of serum insulin, glucose and HOMA-IR, pyruvate, TAG, total cholesterol and NEFA in the liver were decreased ($P < 0.05$). The concentrations of glycogen and activities of lipolysis enzymes in the liver were increased ($P < 0.05$) in the IUGR-C group compared with the IUGR group. These results were associated with lower ($P < 0.05$) phosphorylated insulin receptor substrate 1, protein kinase B or Akt, glycogen synthase kinase 3 β and expressions of sterol regulatory element binding protein 1 and fatty acid synthase (FASN); decreased expressions for *Cd36*, sterol regulatory element binding protein 1c (*Srebf1*) and *Fasn*; increased ($P < 0.05$) expression of PPAR α ; and expressions for *Ppara* and hormone-sensitive lipase in the liver of IUGR-C rats than the IUGR rats. Maternal malnutrition caused IR and lipid accumulation in the liver. Curcumin supplementation prevented IR by regulating insulin signalling pathways and attenuated hepatic lipid accumulation.

Key words: Intra-uterine growth restriction: Insulin resistance: Lipid metabolism: Curcumin: Rats

Intra-uterine growth restriction (IUGR) is a serious problem in human beings and mammals, which affects approximately 5–10% of human infants worldwide and 15–20% of newborn animals^(1,2). It has been observed that at least 13.7 million infants were born with low birth weight each year⁽³⁾. IUGR leads to high morbidity and mortality, low feed utilisation and permanent adverse effects on postnatal life. Barker *et al.*⁽⁴⁾ originally stated the ‘early’ or ‘fetal’ origins of adult disease hypothesis. The hypothesis described that environmental factors, particularly poor nutrition, act in early life to programme the risks for the early onset of metabolic diseases in adult life and premature death. A significant relationship was observed between children born with IUGR and the development

of a variety of adult diseases, including hyperinsulinaemia, dyslipidaemia and non-alcoholic fatty liver disease^(5,6).

Liver is a major organ for lipid metabolism and insulin target organ, often influenced by IUGR during pregnancy^(7,8). The liver also plays a critical role in maintaining blood glucose homeostasis by controlling hepatic glucose production. IUGR has been linked to glucose intolerance and insulin resistance (IR)⁽⁹⁾. Previous studies suggested that IUGR not only closely linked with lipid dysfunction but also related to fatty liver disease⁽¹⁰⁾.

Curcumin is a naturally occurring phenolic compound, which is widely used in food, beverage, medicine and so forth⁽¹¹⁾. Curcumin (C₂₁H₂₀O₆), first described in 1910 by Lampe and

Abbreviations: Akt, protein kinase B; FASN, fatty acid synthase; GSK, glycogen synthase kinase; HL, hepatic lipase; HOMA-IR, homeostasis model assessment for insulin resistance; HSL, hormone-sensitive lipase; IR, insulin resistance; IRS-1, insulin receptor substrate 1; IUGR, intra-uterine growth restriction; IUGR-C, intra-uterine growth restriction + curcumin; LPL, lipoprotein lipase; NBW, normal birth weight; NBW-C, normal birth weight + curcumin; PI3K, phosphoinositide 3-kinase; PK, pyruvate kinase; *Srebf1*, sterol regulatory element binding protein 1c; SREBP1, sterol regulatory element binding protein 1; TC, total cholesterol; TL, total lipase.

* **Corresponding author:** Tian Wang, fax +86 25 84396195, email tianwangnjau@163.com

Table 1. Primers

Primer	Sequence		Product size (bp)
	Forward	Reverse	
<i>Cd36</i>	TGTA CTCTCTCCTCGGATGG	GAGCACTTGCTTCTTGCCAAC	198
<i>Ppara</i>	TCGTGGAGTCCTGGA ACTGA	GTTACGCCCAAATGCACCAC	118
<i>Srebf1</i>	CCATGGACGAGCTACCCTTC	GGCATCAAATAGGCCAGGGA	149
<i>Fasn</i>	TATTGACCCTGCAACCCACC	TAGACACCACCAGAGACGGT	111
<i>HSL</i>	TCCGTGCTCTAGCCTACTAC	GTGAGTCTGGAAGGACATGG	189
<i>Actb</i>	CCCGCGAGTACAACCTTCTT	CGCAGCGATATCGTCATCCA	83

Srebf1, sterol regulatory element binding protein 1c; *Fasn*, fatty acid synthase; *HSL*, hormone-sensitive lipase; *Actb*, β -actin.

Milobedeska, is the most active ingredient of turmeric and makes up 2–5 % of this spice⁽¹²⁾. The beneficial effects of curcumin on IR have been widely researched in animal models⁽¹³⁾. Curcumin treatment also has been illustrated to attenuate IR by decreasing insulin receptor substrate 1 (IRS-1) phosphorylation in the muscle of Wistar rats fed with high fructose⁽¹⁴⁾. In addition, they found that curcumin attenuated hyperinsulinaemia and the homeostasis model assessment for IR (HOMA-IR) index. However, related studies about curcumin on IR in IUGR are very limited.

Accordingly, we hypothesised that dietary supplementation of curcumin has a protective effect on IUGR IR by modulating the insulin signalling pathway. Furthermore, we took advantage of a maternal malnutrition rat model, which is an identified model for human IUGR study, and determined the hepatic lipid content and protein expressions related to fatty acid synthesis and lipid oxidation, to investigate whether these changes of IUGR-induced hyperlipidaemia were alleviated after curcumin administration.

Materials and methods

Curcumin preparation

The curcumin used in the present study was kindly provided by the Kehu Bio-technology Research Center (Guangzhou, People's Republic of China). The content of curcumin was 98 % measured by using HPLC.

Animal experiment design

The experimental design and procedures were approved by the Institutional Animal Care and Use Committee of Nanjing Agricultural University following the requirements of the Regulations for the Administration of Affairs Concerning Experimental Animals of China (SYXK(Su)2017-0007). The feed restriction method was used for maternal rats during pregnancy lead to induction of IUGR mode according to a previous study⁽¹⁵⁾. First-time-pregnant Sprague–Dawley rats (Nanjing Qinglongshan Experiment Animal Center, Nanjing, People's Republic of China) were housed in a facility at a constant temperature and humidity. The light regimen was adjusted to a 12 h light–12 h dark cycle. At day 10 of gestation, rats were provided either a diet of standard laboratory diet (LabDiet 5001) (n 12) *ad libitum* or a 50 % feed-restricted (n 12) diet determined by the quantification of normal feed intake in *ad libitum*-fed rats. Dams gave birth normally, and birth weights of offspring were recorded on day 1 of postnatal

life and divided into normal or IUGR groups. Rats were limited to 10 per litter to normalise rearing (average 11.83 and 12.1 rats per litter in normal and IUGR groups, respectively). During the 21 d of lactation period, each litter of rats from the normal or IUGR group were nursed by their own dams. During the lactation, all dams were free to feed. At 3 weeks of age, offspring in each litter were weaned and housed individually until they were 6 weeks old for observing the early growth of rats. At 6 weeks of age, twenty-four female rats with nearly equal body weights (within each group) were allocated to the NBW (normal birth weight), NBW-C (NBW with curcumin supplementation), IUGR and IUGR-C (IUGR with curcumin supplementation) groups (one rat per litter, (n 6) per group), respectively. The rats were allowed water and a standard granulated diet (American Institute of Nutrition (AIN)-93G diet) *ad libitum*. During the entire experimental period, rats in the NBW-C and IUGR-C groups were fed a standard diet supplemented with 400 mg curcumin/kg. Curcumin was added to the feed before it was made into pellets. The light regimen was a 12 h light–12 h dark cycle and the temperature was maintained at 20 to 24°C. At 12 weeks of age, all rats were fasted overnight, and blood was collected via cardiac puncture after anaesthesia. Serum was obtained from the blood via centrifugation for 15 min at 3000 **g** at 4°C. Liver tissue (the same area for each sample) was removed after death and snap-frozen in liquid N₂ and then stored at –80°C for further analysis.

Serum biochemistry parameters

Concentrations of total cholesterol (TC) and TAG in serum and liver were measured according to previous studies^(16,17). Concentrations of VLDL in serum and liver were determined by ELISA kit from Shanghai YILI Biological Technology Co., Ltd. Concentrations of HDL-cholesterol, LDL-cholesterol, glycogen, pyruvate and NEFA and activities of hepatic pyruvate kinase (PK), hepatic lipase (HL) and lipoprotein lipase (LPL) were determined using colorimetric kits (Nanjing Jiancheng Institute of Bioengineering) with a spectrophotometer, according to the manufacturer's instructions. Total lipase (TL) activity was defined as equal to HL and LPL activities. The concentrations of curcumin in serum and liver of rats were determined by HPLC-MS/MS system according to the previous study⁽¹⁸⁾.

Gene expression assays

Total RNA from the liver samples stored at –80°C was isolated using the Trizol reagent (Invitrogen). The determination of RNA content, mRNA quantification and real-time PCR (Applied

Table 2. Effect of curcumin on the serum concentrations of insulin, glucose and homeostasis model of assessment for insulin resistance (HOMA-IR) index of rats with intra-uterine growth restriction (IUGR)* (Mean values and standard deviations; *n* 6 per group)

Items	Experimental groups								<i>P</i>		
	NBW		NBW-C		IUGR		IUGR-C				
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	B	D	B×D
Insulin (mU/l)	8.46 ^{a,b}	0.07	7.85 ^b	0.66	8.89 ^a	0.54	6.52 ^c	0.71	0.06	<0.01	<0.01
Glucose (mmol/l)	3.60 ^{b,c}	0.18	3.79 ^b	0.06	4.41 ^a	0.24	3.50 ^c	0.07	<0.01	<0.01	<0.01
HOMA-IR†	1.43 ^b	0.14	1.38 ^b	0.06	1.72 ^a	0.01	1.02 ^c	0.04	0.28	<0.01	<0.01

NBW, normal birth weight rats; NBW-C, normal birth weight rats fed diets supplemented with 400 mg/kg curcumin; IUGR, IUGR rats fed diets supplemented with 400 mg/kg curcumin; B, birth weight; D, dietary curcumin supplementation; B × D, interaction between the corresponding parameters.

^{a,b,c} Mean values within a row with unlike superscript letters were significantly different (*P* < 0.05).

* Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate.

† HOMA-IR = (fasting glucose (mmol/l) × fasting insulin (μU/ml))/22.5.

Biosystems) were performed according to previously described methods⁽⁹⁾. The primer sequences for the target and housekeeping genes (*Cd36*, *Ppara*, sterol regulatory element binding protein 1c (*Srebp1c*), fatty acid synthase (*Fasn*), hormone-sensitive lipase (*Hsl*) and β-actin (*Actb*)) used for real-time PCR are listed in Table 1. Glyceraldehyde 3-phosphate dehydrogenase (*Gapdh*) was also used as a control gene to normalise the expression of target genes. Briefly, a reaction system of 20 μl was composed of 0.4 μl of forward primers, 0.4 μl of reverse primers, 0.4 μl of ROX reference dye, 10 μl of SYBR Premix Ex Taq (TaKaRa Biotechnology Co. Ltd), 6.8 μl of double-distilled water and 2 μl of complementary DNA. The 2^{-ΔΔCt} method was used to calculate relative levels of mRNA expression after normalisation with housekeeping genes⁽¹⁹⁾. The values for the NBW group were used for calibration.

Protein analysis

Primary antibodies against IRS-1 (1:1000; Cell Signaling Technology); glycogen synthase kinase-3 α/β (GSK3α/β; 1:1000; Cell Signaling Technology); protein kinase B (Akt; 1:1000; Cell Signaling Technology); phosphoinositide 3-kinase p85 (PI3K; 1:1000; Cell Signaling Technology); sterol regulatory element binding protein 1 (SREBP1; 1:1000; Affinity); FASN (1:1000; Cell Signaling Technology); PPARα (1:500; Affinity); phosphorylated IRS-1^{Ser302} (pIRS-1^{Ser302}; 1:1000; Cell Signaling Technology); phosphorylated GSK3α^{Ser21}/β^{Ser9} (pGSK3α^{Ser21}/β^{Ser9}; 1:1000; Cell Signaling Technology); phosphorylated Akt^{Ser473} (pAkt^{Ser473}; 1:1000; Cell Signaling Technology); phosphorylated PI3K^{Tyr458} (pPI3K^{Tyr458}; 1:1000; Cell Signaling Technology); and antibodies against *Actb* (1:1000; Cell Signaling Technology) were used in the present study. The total protein and cytomembrane protein of the liver were extracted using assay kits according to the manufacturer's instructions (Beyotime). The protein content of the sample was measured using the bicinchoninic acid (BCA) Protein Assay Kit (Beyotime). For Western blotting analyses, 40 μg of protein from each sample was subjected to sodium dodecylsulfate-polyacrylamide gel electrophoresis. After electrophoresis, proteins were separated and transferred to polyvinylidene difluoride membranes. The membranes were blocked with blocking buffer (5 % non-fat dry milk) for 12 h at 4°C. Then, the membranes were probed with appropriate primary and

secondary antibodies (horseradish peroxidase-conjugated goat anti-rabbit or anti-mouse IgG; Cell Signaling Technology 1:5000 dilution in 1 × Tris-buffered saline with 0.1 % Tween 20). The blots were detected using enhanced chemiluminescence reagents (ECL-Kit, Beyotime) followed by autoradiography. Photographs of the membranes were taken using the Luminescent Image Analyzer LAS-4000 system (Fujifilm Co.) and quantified by Gel-Pro Analyzer 4.0 software (Media Cybernetics). Results were corrected for total protein.

Statistical analysis

Differences between groups were analysed using a two-way ANOVA. The classification variables were birth weight (NBW + NBW-C × IUGR + IUGR-C), diet (NBW + IUGR × NBW-C + IUGR-C), and the interaction between birth weight and diet (NBW × NBW-C × IUGR × IUGR-C). A Tukey's *post hoc* analysis was used to determine the differences between the four groups when a statistically significant birth weight × diet interaction was observed. SPSS 17.0 (SPSS, Inc.) was used for these analyses. A probability level of *P* < 0.05 was considered statistically significant, and *P* < 0.01 was considered very significant. Data are presented as mean values and standard deviations.

Results

Serum hormone levels

IUGR rats exhibited higher concentrations of serum insulin (*P* > 0.05), glucose (*P* < 0.05) and HOMA-IR (*P* < 0.05) compared with NBW rats. Curcumin supplementation reduced (*P* < 0.01) the concentrations of serum insulin, glucose and HOMA-IR in IUGR rats. In addition, a significant effect of birth weight × dietary interaction (*P* < 0.05) for concentrations of serum insulin, glucose and HOMA-IR were observed. There were no significant differences in the concentrations of serum insulin, glucose and HOMA-IR between rats of the NBW and NBW-C groups (*P* > 0.05) (Table 2).

Serum biochemistry and hepatic lipid metabolic parameters

The concentrations of TC and TAG in the serum of IUGR rats were significantly lower (*P* < 0.05) compared with the NBW rats. Dietary

Table 3. Effect of curcumin on serum and liver lipid metabolic measurements of rats with intra-uterine growth restriction (IUGR)* (Mean values and standard deviations; *n* 6 per group)

Items	Experimental groups								<i>P</i>		
	NBW		NBW-C		IUGR		IUGR-C				
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	B	D	B×D
Serum											
TC (mmol/l)	2.52 ^a	0.05	2.48 ^a	0.08	2.30 ^b	0.13	2.19 ^b	0.10	<0.01	0.07	0.33
TAG (mmol/l)	0.52 ^a	0.04	0.42 ^b	0.05	0.41 ^b	0.03	0.42 ^b	0.01	<0.01	<0.01	<0.01
HDL-cholesterol (mmol/l)	0.53 ^a	0.10	0.46 ^b	0.03	0.55 ^a	0.07	0.34 ^c	0.05	0.02	<0.01	<0.01
LDL-cholesterol (mmol/l)	1.32 ^b	0.27	1.64 ^a	0.26	1.55 ^b	0.36	1.58 ^a	0.23	0.30	0.04	0.08
VLDL-cholesterol (mmol/l)	11.41	1.51	10.95	0.81	11.74	0.83	10.72	0.49	0.89	0.08	0.50
Curcumin (ng/ml)	–	–	1.79	0.29	–	–	2.21	0.18	–	–	–
Liver											
TC (mmol/g protein)	1.48 ^b	0.40	1.97 ^{a,b}	0.66	2.49 ^a	0.70	1.78 ^b	0.25	0.08	0.62	<0.05
TAG (mmol/g protein)	1.81 ^b	0.22	1.55 ^b	0.18	2.59 ^a	0.29	1.72 ^b	0.39	<0.01	<0.01	0.01
NEFA (μmol/g protein)	294.44 ^c	12.63	269.52 ^d	18.43	341.18 ^a	1.64	320.35 ^b	31.99	<0.01	<0.01	0.80
VLDL-cholesterol (mmol/g protein)	1.10 ^a	0.11	0.93 ^b	0.15	1.10 ^a	0.06	1.07 ^a	0.04	0.10	0.10	0.02
Curcumin (μg/g)	–	–	2.73	0.24	–	–	2.68	0.15	–	–	–

NBW, normal birth weight rats; NBW-C, normal birth weight rats fed diets supplemented with 400 mg/kg curcumin; IUGR-C, IUGR rats fed diets supplemented with 400 mg/kg curcumin; B, birth weight; D, dietary curcumin supplementation; B × D, interaction between the corresponding parameters; TC, total cholesterol; –, means not determined.

^{a,b,c,d} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

* Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate.

Table 4. Effect of curcumin on hepatic glycogen and enzymes of rats with intra-uterine growth restriction (IUGR)* (Mean values and standard deviations; *n* 6 per group)

Items	Experimental groups								<i>P</i>		
	NBW		NBW-C		IUGR		IUGR-C				
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	B	D	B×D
Glycogen (mg/g protein)	5.36 ^a	0.98	3.56 ^c	0.47	1.87 ^d	0.12	4.28 ^b	0.99	<0.01	<0.01	0.32
Pyruvate (μmol/g protein)	13.21 ^b	0.84	13.82 ^b	1.03	17.33 ^a	1.28	13.22 ^b	1.52	<0.01	<0.01	<0.01
Pyruvate kinase (U/g protein)	14.77 ^a	0.76	11.23 ^c	1.87	13.00 ^b	0.97	8.78 ^d	1.31	<0.01	<0.01	0.53

NBW, normal birth weight rats; NBW-C, normal birth weight rats fed diets supplemented with 400 mg/kg curcumin; IUGR-C, IUGR rats fed diets supplemented with 400 mg/kg curcumin; B, birth weight; D, dietary curcumin supplementation; B × D, interaction between the corresponding parameters.

^{a,b,c,d} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

* Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate.

curcumin supplementation significantly decreased ($P < 0.05$) the concentration of HDL-cholesterol and increased ($P < 0.05$) the concentration of LDL-cholesterol in the serum of NBW and IUGR rats. In addition, a birth weight × dietary interaction effect ($P < 0.05$) was observed for concentrations of TAG and HDL-cholesterol in the serum. The concentrations of TC, TAG and NEFA in the liver of IUGR rats were significantly higher ($P < 0.05$) than in the liver of NBW rats. In the IUGR-C group, the concentrations of TC, TAG and NEFA in the liver of rats were lower ($P < 0.05$) than in the IUGR group. A birth weight × dietary interaction effect was observed in the concentrations of TC and TAG in the liver of rats. The NBW-C group had lower ($P < 0.05$) concentrations of VLDL and NEFA in the liver than in the NBW group rats (Table 3).

Hepatic glycogen and metabolic parameters

IUGR rats exhibited lower ($P < 0.05$) concentration of glycogen and higher ($P < 0.05$) concentration of pyruvate in the liver compared with the NBW rats. Dietary curcumin supplementation significantly increased the concentration of glycogen and decreased the concentration of pyruvate and activity of PK in the liver of IUGR rats ($P < 0.05$). Dietary curcumin supplementation, the activity of PK

and the concentration of glycogen were significantly lower in the liver of the NBW-C group than the NBW group ($P < 0.05$). In addition, a birth weight × dietary interaction effect ($P < 0.05$) was observed for concentrations of pyruvate (Table 4).

Hepatic lipolysis enzymes

The IUGR group showed lower activities of LPL ($P < 0.01$), HL ($P < 0.05$) and TL ($P < 0.05$) in the liver of rats compared with the NBW group rats. The activities of hepatic LPL ($P < 0.01$), HL ($P < 0.05$) and TL ($P < 0.01$) were higher ($P < 0.05$) in the IUGR-C group rats than in the IUGR group rats. Increased ($P < 0.05$) activities of hepatic LPL, HL and TL were observed in the NBW-C group rats (Table 5).

Hepatic protein expression

The IUGR rats exhibited higher phosphorylated IRS-1 ($P > 0.05$) (Fig. 1(A)), Akt ($P < 0.05$) (Fig. 1(C)) and glycogen synthase 3β (GSK3β) ($P < 0.05$) (Fig. 2) levels, and lower ($P > 0.05$) phosphorylated GSK3α and PI3K levels (Fig. 1(B)) in the liver compared with the NBW rats. Dietary curcumin supplementation significantly decreased ($P < 0.05$) the phosphorylated levels of IRS, Akt and

Table 5. Effect of curcumin on hepatic lipolysis enzymes of rats with intra-uterine growth restriction (IUGR)* (Mean values and standard deviations; *n* 6 per group)

Items	Experimental groups								<i>P</i>		
	NBW		NBW-C		IUGR		IUGR-C				
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	B	D	B×D
LPL (U/mg protein)	0.97 ^b	0.03	1.26 ^a	0.21	0.88 ^c	0.07	0.99 ^b	0.10	<0.01	<0.01	0.09
HL (U/mg protein)	1.05 ^b	0.14	1.18 ^a	0.17	0.92 ^c	0.13	1.06 ^b	0.10	0.04	0.03	0.93
TL (U/mg protein)	1.92 ^c	0.20	2.43 ^a	0.34	1.89 ^d	0.11	2.10 ^b	0.10	0.05	<0.01	0.09

NBW, normal birth weight rats; NBW-C, normal birth weight rats fed diets supplemented with 400 mg/kg curcumin; IUGR-C, IUGR rats fed diets supplemented with 400 mg/kg curcumin; B, birth weight; D, dietary curcumin supplementation; B × D, interaction between the corresponding parameters; I, IUGR; LPL, lipoprotein lipase; HL, hepatic lipase; TL, total lipase, equal to HL and LPL activities.

^{a,b,c,d} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

* Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate.




Fig. 1. Abundance of phosphorylated proteins in liver of normal birth weight (NBW) rats, NBW rats supplemented with curcumin (NBW-C), intra-uterine growth restriction (IUGR) rats and IUGR rats supplemented with curcumin (IUGR-C). (A) Insulin receptor substrate-1 (IRS-1); (B) phosphoinositide 3-kinase p85 (PI3K); (C) protein kinase B (Akt). Data are mean values (*n* 6 per group), with standard deviations represented by vertical bars. Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate. ^{a,b,c} Mean values with unlike letters were significantly different ($P < 0.05$). Results were corrected for total protein. B, birth weight; D, diet; B × D, interaction between the corresponding parameters. pIRS-1^{Ser302}, phosphorylated IRS-1^{Ser302}; pPI3K^{Tyr458}, phosphorylated phosphoinositide 3-kinase p85^{Tyr458}.

GSK3 β , and had a tendency to increase ($P > 0.05$) the phosphorylated levels of GSK3 α and PI3K in the liver of the IUGR-C group. A birth weight × dietary interaction effect ($P < 0.05$) was observed for the phosphorylated levels of IRS, GSK3 β and PI3K.

The protein expressions of SREBP1 (Fig. 3(A)) and FASN (Fig. 3(B)) were significantly increased ($P < 0.05$) in the liver of IUGR rats compared with the NBW rats. The protein expressions of SREBP1 and FASN in the liver of the IUGR-C group were lower ($P < 0.05$) than in the IUGR rats. The IUGR rats also showed lower ($P < 0.05$) protein expression of PPAR α in the liver compared with the NBW group rats (Fig. 3(C)). Dietary curcumin supplementation significantly increased ($P < 0.05$) the protein expression of PPAR α in the IUGR-C group rats. Dietary curcumin supplementation had no significant effects ($P > 0.05$) on the protein expressions of SREBP1, FASN and PPAR α in the liver of NBW rats. Additionally, a birth weight × dietary interaction effect ($P < 0.01$) was noted for expressions of SREBP1, FASN and PPAR α .

Gene expression

In the liver of the IUGR rats, the mRNA expression levels for *Cd36*, *Srebf1* and *Fasn* were higher ($P < 0.05$) and the mRNA expression levels for *Ppara* and *HSL* were lower ($P < 0.05$) than

in the liver of NBW rats. Dietary curcumin supplementation significantly increased ($P < 0.05$) the mRNA expression levels for *Ppara* and *HSL*, and decreased ($P < 0.05$) the mRNA expression levels for *Cd36*, *Srebf1* and *Fasn* in the liver of the IUGR-C group. A birth weight × dietary interaction effect was noted for the mRNA expressions for *Cd36*, *Ppara*, *Srebf1* and *HSL* in the liver of rats ($P < 0.01$) (Fig. 4).

Discussion

IUGR impairs liver metabolism during the early period in piglets⁽⁹⁾ and induces IR in rats⁽²⁰⁾. Previous studies of Magee *et al.*⁽¹⁰⁾ found that the IUGR male rats increased hepatic fatty synthase and TAG contents. Mina *et al.*⁽¹⁵⁾ found that IUGR female adult had higher per cent of body fat. Our previous study also found that curcumin was beneficial in preventing IUGR-induced inflammation, oxidative damage and injury in the liver of IUGR rats⁽²¹⁾. Nevertheless, the studies on IR and the relationship between glycogen and lipid metabolism in IUGR female rats are very limited. Therefore, we chose IUGR female rats as our research animal model and investigated the effect of dietary curcumin supplementation on insulin levels, hepatic lipid and glycogen metabolism of female rats.




Fig. 2. Abundance of phosphorylated glycogen synthesis kinase-3 α/β (GSK3 α/β) in liver of normal birth weight (NBW) rats, NBW rats supplemented with curcumin (NBW-C), intra-uterine growth restriction (IUGR) rats and IUGR rats supplemented with curcumin (IUGR-C). Data are mean values (n 6 per group), with standard deviations represented by vertical bars. Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate. ^{a,b} Mean values with unlike letters were significantly different ($P < 0.05$). Results were corrected for total protein. B, birth weight; D, diet; $B \times D$, interaction between the corresponding parameters. pGSK3 α ^{Ser21}, phosphorylated glycogen synthase kinase 3 α ^{Ser21}; pGSK3 β ^{Ser9}, phosphorylated glycogen synthase kinase 3 β ^{Ser9}.




Fig. 3. Abundance of proteins in liver of normal birth weight (NBW) rats, NBW rats supplemented with curcumin (NBW-C), intra-uterine growth restriction (IUGR) rats and IUGR rats supplemented with curcumin (IUGR-C). (A) Sterol regulatory element binding protein 1 (SREBP1); (B) fatty acid synthase (FASN); (C) PPAR α . Data are mean values (n 6 per group), with standard deviations represented by vertical bars. Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate. ^{a,b,c} Mean values with unlike letters were significantly different ($P < 0.05$). Results were corrected for total protein. B, birth weight; D, diet; $B \times D$, interaction between the corresponding parameters.




Fig. 4. Effect of curcumin on the hepatic gene expressions of rats with intra-uterine growth restriction (IUGR). Data are mean values (n 6 per group), with standard deviations represented by vertical bars. Data were analysed by using two-way ANOVA and Tukey's *post hoc* testing, where appropriate. ^{a,b,c} Mean values with unlike letters were significantly different ($P < 0.05$). *Srebf1*, sterol regulatory element binding protein 1c; *Fasn*, fatty acid synthase; *HSL*, hormone-sensitive lipase; B, birth weight; D, diet; B \times D, interaction between the corresponding parameters. \square , Normal BW (NBW) rats; \square (hatched), NBW rats supplemented with curcumin (NBW-C); \blacksquare , IUGR rats; \square (white with diagonal lines), IUGR rats supplemented with curcumin (IUGR-C).

In the present study, IUGR increased serum HOMA-IR, insulin and glucose levels, which were considered as markers of IR. The results were consistent with the previous study on insulin sensitivity index, which first formally reported the sequence of IR in short children with IUGR⁽²²⁾. IR is related to a postreceptor defect in the intracellular insulin signalling pathway, leading to the failure of insulin to reduce the level of glucose and improve hepatic glycogen synthesis⁽²³⁾. Insulin regulation effect on glucose metabolism are mediated by IRS-1, which activates PI3K and Akt. The phosphorylation of Akt leads to inactivation of glycogen synthase kinase-3 (GSK3) and thus enhances the glycogen synthesis⁽²⁴⁾. Our results showed that IUGR induced rats increased levels of phosphorylated IRS-1 and Akt. It has been demonstrated that the increased serine phosphorylation of IRS-1 plays an important role in the pathogenesis of IR⁽²⁵⁾. The activation of Akt induced the significant phosphorylation level of GSK3 β in IUGR, not GSK3 α in our study. GSK3 α and GSK3 β are two functional isoforms of GSK3, which originate from different genes, but share 97 % amino acid homology. GSK3 β is the primary kinase in regulating glycogen synthesis in muscle which has a positive effect on glycogen deposition as compared with GSK3 α ^(26,27). The different effects of GSK3 α and β on liver are not well documented. Our research implied that the causes of IR in IUGR might be related to the failure of Akt to suppress GSK3 β . Nachimuthu *et al.*⁽¹⁴⁾ reported that curcumin treatment attenuated the IR by decreasing IRS-1 phosphorylation in rats and alleviated hyperinsulinaemia and HOMA-IR levels. In the present study, the concentrations of curcumin in serum and liver were accumulated in NBW and IUGR rats after treated with curcumin. We found that dietary curcumin supplementation decreased the levels of serum insulin, glucose and HOMA-IR in IUGR rats. Furthermore, results also showed that the levels of IRS-1 serine phosphorylation, Akt and GSK3 β phosphorylation were lower in IUGR rats when supplemented with dietary curcumin. PK acts as a key glycolytic enzyme that catalyses the final step of glycolysis and generates pyruvate in the metabolic process. In our present study, the activity of PK and concentration of pyruvate were decreased and concentration of hepatic glycogen was increased in liver of IUGR rats

supplemented with curcumin. These results implied that curcumin could attenuate IR, inhibit hepatic glycolysis and improve hepatic glycogen deposition through regulating IRS-1/Akt/GSK3 pathway and decreasing glycolytic enzyme activity.

TAG, the most common non-toxic form of fatty acid, is the main lipid stored in the liver of patients with non-alcoholic fatty liver disease⁽²⁸⁾. In the present study, IUGR rats significantly increased the liver concentration of TAG and TC and decreased the serum concentration of TAG. These results indicated that more TAG was exported to liver in IUGR rats. Some previous studies revealed that inhibiting TAG synthesis was helpful for the treatment of hepatic steatosis⁽²⁹⁾. In patients with non-alcoholic fatty liver disease, NEFA provide most of lipid content for hepatic TAG synthesis⁽³⁰⁾. In the present experiment, we observed that hepatic NEFA concentration was increased in IUGR rats. It is well known that accumulation of TAG and increase of NEFA in liver is due to the imbalance between hepatic lipids acquisition and removal⁽³¹⁾. Notably, the activities of enzymes related to lipolysis in liver were decreased in IUGR rats. LPL catalyses the hydrolysis of TAG in circulating chylomicrons, thus supplying NEFA for tissue utilisation⁽³²⁾. HL plays a critical role in hydrolysing TAG and phospholipids in the blood. In our experiment, the decreased activities of LPL and HL in IUGR rats might be the main reasons for inefficient NEFA utilisation and excessive TAG accumulation in liver. More importantly, after dietary curcumin supplementation, the activities of LPL, HL and TL in IUGR rats were increased and TAG accumulation was obviously attenuated to a normal level. Moreover, curcumin significantly decreased the concentrations of hepatic NEFA in IUGR rats. The findings were in agreement with the previous results reported by Asai *et al.*⁽³¹⁾ and Jang *et al.*⁽³³⁾, and reported that curcumin was beneficial to inhibit fatty acid synthases and attenuate hepatic TAG accumulation in high-fat-fed rats. These results suggested that IUGR might have a high risk of hepatic lipid metabolic disorder, and dietary curcumin supplementation could alleviate the accumulation of TAG in liver.

Insulin regulates the lipogenesis through the activation of SREBP-1⁽³⁴⁾. Liver plays a vital role in the regulation of gene expression for lipid metabolism, including SREBP-1c and PPAR α ⁽³⁵⁾. SREBP-1c is one of the isoforms of SREBP and

first enhances the transcription of genes associated with biosynthesis of fatty acids and TAG. FASN is well known to catalyse the last step in the biosynthesis of fatty acids, directly activated by SREBP and involved in TAG synthesis. Overexpression of SREBP-1 c led to the increase in TAG content by more than 4-fold and expression of FASN to be increased by 3.9-fold in the liver⁽³⁶⁾. Emerging evidence has suggested that CD36-mediated fatty acid uptake and increased CD36 expression resulted in the dyslipidemia and hepatic TAG storage⁽³⁷⁾. We found that the transcription levels of *Cd36*, *Srebf1* and *Fasn* were increased in IUGR rats. The protein expressions of SREBP1 and FASN in the liver of IUGR rats also were increased. A previous study indicated that IUGR could easily cause hepatic TAG accumulation by increasing the levels of SREBP and FASN proteins during early life⁽¹⁰⁾, which were similar to our results. PPAR α promotes β -oxidation of fatty acids which is mainly expressed in the liver. HSL is believed to be a key enzyme for improving the decomposition of TAG⁽³⁸⁾ and inhibited by insulin⁽³⁹⁾. Our results showed that the transcription levels of *HSL* and the protein and mRNA expressions of PPAR α were decreased in the liver of IUGR rats. The present study suggested that IUGR inhibited the hepatic β -oxidation of fatty acids by regulating *HSL* and PPAR α . In mice administrated with a methionine choline-deficient diet, PPAR α -deficiency provoked more severe steatosis, whereas PPAR α activation enhanced hepatic lipid turnover⁽⁴⁰⁾. Furthermore, activation of PPAR α obviously reduced adiposity in mice fed with a high-fat diet⁽⁴¹⁾. These results implied that IUGR could improve lipid accumulation and inhibit lipolysis in the liver which might be related to the alterations of the expression of SREBP1 and target genes. Interestingly, curcumin had a potential ability of attenuating hepatic lipid deposition in IUGR rats. In the present study, dietary curcumin supplementation significantly decreased the protein expressions of SREBP1, FASN and increased the protein expression of PPAR α in the liver of IUGR rats. Curcumin also down-regulated the transcription levels of *Cd36*, *Srebf1* and *Fasn*, and up-regulated the transcription levels of *Ppara* and *HSL* in IUGR rats. Previous studies indicated that curcumin could suppress the gene expression of *Fasn*⁽⁴²⁾ and activate protein expression of PPAR α ⁽⁴³⁾. According to the results of the present study, we could confirm that curcumin could promote lipolysis and fatty acid oxidation in the liver of IUGR rats.

In conclusion, our present data suggested that IUGR rats exhibited a high risk of IR and hepatic lipid accumulation in the liver. Curcumin efficiently attenuated IR of IUGR rats, which might contribute to reduce insulin level in serum and regulate the insulin signalling pathways in liver. Curcumin also alleviated hepatic lipid accumulation of IUGR rats by regulating SREBP target genes and lipid metabolism-related genes in liver, which might contribute to the inhibition of lipogenesis and promotion of lipolysis. Our findings may be helpful in finding a new nutritional strategy for the prevention or treatment of IUGR in humans in future.

Acknowledgements

We are thankful to Kehu Bio-technology Research Center (Guangzhou, People's Republic of China) for kindly supplying the curcumin.

This work was supported by the National Natural Science Foundation of China (nos. 31572418, 31802101, 31472129 and 31601948); the National key research and development programme of China (2018YFD0501101); and the Fundamental Research Funds for the Central Universities (KJQN201935).

The authors' responsibilities were as follows: Y. N. designed the research, conducted the research, analysed the data, and wrote and revised the manuscript; H. A. revised the manuscript; J. T. H. and C. W. designed the research; T. C., L. L. Z., X. Z. and J. F. Z. conducted the research; T. W. had primary responsibility for final content. All authors read and approved the final manuscript.

None of the authors declared a conflict of interest.

References

- Garite TJ, Clark R & Thorp JA (2004) Intrauterine growth restriction increases morbidity and mortality among premature neonates. *Am J Obstet Gynecol* **191**, 481–487.
- Wu G, Bazer FW, Wallace JM, *et al.* (2006) Board-invited review: intrauterine growth retardation: implications for the animal sciences. *J Anim Sci* **84**, 2316–2337.
- De Onis M, Blossner M & Villar J (1998) Levels and patterns of intrauterine growth retardation in developing countries. *Eur J Clin Nutr* **52**, S5–S15.
- Barker DJ, Eriksson JG, Forsen T, *et al.* (2002) Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol* **31**, 1235–1239.
- Alisi A, Panera N, Agostoni C, *et al.* (2011) Intrauterine growth retardation and nonalcoholic fatty liver disease in children. *Int J Endocrinol* **2011**, 269853.
- Tarry-Adkins JL & Ozanne SE (2011) Mechanisms of early life programming: current knowledge and future directions. *Am J Clin Nutr* **94**, 1765S–1771S.
- Liu C, Lin G, Wang X, *et al.* (2013) Intrauterine growth restriction alters the hepatic proteome in fetal pigs. *J Nutr Biochem* **24**, 954–959.
- Tarryadkins JL, Fernandeztwinn DS, Hargreaves IP, *et al.* (2015) Coenzyme Q10 prevents hepatic fibrosis, inflammation, and oxidative stress in a male rat model of poor maternal nutrition and accelerated postnatal growth. *Am J Clin Nutr* **103**, 579–588.
- He J, Dong L, Xu W, *et al.* (2015) Dietary tributyrin supplementation attenuates insulin resistance and abnormal lipid metabolism in suckling piglets with intrauterine growth retardation. *PLOS ONE* **10**, e0136848.
- Magee TR, Han G, Cherian B, *et al.* (2008) Down-regulation of transcription factor peroxisome proliferator-activated receptor in programmed hepatic lipid dysregulation and inflammation in intrauterine growth-restricted offspring. *Am J Obstet Gynecol* **199**, e1–e5.
- Aggarwal BB, Sundaram C, Malani N, *et al.* (2007) Curcumin: the Indian solid gold. *Adv Exp Med Biol* **595**, 1–75.
- Miłobędzka J, Kostanecki SV & Lampe V (1910) Zur Kenntnis des Curcumins (Knowledge of curcumin). *Eur J Inorg Chem* **43**, 2163–2170.
- Jang EM, Choi MS, Jung UJ, *et al.* (2008) Beneficial effects of curcumin on hyperlipidemia and insulin resistance in high-fat-fed hamsters. *Metab Clin Exp* **57**, 1576–1583.
- Maithilikarpagaselvi N, Sridhar MG, Swaminathan RP, *et al.* (2016) Curcumin prevents inflammatory response, oxidative stress and insulin resistance in high fructose fed male Wistar rats: potential role of serine kinases. *Chem Biol Interact* **244**, 187–194.

15. Desai M, Gayle D, Babu J, *et al.* (2005) Programmed obesity in intrauterine growth-restricted newborns: modulation by newborn nutrition. *Am J Physiol Regul Integr Comp Physiol* **288**, R91–R96.
16. Bucolo G & David H (1973) Quantitative determination of serum triglycerides by the use of enzymes. *Clin Chem* **19**, 476–482.
17. Yokode M, Hammer RE, Ishibashi S, *et al.* (1990) Diet-induced hypercholesterolemia in mice: prevention by overexpression of LDL receptors. *Science* **250**, 1273–1275.
18. Kunati SR, Yang S, William BM, *et al.* (2018) An LC-MS/MS method for simultaneous determination of curcumin, curcumin glucuronide and curcumin sulfate in a phase II clinical trial. *J Pharm Biomed Anal* **156**, 189–198.
19. Schmittgen TD & Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* **3**, 1101–1108.
20. Lim K, Armitage JA, Stefanidis A, *et al.* (2011) IUGR in the absence of postnatal 'catch-up' growth leads to improved whole body insulin sensitivity in rat offspring. *Pediatr Res* **70**, 339–344.
21. He J, Niu Y, Wang F, *et al.* (2018) Dietary curcumin supplementation attenuates inflammation, hepatic injury and oxidative damage in a rat model of intra-uterine growth retardation. *Br J Nutr* **120**, 537–548.
22. Hofman PL, Cutfield WS, Robinson EM, *et al.* (1997) Insulin resistance in short children with intrauterine growth retardation. *J Clin Endocrinol Metab* **82**, 402–406.
23. Barthel A & Schmolz D (2003) Novel concepts in insulin regulation of hepatic gluconeogenesis. *Am J Physiol Endocrinol Metab* **285**, E685–E692.
24. Rayasam GV, Tulasi VK, Sodhi R, *et al.* (2009) Glycogen synthase kinase 3: more than a namesake. *Br J Pharmacol* **156**, 885–898.
25. Taniguchi CM, Emanuelli B & Kahn CR (2006) Critical nodes in signalling pathways: insights into insulin action. *Nat Rev Mol Cell Biol* **7**, 85–96.
26. Macaulay K, Blair AS, Hajduch E, *et al.* (2005) Constitutive activation of GSK3 down-regulates glycogen synthase abundance and glycogen deposition in rat skeletal muscle cells. *J Biol Chem* **280**, 9509–9518.
27. Pearce NJ, Arch JR, Clapham JC, *et al.* (2004) Development of glucose intolerance in male transgenic mice overexpressing human glycogen synthase kinase-3 β on a muscle-specific promoter. *Metab Clin Exp* **53**, 1322–1330.
28. Tilg H & Moschen AR (2008) Insulin resistance, inflammation, and non-alcoholic fatty liver disease. *Trends Endocrinol Metab* **19**, 371–379.
29. Yamaguchi K, Yang L, McCall S, *et al.* (2007) Inhibiting triglyceride synthesis improves hepatic steatosis but exacerbates liver damage and fibrosis in obese mice with nonalcoholic steatohepatitis. *Hepatology* **45**, 1366–1374.
30. Musso G, Gambino R & Cassader M (2009) Recent insights into hepatic lipid metabolism in non-alcoholic fatty liver disease (NAFLD). *Prog Lipid Res* **48**, 1–26.
31. Asai A & Miyazawa T (2001) Dietary curcuminoids prevent high-fat diet-induced lipid accumulation in rat liver and epididymal adipose tissue. *J Nutr* **131**, 2932–2935.
32. Mead JR, Irvine SA & Ramji DP (2002) Lipoprotein lipase: structure, function, regulation, and role in disease. *J Mol Med* **80**, 753–769.
33. Jang EM, Choi MS, Jung UJ, *et al.* (2008) Beneficial effects of curcumin on hyperlipidemia and insulin resistance in high-fat-fed hamsters. *Metab Clin Exp* **57**, 1576–1583.
34. Tessari P, Coracina A, Cosma A, *et al.* (2009) Hepatic lipid metabolism and non-alcoholic fatty liver disease. *Nutr Metab Cardiovasc Dis* **19**, 291–302.
35. Jump DB, Botolin D, Wang Y, *et al.* (2005) Fatty acid regulation of hepatic gene transcription. *J Nutr* **135**, 2503–2506.
36. Shimano H, Horton JD, Shimomura I, *et al.* (1997) Isoform 1 c of sterol regulatory element binding protein is less active than isoform 1a in livers of transgenic mice and in cultured cells. *J Clin Invest* **99**, 846–854.
37. Koonen DPY, Jacobs RL, Febbraio M, *et al.* (2007) Increased hepatic cd36 expression contributes to dyslipidemia associated with diet-induced obesity. *Diabetes* **56**, 2863–2871.
38. Langin D, Laurell H, Holst LS, *et al.* (1993) Gene organization and primary structure of human hormone-sensitive lipase: possible significance of a sequence homology with a lipase of Moraxella TA144, an antarctic bacterium. *Proc Natl Acad Sci U S A* **90**, 4897–4901.
39. Haemmerle G, Zimmermann R, Hayn M, *et al.* (2002) Hormone-sensitive lipase deficiency in mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *J Biol Chem* **277**, 4806–4815.
40. Ip E, Farrell GC, Robertson G, *et al.* (2003) Central role of PPAR α -dependent hepatic lipid turnover in dietary steatohepatitis in mice. *Hepatology* **38**, 123–132.
41. Goto T, Lee JY, Teraminami A, *et al.* (2011) Activation of peroxisome proliferator-activated receptor- α stimulates both differentiation and fatty acid oxidation in adipocytes. *J Lipid Res* **52**, 873–884.
42. Younesian O, Kazerouni F, Dehghan-Nayeri N, *et al.* (2017) Effect of curcumin on fatty acid synthase expression and enzyme activity in breast cancer cell line SKBR3. *Int J Cancer Manag* **10**, e8173.
43. Kang O, Kim S, Seo Y, *et al.* (2013) Curcumin decreases oleic acid-induced lipid accumulation via AMPK phosphorylation in hepatocarcinoma cells. *Eur Rev Med Pharmacol Sci* **17**, 2578–2586.