

Zinc-clotrimazole complexes are effective against *Trichomonas vaginalis*

Victor Midlej¹, Felipe Rubim¹, Wilmer Villarreal², Érica S. Martins-Duarte³, Maribel Navarro^{4,5}, Wanderley de Souza¹ and Marlene Benchimol^{1,6}

Research Article

Cite this article: Midlej V, Rubim F, Villarreal W, Martins-Duarte ÉS., Navarro M, de Souza W, Benchimol M (2019). Zinc-clotrimazole complexes are effective against *Trichomonas vaginalis*. *Parasitology* **146**, 1206–1216. <https://doi.org/10.1017/S003118201900043X>

Received: 11 December 2018

Revised: 19 March 2019

Accepted: 20 March 2019

First published online: 3 May 2019

Key words:

Chemotherapy; clotrimazole; electron microscopy; hydrogenosomes; trichomoniasis; zinc complexes; ultrastructure

Author for correspondence:

Marlene Benchimol, E-mail: marlenebenchimol@gmail.com

¹Laboratório de Ultraestrutura Celular Hertha Meyer, Instituto de Biofísica Carlos Chagas Filho, Instituto Nacional de Ciência e Tecnologia, and Núcleo de Biologia Estrutural e Bioimagens, Universidade Federal do Rio de Janeiro – UFRJ, Rio de Janeiro, Brasil; ²Departamento de Química, Universidade Federal de São Carlos-UFSCar, São Carlos, SP, Brasil; ³Departamento de Parasitologia, Universidade Federal de Minas Gerais, Belo Horizonte, Brasil; ⁴Instituto Nacional de Metrologia, Qualidade e Tecnologia-Inmetro, Duque de Caxias, Rio de Janeiro, Brasil; ⁵Departamento de Química, ICE, Universidade Federal de Juiz de Fora, Juiz de Fora-MG, Brasil and ⁶Universidade do Grande Rio-Unigranrio, Duque de Caxias, Rio de Janeiro, Brasil

Abstract

Trichomonas vaginalis is a protozoan parasite that causes trichomoniasis in humans, the most prevalent non-viral sexually transmitted disease (STD). Imidazole compounds are used for the treatment of trichomoniasis, and metronidazole is the most commonly prescribed. However, these compounds can lead to parasite resistance and unwanted side effects. Therefore, there is a need for an alternative treatment for this disease. Here, we explored the potential of clotrimazole (CTZ) and zinc compounds, as well as CTZ complexed with zinc salts ([1] acetate $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ and [2] a chloride $[\text{Zn}(\text{CTZ})_2\text{Cl}_2]$ complexes) against *T. vaginalis*. We synthesized the zinc complexed CTZ compounds and determined their concentration values that inhibited parasite growth by 50% (IC₅₀). We used scanning and transmission electron microscopy to visualize the ultrastructural alterations induced by CTZ and their zinc complexes. The incubation of the parasites with $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ complex inhibited their growth, yielding an IC₅₀ of 4.9 μM . Moreover, there were changes in the shape of treated parasites, including the formation of surface projections that subsequently detached from the cell, in addition to changes in the hydrogenosomes, endoplasmic reticulum and Golgi complex. We found $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ to be a highly effective compound against *T. vaginalis* *in vitro*, suggesting its potential utility as an alternative chemotherapy for trichomoniasis.

Introduction

Human trichomoniasis is one of the most common non-viral sexually transmitted diseases. It affects about 278 million people worldwide (WHO, 2012) and is caused by the parasite *Trichomonas vaginalis*. Trichomoniasis induces an infectious process in the urogenital tract, and in pregnant women, the disease can lead to severe complications: increased risk of placental membrane rupture, premature birth and chronic pulmonary disease acquired during childbirth (Hirt, 2013). In men, the disease is often asymptomatic, but it can cause urethritis, prostatitis and infertility (Secor, 2012). *T. vaginalis* can cause infertility in both men and women (Lewis, 2010; Mielczarek and Blaszkowska, 2016). Trichomoniasis has received a great deal of attention due to the severe consequences of the disease, such as increased risks for HIV infection and cervical and prostate cancers (Secor, 2012; Hirt, 2013).

The 5-Nitroimidazole compounds, such as metronidazole [1-(2-hydroxyethyl)-2-methyl-5-nitroimidazole] (Edwards, 1993), are recommended as trichomoniasis treatments. Although metronidazole has high efficacy, it has several negative side effects (Secor, 2012). These include nausea, vomiting, metallic taste in the mouth, diarrhea and peripheral neuropathies (Rosa *et al.*, 2011). Moreover, several studies documented metronidazole resistance (Houang *et al.*, 1997; Schwebke and Barrientes, 2006). Therefore, the development of a safer compound for the treatment of this infection is needed.

Clotrimazole (CTZ) has been used for *Candida albicans* infections and other diseases, such as sickle cell disease and some cancers (Crowley and Gallagher, 2014). In addition, clotrimazole was previously used for the treatment of metronidazole-resistant trichomoniasis (Cudmore *et al.*, 2004). When clotrimazole is combined with other molecules, such as metals, it displays promising pharmacological characteristics (Navarro *et al.*, 2006, 2009). Assuming that the zinc present in the prostatic fluid has microbicidal effects on *T. vaginalis* (Levy and Fair, 1973; Benchimol *et al.*, 1993), we investigated the activity of zinc-clotrimazole complexes against the parasite. Here, we demonstrate a significant anti-proliferative effect of the zinc-clotrimazole complexes *in vitro* against *T. vaginalis*.

Materials and methods

Parasite and cell culture

For the analyses of drug activity against *T. vaginalis*, we used the JT strain isolated in the Hospital of Federal University of Rio de Janeiro in Brazil. Trophozoites were maintained in

Diamond's trypticase yeast-extract maltose (TYM) medium supplemented with 10% foetal bovine serum (FBS), and the cells were grown for 24 h at 37 °C. A human foreskin fibroblast (HFF; ATCC) cell line was chosen to evaluate the drug's cytotoxicity. HFFs were grown in 25-cm² culture flasks containing Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 2 mM L-glutamine and 10% FBS at 37 °C under a 5% CO₂ atmosphere.

Chemicals and instrumentation

All commercial reagents (CTZ, ZnCl₂ and ZnAc₂·2H₂O) were used without further purification. Analytical-grade solvents were distilled from appropriate drying agents prior to use. The Infrared (IR) spectra were obtained using a Shimadzu spectrometer in the 4000–400 cm⁻¹ region. The elemental composition of the metal complexes was determined using a Perkin Elmer 2400 CHNS/O Series II microanalyzer. Conductivity values were obtained using a Meter Lab CDM2300 instrument. Nuclear Magnetic Resonance (NMR) experiments were performed at 298 K on a Bruker DRX 400 MHz spectrometer at 9.4 T, and ¹H was observed at 400.13 MHz, while ¹³C{¹H} was observed at 100.62 MHz. The NMR spectra were recorded in dimethyl sulfoxide (DMSO)-d₆ solutions with tetramethylsilane TMS.

Synthesis of zinc-clotrimazole complexes

[Zn(CTZ)₂(Ac)₂].2H₂O

To prepare [Zn(CTZ)₂(Ac)₂].2H₂O, 710 mg (2.06 mM) of CTZ was dissolved in methanol (30 mL), and 205 mg (0.93 mM) of ZnAc₂·2H₂O was added to the solution, which was then stirred at room temperature for 48 h. The volume of the solvent was reduced, and distilled water was added. The white solid obtained was filtered, washed with cold methanol and dried under vacuum (yield: 778 mg, 94%). Elemental analysis: Calculated for C₄₈H₄₀N₄Cl₂O₄Zn·2H₂O: C, 63.41; H, 4.88; N, 6.16. Found: C, 63.41; H, 5.30; N, 6.05. IR (cm⁻¹): ν(O-H) 3440, ν(C=N) 1585, ν(C=C) 1492, ν_a(COO) 1618 and ν_s(COO) 1433. ¹H NMR [DMSO; δ ppm (multiplicity, integration, assignation): 7.77 (s, 1H, H2); 7.51 (m, 2H, H9 and H8); 7.41 (m, 7H, H7, H10, H12, H14, H15, H17, H19); 7.11 (s, 1H, H5); 7.06 (m, 4H, H11, H13, H16, H18); 6.96 (s, 1H, H4); 6.91 (d, 1H, H6, *J*_{ortho} = 7.84 Hz), 1.75 (s, 3H, Ac). Molar Conductivity in DMSO, Λ_M = 0.35 ± 0.03 Ω⁻¹ cm² M⁻¹.

[Zn(CTZ)₂Cl₂]

To prepare [Zn(CTZ)₂Cl₂], 308 mg (0.89 mM) of CTZ was dissolved in 20 mL of methanol, and a solution of 58 mg (0.43 mM) of ZnCl₂ in 10 mL of methanol was added. The mixture was stirred at room temperature for 24 h. The white solid obtained was filtered, washed with distilled water and dried under a vacuum (yield, 291 mg, 82%). Elemental analysis: Calculated for C₄₄H₃₄N₄Cl₂Zn: C, 63.98; H, 4.15; N, 6.78. Found: C, 64.52; H, 3.53; N, 6.51. IR (cm⁻¹): ν(C=N) 1589, ν(C=C) 1492. ¹H NMR [DMSO; δ ppm (multiplicity, integration, assignation): 7.67 (s, 1H, H2); 7.52 (m, 2H, H9 and H8); 7.42 (m, 7H, H7, H10, H12, H14, H15, H17, H19); 7.11 (s, 1H, H5); 7.07 (m, 4H, H11, H13, H16, H18); 7.03 (s, 1H, H4); 6.93 (d, 1H, H6, *J*_{ortho} = 7.71 Hz). Molar Conductivity in DMSO, Λ_M = 3.57 ± 0.05 Ω⁻¹ cm² M⁻¹. The same compound was independently synthesized using a slightly different procedure (Betanzos-Lara *et al.*, 2012).

In vitro anti-proliferative activities

Parasites were grown to a density of 2 × 10⁴ cells mL⁻¹. Next, CTZ and the zinc-clotrimazole complexes, [Zn(CTZ)₂(Ac)₂] and [Zn

(CTZ)₂Cl₂], were added at different concentrations (1, 5, 10 or 20 μM) from stock solutions diluted in DMSO. The final concentration of DMSO in the growth medium never exceeded 1% (v/v), and it had no effect on cell growth or cell morphology. Cell densities were determined using a haemocytometer.

Combination treatment

We performed a combination treatment in order to evaluate the combined effect of CTZ with ZnAc₂. The drugs were added as separated compounds following the ratio of 2:1 (CTZ: ZnAc₂). In [Zn(CTZ)₂(Ac)₂], for each zinc atom there are two molecules of clotrimazole molecule in the complex. Based in the concentration of zinc-clotrimazole complexes we set the combination treatment equalizing both compounds with the same number of available molecules. For construction of the association growth curve, we treated the parasites simultaneously with 1:2, 5:10, 10:20 and 20:40 μM of ZnAc₂: Clotrimazole, respectively. Cell densities were determined using a haemocytometer.

Determination of IC50

To determine the drug concentration that inhibited parasite growth by 50% (IC₅₀), the percentage of growth inhibition was plotted as a function of the drug concentration, and the values were fitted in a non-linear curve. Regression analyses were performed using the Sigma Plot 8.0 software. Data were plotted using the Graphpad Prism 6.0[®] software.

Immunofluorescence

Control or treated parasites were washed in warm PBS (pH 7.2) and allowed to adhere to glass coverslips with poly-L-lysine for 10 min at 37 °C. Cells were fixed for 1 h with 4% formaldehyde, permeabilized with 3% Nonidet (NP-40) and incubated with 50 mM NH₄Cl and 3% of bovine serum albumin in PBS (PBS/BSA) as blocking solutions, after each step mentioned above the samples were washed with PBS (pH 8.0). Trichomonas were labeled with polyclonal antibody anti-pyruvate:ferrodoxin oxidoreductase (PFOR) (Provided by Dr Jan Tachezy, Charles University, Czech Republic). Afterwards, the coverslips were washed with PBS (pH 8.0) and incubated for 1 h with an Alexa Fluor 488-conjugated anti-mouse IgG antibody (Life Technologies) that was diluted 1:100 in PBS/BSA. Finally, the coverslips were washed and examined using an Axio Observer fluorescence microscope (Zeiss, Germany).

Scanning electron microscopy (S.E.M.)

Parasites were adhered as described above and fixed in 2.5% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.2). They were then post-fixed for 15 min in 1% OsO₄. The parasites were dehydrated in an ascending series of ethanol into 100% ethanol, critical-point dried with liquid CO₂, coated with a 15-nm thick layer of gold and observed using a Quanta S.E.M. (FEI company, The Netherlands). For high-resolution observations, the sputter-coating step was performed using a thin layer (2 nm) of platinum, and cells were observed on an Auriga High-Resolution S.E.M. (Zeiss, Germany).

Transmission electron microscopy (TEM)

Control and treated trichomonas were fixed as described previously for S.E.M. After fixation, cells were post-fixed for 40 min in a solution containing 1% OsO₄ and 0.8% potassium ferrocyanide in 0.1 M cacodylate buffer. They were then washed in PBS,

dehydrated in acetone and embedded in Epon (an epoxy resin). Ultrathin sections (60–70-nm thick) were stained with uranyl acetate and lead citrate and were observed using a Tecnai Spirit TEM (FEI company, the Netherlands).

Viability analysis

To measure parasite viability, non-treated and IC50 concentration-treated *T. vaginalis* were stained with 6 μM of propidium iodide (PI) (Sigma, USA) for 30 min. The analyses were performed after 48 h of incubation with the drug. Cell fluorescence was analyzed using an Accuri flow cytometer (BD Biosciences, USA). Overall, three independent experiments were performed in duplicate. The cytotoxic effects of the compounds on HFF cells and the viability of *T. vaginalis* when treated with all compounds were evaluated based on the reduction of MTS [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfo-phenyl) 2H-tetrazolium] into the soluble formazan dye (Promega, USA). For this assay, 2×10^5 HFF cells were seeded into 96-well plates and cultured with DMEM supplemented with 2 mM L-glutamine and 10% FBS at a final volume of 200 μL per well. After the cells reached confluence, they were incubated for 48 h with clotrimazole and complexes (1) and (2) at different concentrations in DMEM supplemented with 2 mM L-glutamine and 10% FBS. The dilution concentrations ranged from 10–50 μM for clotrimazole and 5–30 μM for complexes 1 and 2. At the end of the experiment, the HFF cells were washed with PBS and incubated with 100 μL of 10 mM glucose in PBS plus 20 μL of MTS (Promega, USA). The same methodology was applied for *T. vaginalis*, where the initial amount of cells was 2×10^4 parasites per well. The absorbance was determined at 490 nm after 2 h of incubation in the microplate reader SpectraMax M2e (Molecular Devices). Cytotoxicity was calculated as a percentage of viable cells vs untreated cells (control, 100%). Control cells were treated with 4% formaldehyde.

Results

Synthesis and characterization

Figure 1 shows the synthesis scheme for $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ [1] and $[\text{Zn}(\text{CTZ})_2\text{Cl}_2]$ [2], which were obtained at a high yield. All of these metal complexes were obtained as air-stable white solids. The elemental analyses determined were in agreement with the molecular formula proposed for each Zn-CTZ complex. In order to analyze if these metal complexes were neutral or cationic, their molar conductivity were measured in DMSO, finding that they are in the range for non-electrolyte complexes (17). The IR spectra of complexes 1 and 2 showed frequencies clearly associated with the characteristic functional group of the clotrimazole ligand, as well as the characteristic bands for each metal salts used in this synthesis. In the case of complex 1, signals that corresponded to the stretching vibrations could be observed due to the acetate ligand-binding unidentate form of the metal centre [which resulted from the differences between the values of ν_a and ν_s COO^- (185 cm^{-1}) that were greater than the free ligand (164 cm^{-1})]. Based on the ^1H -NMR spectra results, it is possible to propose that CTZ bound to zinc by the unsubstituted N1 atom of the imidazole, due to the ^1H chemical shift variation observed for the Zn-CTZ complexes with respect to those of the free CTZ ($\Delta\delta$). A significant downfield shifts ($\Delta\delta$) was observed for the imidazole protons attached to the zinc, as an example, H_2 was 0.36 ppm for complex 1 and 0.25 ppm for complex 2 with respect to free CTZ (Fig. 2). It was also possible observed for complex 1, than the relative integral of the signals for the methyl protons of the acetate group in the spectrum was in agreement with an acetate group:CTZ ratio of 1:1.




Fig. 1. Scheme of synthesis of Zn-CTZ complexes.




Fig. 2. ^1H NMR spectra of complexes $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$, $[\text{Zn}(\text{CTZ})_2\text{Cl}_2]$ and CTZ free in DMSO-d_6 .

Anti-proliferative effects

When we added CTZ and its zinc complexes in different concentrations (1, 5, 10 and 20 μM) to *T. vaginalis* cultures after 24 h of initial parasite growth, they exhibited a dose-dependent reduction in cell growth (Fig. 3, Supplementary Fig. 1). The IC50 (Supplementary Fig. 2) and IC90 values obtained after a 48 h incubation period for all compounds were as follows: CTZ, 17.2 μM and $>40\text{ }\mu\text{M}$ (Fig. 3a); $[\text{Zn}(\text{CTZ})_2\text{Cl}_2]$, 10.5 and 22.9 μM (Fig. 3c) and $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$, 4.9 and 17.2 μM (Fig. 3d). In addition, it is important to highlight that at the final concentrations used, DMSO affected neither parasite growth (Fig. 3) nor morphology when observed by light, s.e.m. and TEM (data not shown). These results indicated that the $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ compound was more effective against the parasite, it reduced IC50 values by around 70% (Fig. 3, Supplementary Fig. 1). To confirm the superior effect of the $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ complex, we evaluated the combination of CTZ plus the salt ZnAc_2 against *T. vaginalis* (Fig. 3e, Supplementary Fig. 1). For this experiment we used a ratio of 2:1 CTZ: ZnAc_2 , once in the Zn-CTZ complexes formula the CTZ is twice per mole. The concentrations used in this experiment were 1:2, 5:10, 10:20 and 20:40 μM of ZnAc_2 : Clotrimazole, respectively. The IC50 value for CTZ added in combination with ZnAc_2 after 48 h of treatment was 15.7 μM (Fig. 3e, Supplementary Fig. 2), which very close to the IC50 value of CTZ alone (17.2 μM). Thus, this result confirms that the $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ compound




Fig. 3. Antiproliferative effect of clotrimazole and zinc-clotrimazole complexes against *T. vaginalis*. The growth curve of parasites treated with different concentrations of (a) non-complexed clotrimazole, CTZ; (b) zinc acetate salt, ZnAc₂; and zinc-clotrimazole complexes (c) [Zn(CTZ)₂Cl₂] and (d) [Zn(CTZ)₂(Ac)₂]; (e) zinc acetate salt with non-complexed clotrimazole combination, ZnAc₂ + CTZ. Combined treatment (e) were performed using the ratio of 2:1 (CTZ:ZnAc₂). The compounds were added after 24 h (arrow) of cell growth. The number of cells was determined by light microscopy using a haemocytometer. (f) Percentage of parasite growth treated with different concentrations of non-complexed CTZ, ZnAc₂, Zn-CTZ complexes: [Zn(CTZ)₂Cl₂] and [Zn(CTZ)₂(Ac)₂], and ZnAc₂ + CTZ combination after 48 h of treatment. Note that in 48 h of treatment (f) there is a significant alteration comparing control with treated parasites, CTZ and ZnAc₂ + CTZ combination with [Zn(CTZ)₂(Ac)₂] compound. Statistical significance of differences among the groups was assessed using the two-way analysis of variance (2way-ANOVA) test followed by Turkey's multiple comparison test in the GraphPad Prism 6 Software. Results were considered statistically significant when $P \leq 0.03$. (**** $P < 0.0001$; *** $P < 0.001$). The data represent the mean \pm Standard Deviation (s.d.) of three independent experiments performed in triplicate for each compound.

(Fig. 3, Supplementary Fig. 1) is more effective against *T. vaginalis* than the CTZ, even when its used twice concentration.

A comparison of parasite growth in percentage was performed (Fig. 3f, Supplementary Fig. 1e). We assumed in each time point the control as 100% and the percentage value of all compounds used was relative to its control. This data corroborates the growth curves (Fig. 3a-e); the [Zn(CTZ)₂(Ac)₂] compound presented the most significant effect when compared with CTZ treated parasites

(P value < 0.0001) and the ZnAc₂ + CTZ combined treatment (P value < 0.001) (Fig. 3f).

Ultrastructural effects

Untreated *T. vaginalis* (Fig. 4a-b) displayed a normal morphology when visualized both by S.E.M. (Fig. 4a) and TEM (Fig. 4b). By S.E.M., control cells showed a pear-shaped body, a protruding




Fig. 4. Ultrastructural features of *T. vaginalis* by electron microscopy. (a) s.e.m. and (b) TEM of untreated parasites are seen. The cells exhibited regular morphology, including a pear shape, four anterior flagella (AF), a recurrent flagellum (RF) and an axostyle (Ax). (c–h) Morphological alterations in drug-treated parasites visualized by s.e.m. *T. vaginalis* were incubated with 19 μM of clotrimazole (c–d), 10.3 μM of [Zn(CTZ)₂Cl₂] (e–f) and 5.1 μM of [Zn(CTZ)₂(Ac)₂] (g–h). All compounds altered the morphology of the parasites, including cell rounding (arrowhead) induced by both non-complexed (c–d) and zinc-complexed clotrimazole agents, [Zn(CTZ)₂Cl₂] (e–f) and [Zn(CTZ)₂(Ac)₂] (g). In rounded cells, the flagella are externalized (c–e, g). Membrane projections (filopodia) are visible (arrows) (f, h), which are enhanced with [Zn(CTZ)₂(Ac)₂] treatment (g–h). AF, anterior flagella; Ax, axostyle; C, costa; N, anteriorly positioned nucleus; H, hydrogenosomes with peripheral vesicle (arrows); RF, recurrent flagellum; UM, undulating membrane; V, vacuoles.




Fig. 5. Ultrastructural alterations in treated *T. vaginalis*. The cells treated with non-complexed (**a–b**) and zinc-complexed (**c–f**) clotrimazole are observed by TEM. Parasites treated with $19\ \mu\text{M}$ of clotrimazole (**a–b**) displays fewer membrane projections (arrows) when compared with the cells treated with $101\ \mu\text{M}$ of $[\text{Zn}(\text{CTZ})_2\text{Cl}_2]$ (**c–d**) and $5.1\ \mu\text{M}$ of $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ (**e–f**), which exhibited a greater number of filopodia (arrows). An irregular distribution of Golgi lamellas (**a**, **c**, **e**) and abnormal hydrogenosomes (arrowhead) are observed (**a–b**, **d**). The ER is atypical and enlarged following $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ treatment (**e**), and involves cytoplasmic structures (**e**). Ax, Axostyle; ER, endoplasmic reticulum; G, Golgi complex; H, hydrogenosome; N, nucleus; V, vacuoles.




Fig. 6. Quantitative analysis of the effects of clotrimazole on parasite ultrastructure. One hundred parasites were counted using TEM. The values are expressed in percentages. Hydrogenosomes and the plasma membrane were primarily affected in both non-complexed and zinc-complexed clotrimazole. The [Zn(CTZ)₂(Ac)₂] compound induces atypical hydrogenosomes and membrane projections in approximately 56% and 52% of the treated parasites, respectively.




Fig. 7. Effects of [Zn(CTZ)₂(Ac)₂] treatment on *T. vaginalis*. Trophozoites were treated with 5.1 μ M of [Zn(CTZ)₂(Ac)₂] and observed by High resolution-scanning electron microscopy (HR-s.E.M.) (a–c) and TEM (d–f). Protrusions of plasma membrane are identified (arrows) by HR-s.E.M. (a–c). Membrane projections (arrowhead) welled up from the plasma membrane (b–c). Atypical Golgi complex (G) (d) and hydrogenosomes (H) are observed by TEM (e). Cytoplasmic components are seen in the membrane projections (arrows) (f). ER, endoplasmic reticulum; N, nucleus; V, vacuoles.

axostyle, an adhered recurrent flagellum and four anterior flagella emerging from the anterior region of the parasite's cell body (Fig. 4a). Moreover, using TEM, the nucleus of non-treated trichomonas was positioned at the anterior region, and hydrogenosomes, the membranous profiles of the endoplasmic reticulum, Golgi complex and vacuoles were observable (Fig. 4b). However, parasites incubated in the presence of the compounds displayed several ultrastructural alterations (Figs 4–7): rounded cells, the presence of surface projections, atypical hydrogenosomes and endoplasmic reticulum (Fig. 6).

The extent of ultrastructural alterations found in 48 h-treated parasites and were different when CTZ, [Zn(CTZ)₂Cl₂] and [Zn(CTZ)₂(Ac)₂] were compared (Fig. 6). We observed many altered cells when zinc-CTZ complexes were used compared to non-complexed CTZ (Fig. 6). Furthermore, [Zn(CTZ)₂(Ac)₂] was the most effective compound, and quantitative analyses revealed that more than 50% of treated parasites exhibited non-organized

membrane projections (Figs 4–7) and atypical hydrogenosomes (Figs 4–8). Although rounded cells were frequently observed following the use of all compounds (Figs 4–6), endoflagellar forms were not identified (Fig. 4a), and cells displayed externalized flagella (Fig. 4a–c and e). Membrane projections were only seen in treated cells (Figs 4, 5 and 7) as highlighted in Fig. 6. They corresponded to surface projections on the protozoan's plasma membrane filled with cytoplasmic content (Figs 5c, 5f and 7f). When *T. vaginalis* was treated with [Zn(CTZ)₂Cl₂] and [Zn(CTZ)₂(Ac)₂], these surface extensions detached from the cell body (Figs 5f and 7f). Using high-resolution s.E.M., we observed some membrane projections that appeared as stretched structures welling up from the plasma membrane (Fig. 7a–c). Besides the membrane projections, other important alterations in [Zn(CTZ)₂(Ac)₂]-treated parasites were observed (Fig. 7d–e), including disorganized Golgi lamellae (Figs 5e and 7d) and altered hydrogenosomes (Fig. 7e).




Fig. 8. *T. vaginalis* hydrogenosomes after treatment with clotrimazole compounds under fluorescence microscopy. (**a–c**): Non-treated parasites. (**d–f**): *T. vaginalis* treated with $19\ \mu\text{M}$ of non-complexed clotrimazole. (**g–i**): Parasites treated with $5.1\ \mu\text{M}$ of $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$. (**b, e, h**): Hydrogenosomes labelled with the antibody anti-pyruvate:ferridoxin oxidoreductase (anti-PFOR). Treated cells (**d–i**) exhibited a reduction in fluorescence. *T. vaginalis* treated with $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ (**g–i**) displays less signal than cells treated with non-complexed CTZ (**d–f**). (**a, d, g**): Differential interference contrast (DIC). (**c, f, i**): overlay. Nuclei stained in blue with DAPI.

Hydrogenosomes

Hydrogenosomes are a good target for developing novel chemotherapies against *T. vaginalis*. Here, we observed alterations in organelle morphology (Figs 5d and 7e) and number, which was indicated by the visualization of hydrogenosomes by immunofluorescence microscopy after immunostaining cells with the anti-PFOR antibody (Fig. 8d–i) following incubation with the IC50 concentration of CTZ and its zinc complexes.

It was possible to observe hydrogenosomes with no visible peripheral vesicles (Figs 5a–d and 7e) and no spherical shape (Figs 5c–d and 7e). Immunofluorescence microscopy (Fig. 8) showed a striking difference in hydrogenosome distribution when the parasites were treated with either CTZ (Fig. 8d–f) or $[\text{Zn}(\text{CTZ})_2(\text{Ac})_2]$ (Fig. 8g–i). Non-treated cells displayed a regular distribution of hydrogenosomes and immunolabelling (Fig. 8a–c).




Fig. 9. Cytotoxic effects of clotrimazole and its zinc complexes on cultures *in vitro*. Quantitative analysis of viability of *T. vaginalis* using PI (a) and the HFF cell line using MTS (b). (a) Viability analyses of control and treated parasites were performed after 48 h of drug incubation. The following concentrations of the compounds were used: 19 μM of CTZ, 10.3 μM of [Zn(CTZ)₂Cl₂] and 5.1 μM of [Zn(CTZ)₂(Ac)₂]. No significant differences in viability are observed. The values are expressed as the mean ± s.d. for each of the three independent experiments performed in triplicate. Fifty-thousand parasites/events per sample were counted by flow cytometry. One-way ANOVA was performed to assess significance (b–d). Treatment of HFF with different concentrations of CTZ (b), [Zn(CTZ)₂Cl₂] (c) and [Zn(CTZ)₂(Ac)₂] (d) showed that the compounds were innocuous to human cells at IC₅₀ concentrations, but they become toxic at higher concentrations.

Viability assay

Quantitative viability assays were performed using PI for parasites (Fig. 9a), MTS for human cells (Fig. 9b) and *T. vaginalis* (Supplementary Fig. 3). The administration of CTZ and zinc complexes against *T. vaginalis* revealed that, for all compounds, there was no observable change in cell permeability due to membrane loss in the remaining protozoa when compared with controls and DMSO-treated cells (Fig. 9a). The plasma membrane integrity of parasites treated with these compounds for 48 h was similar to that of untreated cells. After 72 h of culture growth, 94.5% of untreated parasites were viable (Fig. 9a) as determined using PI. At 48 h following drug incubation, 96% of CTZ-treated cells, 95% of [Zn(CTZ)₂Cl₂]-treated cells and 96% of [Zn(CTZ)₂(Ac)₂]-treated cells were viable (Fig. 9a). The same profile was observed with MTS assay (Supplementary Fig. 3), there were no significant differences in the viability percentages among these treatment groups.

We performed cytotoxicity assays to evaluate the effects of CTZ and its zinc complex compounds against primary cultures of Human Foreskin Fibroblast (HFF) (Fig. 9b–d). Treatment with different concentrations of CTZ and zinc-CTZ complexes (Fig. 9b–d) showed that the compounds did not affect human cells at IC₅₀ concentrations. The Toxic Concentration of 50% (TC₅₀) and the Selective Index (SI) for HFF were 34.7 μM (SI = 1.8) for CTZ (Fig. 9b), 29.7 μM (SI = 2.9) for [Zn(CTZ)₂Cl₂] (Fig. 9c) and 28.4 μM (SI = 5.6) for [Zn(CTZ)₂(Ac)₂] (Fig. 9d).

Discussion

For many years, public health policies have neglected trichomoniasis, mainly because of the lack of information regarding the disease and the assumption that is just a nuisance infection.

Metronidazole exhibits high efficacy, and at a single 2-g oral dose, the cure rate is 90–95% (Workowski *et al.*, 2010). Despite the clinical effectiveness of 5-Nitroimidazole compounds, these drugs can lead to resistance against *T. vaginalis* (Lossick and Kent, 1991; Schwebke and Barrientes, 2006). Clotrimazole has been used as an alternative treatment for recalcitrant-Metronidazole-treated infections of *T. vaginalis in vitro* or *in vivo* when combined with other compounds (Lossick and Kent, 1991; Debbia *et al.*, 1996; du Bouchet *et al.*, 1997; Mammen-Tobin and Wilson, 2005).

Zinc salts have been previously tested against trichomonas *in vitro* (Krieger and Rein, 1982; Benchimol *et al.*, 1993; Figueroa-Angulo *et al.*, 2012; Byun *et al.*, 2015). As a result, it has been suggested that the low level of infections detected in males could be related to the urogenital microenvironment, which accumulates a higher concentration of zinc and affects the pathobiology of trichomonads (Byun *et al.*, 2015). Indeed, previous work showed that the prostatic glands have a zinc-rich environment that creates adverse conditions for *T. vaginalis* (Levy and Fair, 1973). Previous studies demonstrated that Zn²⁺ alters trichomonas virulence, and, as a consequence, affects the morphology and expression of several virulence factors in the prostatic host cells (Vazquez-Carrillo *et al.*, 2011; Figueroa-Angulo *et al.*, 2012). In this study, we showed for the first time (1) the ultrastructural alterations caused by clotrimazole in *T. vaginalis* and (2) the effectiveness of zinc-clotrimazole complexes on the parasites.

We investigated two Zn-clotrimazole complexes and their zinc salts. Our results showed that when the parasites were incubated with IC₅₀ concentrations of clotrimazole and Zn-CTZ compounds, we observed important ultrastructural alterations, but the most severe ultrastructural alterations were seen with the Zn-CTZ complex. We also obtained an IC₅₀ of 4.9 μM after

48 h of treatment with $[Zn(CTZ)_2(Ac)_2]$, which shows that the complex with $ZnAc_2$ improved CTZ activity compared to CTZ alone ($17.2 \mu M$). Although $[Zn(CTZ)_2(Ac)_2]$ is composed by two molecules of CTZ per complex, the IC₅₀ for $[Zn(CTZ)_2(Ac)_2]$ is 3 times lower than for CTZ alone, and this shows that the activity of CTZ was twice improved in the complex. Metronidazole had IC₅₀ concentrations ranging from 1.5–400 μM (Upcroft and Upcroft, 2001; Crowell *et al.*, 2003; Goodhew and Secor, 2013). In the case of the JT strain used here, it was previously shown that the IC₅₀ concentration of metronidazole was 1.8 μM (Rocha *et al.*, 2014b).

The combination of metronidazole with other drugs has been extensively tested for the treatment of *T. vaginalis* (Crowell *et al.*, 2003; Goodhew and Secor, 2013). The dual administration of zinc sulfate and metronidazole was tested, and the authors observed a decrease in the drug doses needed to treat with success recalcitrant trichomoniasis cases (Houang *et al.*, 1997). Here, we synthesized a new compound from two chemicals known to have trichomonocidal effects (clotrimazole and zinc), and both compounds are known to be active against *T. vaginalis*. We evaluated the effect of a combination of CTZ plus $ZnAc_2$ on parasite growth and compared with the zinc-clotrimazole complex. The IC₅₀ of the combined treatment after 48 h was 15.7 μM , which very close to the IC₅₀ found for CTZ alone ($17.2 \mu M$). By contrast, the $[Zn(CTZ)_2(Ac)_2]$ had a better anti-proliferative activity than the combination of CTZ and $ZnAc_2$ and was able to inhibit *T. vaginalis* with an IC₅₀ of 4.9 μM . This confirms that the significant improvement in the activity anti-*T. vaginalis* is exclusively seen when CTZ is coordinated to zinc atom, producing the coordination compound ($[Zn(CTZ)_2(Ac)_2]$).

As mentioned above, clotrimazole is reasonably safe without eliciting serious side effects when administered in humans (Crowley and Gallagher, 2014). We further investigated the effect of CTZ and its zinc derivatives on a human foreskin cell line (HFF) *in vitro* to assess cell toxicity. We observed that in addition to being more active, the complexes also had an improvement in selectivity for HFF when compared to CTZ. The Selective Index (SI) values for $[Zn(CTZ)_2(Ac)_2]$ and $[Zn(CTZ)_2Cl_2]$ were 2.9 and 5.6, respectively, while SI value for CTZ was 1.8. The SI for $[Zn(CTZ)_2(Ac)_2]$ was 3.2 times higher than CTZ. Thus both metal complexes are more tolerable for HFF in the effective concentrations than CTZ. This is evident when the IC₉₀ for $[Zn(CTZ)_2(Ac)_2]$ complex (<20 μM) is non-toxic for HFF (<30 μM), while the IC₉₀ for CTZ (>40 μM) exceeds the toxic concentration for HFF (>30 μM).

Our results demonstrated that the use of CTZ and its zinc derivatives had no deleterious effects on host cells, suggesting that these compounds may have selective anti-parasitic activity. In the case of metronidazole, there was a reduction in viable cells even at low concentrations (i.e. 1.8 μM) for HeLa cells (Rocha *et al.*, 2014a; b). Tinidazole is thought to be less toxic in human cells than metronidazole (Crowell *et al.*, 2003); however, it was demonstrated that at therapeutic concentrations (0.1, 1, 10 and 50 $\mu g mL^{-1}$), tinidazole is genotoxic, cytotoxic and can induce apoptotic cell death *in vitro* (López Nigro and Carballo, 2008).


Ultrastructural analyses of parasites indicated that the compounds used in the present work induced a number of alterations, such as the appearance of rounded cells, membrane projections and atypical organelles. It is important to highlight that we found two or more of these alterations in the same treated cells. Clotrimazole may disrupt the cell membrane, alters both membrane morphogenesis and the membrane ultrastructure (McGinnis and Rinaldi, 1991; Debbia *et al.*, 1996).

Hydrogenosomes are considered the main target organelle of drugs that affect trichomonads (Land *et al.*, 2001; Wright *et al.*, 2010; Rosa *et al.*, 2011). Sterol biosynthesis inhibitors disrupt

hydrogenosomal morphology (Rosa *et al.*, 2011). Parasites treated with metronidazole exhibited alterations in the size of this organelle (Wright *et al.*, 2010) and disorganization of hydrogenosomal proteins that act in the organelle's metabolism (Land *et al.*, 2001). Zinc treatment targets the hydrogenosomes (Benchimol *et al.*, 1993) and provokes alterations in hydrogenosomal vesicles, such as changes in electron density and size (Benchimol *et al.*, 1993). Here, we observed altered hydrogenosomes when zinc-clotrimazole complexes were used, which corroborates previous observations in the literature. These data suggest that $[Zn(CTZ)_2(Ac)_2]$ might be used to treat trichomonads. This treatment could be performed with intravaginal administration mainly by patients who have some nitroimidazole restrictions, such as nitroimidazole-resistant infections or who are allergic to nitroimidazoles.

In summary, we show that new zinc-clotrimazole complex, $[Zn(CTZ)_2(Ac)_2]$, is effective and enhanced CTZ activity against *T. vaginalis*. Although the $[Zn(CTZ)_2(Ac)_2]$ mechanism of action is still unknown, more experiments are in course to clarify this question. Our results suggest that $[Zn(CTZ)_2(Ac)_2]$ may constitute a potential and safe compound that could be used as intravaginal treatment for *T. vaginalis* infections.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S003118201900043X>

Author ORCIDs.  Victor Midlej, 0000-0001-9294-6597; Marlene Benchimol, 0000-0002-4957-8538

Acknowledgements. We acknowledge Dr Jan Tachezy from Charles University, Czech Republic for kindly provided the polyclonal antibody anti-pyruvate:ferredoxin oxidoreductase (PFOR).

Financial support. This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Programa de Núcleos de Excelência (PRONEX), Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), Universidade Federal do Rio de Janeiro (UFRJ) and by Centro Nacional de Bioimagem (CENABIO). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Conflicts of interest. There is no conflict of interest.

Ethical standards. Not applicable.

References

- Benchimol M, Almeida JC, Lins U, Gonçalves NR and de Souza W (1993) Electron microscopic study of the effect of zinc on *Trichomonas foetus*. *Antimicrobial Agents and Chemotherapy* **37**, 2722–2726.
- Betanzos-Lara S, Gómez-Ruiz C, Barrón-Sosa LR, Gracia-Mora I, Flores-Álamo M and Barba-Behrens N (2012) Cytotoxic copper(II), cobalt(II), zinc(II), and nickel(II) coordination compounds of clotrimazole. *Journal of Inorganic Biochemistry* **114**, 82–93.
- Byun JM, Jeong DH, Kim YN, Lee KB, Sung MS and Kim KT (2015) Experience of successful treatment of patients with metronidazole-resistant *Trichomonas vaginalis* with zinc sulfate: a case series. *Taiwanese Journal of Obstetrics & Gynecology* **54**, 617–620.
- Crowell AL, Sanders-Lewis KA and Secor WE (2003) *In vitro* metronidazole and tinidazole activities against metronidazole-resistant strains of *Trichomonas vaginalis*. *Antimicrobial Agents and Chemotherapy* **47**, 1407–1409.
- Crowley PD and Gallagher HC (2014) Clotrimazole as a pharmaceutical: past, present and future. *Journal of Applied Microbiology* **117**, 611–617.
- Cudmore SL, Delgaty KL, Hayward-McClelland SF, Petrin DP and Garber GE (2004) Treatment of infections caused by metronidazole-resistant *Trichomonas vaginalis*. *Clinical Microbiology Reviews* **17**, 783–793.
- du Bouchet L, Spence MR, Rein MF, Danzig MR and McCormack WM (1997) Multicenter comparison of clotrimazole vaginal tablets, oral

- metronidazole, and vaginal suppositories containing sulfanilamide, aminacrine hydrochloride, and allantoin in the treatment of symptomatic trichomoniasis. *Sexually Transmitted Diseases* **24**, 156–160.
- Debbia EU, Campora U, Massaro S, Boldrini E and Schito GC** (1996) *In vitro* activity of metronidazole alone and in combination with clotrimazole against clinical isolates of *Trichomonas vaginalis*. *Journal of Chemotherapy* **8**, 96–101.
- Edwards DI** (1993) Nitroimidazole drugs—action and resistance mechanisms. I. Mechanisms of action. *Journal of Antimicrobial Chemotherapy* **31**, 9–20.
- Figueroa-Angulo EE, Rendón-Gandarilla FJ, Puente-Rivera J, Calla-Choque JS, Cárdenas-Guerra RE, Ortega-López J, Quintas-Granados LI, Alvarez-Sánchez ME and Arroyo R** (2012) The effects of environmental factors on the virulence of *Trichomonas vaginalis*. *Microbes and Infection* **14**, 1411–1427.
- Goodhew EB and Secor WE** (2013) Drug library screening against metronidazole-sensitive and metronidazole-resistant *Trichomonas vaginalis* isolates. *Sexually Transmitted Infections* **89**, 479–484.
- Hirt RP** (2013) *Trichomonas vaginalis* virulence factors: an integrative overview. *Sexually Transmitted Infections* **89**, 439–443.
- Houang ET, Ahmet Z and Lawrence AG** (1997) Successful treatment of four patients with recalcitrant vaginal trichomoniasis with a combination of zinc sulfate douche and metronidazole therapy. *Sexually Transmitted Diseases* **24**, 116–119.
- Krieger JN and Rein MF** (1982) Zinc sensitivity of *Trichomonas vaginalis*: in vitro studies and clinical implications. *The Journal of Infectious Diseases* **146**, 341–345.
- Land KM, Clemens DL and Johnson PJ** (2001) Loss of multiple hydrogenosomal proteins associated with organelle metabolism and high-level drug resistance in trichomonads. *Experimental Parasitology* **97**, 102–110.
- Levy BJ and Fair WR** (1973) The location of antibacterial activity in the rat prostatic secretions. *Investigative Urology* **11**, 173–177.
- Lewis DA** (2010) Trichomoniasis. *Medicine* **38**, 291–293.
- López Nigro MM and Carballo MA** (2008) Genotoxicity and cell death induced by tinidazole (TNZ). *Toxicology Letters* **180**, 46–52.
- Lossick JG and Kent HL** (1991) Trichomoniasis: trends in diagnosis and management. *American Journal of Obstetrics Gynecology* **165**, 1217–1222.
- Mammen-Tobin A and Wilson JD** (2005) Management of metronidazole-resistant *Trichomonas vaginalis*—a new approach. *International Journal of STD & AIDS* **16**, 488–490.
- McGinnis MR and Rinaldi MG** (1991) Antifungal drugs: mechanism of action, drug resistance, susceptibility testing, and assay of activity in biological fluids. In Lorian V (ed), *Antibiotics in Laboratory Medicine*, 3rd edn. Baltimore, USA: Williams & Wilkins Press, pp. 198–257.
- Mielczarek E and Blaszkowska J** (2016) *Trichomonas vaginalis*: pathogenicity and potential role in human reproductive failure. *Infection* **44**, 447–458.
- Navarro M, Peña NP, Colmenares I, González T, Arsenak M and Taylor P** (2006) Synthesis and characterization of new palladium–clotrimazole and palladium–chloroquine complexes showing cytotoxicity for tumor cell lines in vitro. *Journal of Inorganic Biochemistry* **100**, 152–157.
- Navarro M, Higuera-Padilla AR, Arsenak M and Taylor P** (2009) Synthesis, characterization, DNA interaction studies and anticancer activity of platinum–clotrimazole complexes. *Transition Metal Chemistry* **34**, 869–875.
- Rocha DA, de Andrade Rosa I, de Souza W and Benchimol M** (2014a) Evaluation of the effect of miltefosine on *Trichomonas vaginalis*. *Parasitology Research* **113**, 1041–1047.
- Rocha DA, de Andrade Rosa I, Urbina JA, de Souza W and Benchimol M** (2014b) The effect of 3-(biphenyl-4-yl)-3-hydroxyquinuclidine (BPQ-OH) and metronidazole on *Trichomonas vaginalis*: a comparative study. *Parasitology Research* **113**, 2185–2197.
- Rosa IA, Rocha DA, de Souza W, Urbina JA and Benchimol M** (2011) Ultrastructural alterations induced by $\Delta(24(25))$ -sterol methyltransferase inhibitors on *Trichomonas vaginalis*. *FEMS Microbiology Letters* **315**, 72–78.
- Schwebke JR and Barrientes FJ** (2006) Prevalence of *Trichomonas vaginalis* isolates with resistance to metronidazole and tinidazole. *Antimicrobial Agents and Chemotherapy* **50**, 4209–4210.
- Secor WE** (2012) *Trichomonas vaginalis*: treatment questions and challenges. *Expert Review of Ant-Infective Therapy* **10**, 107–109.
- Upcroft JA and Upcroft P** (2001) Drug susceptibility testing of anaerobic protozoa. *Antimicrobial Agents and Chemotherapy* **45**, 1810–1814.
- Vazquez-Carrillo LI, Quintas-Granados LI, Arroyo R, Mendoza-Hernández G, González-Robles A, Carvajal-Gamez BI and Alvarez-Sánchez ME** (2011) The effect of Zn²⁺ on prostatic cell cytotoxicity caused by *Trichomonas vaginalis*. *Journal of Integrated OMICS* **1**, 198–210.
- Workowski KA and Berman S and Centers for Disease Control and Prevention** (2010). Sexually transmitted diseases treatment guidelines. *MMWR Recommendations and Reports* **59**, 1–110.
- World Health Organization** (2012) *Global Incidence and Prevalence of Selected Curable Sexually Transmitted Infections – 2008*. Geneva: WHO Press.
- Wright JM, Webb RI, O'Donoghue P, Upcroft P and Upcroft JA** (2010) Hydrogenosomes of laboratory-induced metronidazole-resistant *Trichomonas vaginalis* lines are downsized while those from clinically metronidazole-resistant isolates are not. *Journal of Eukaryotic Microbiology* **57**, 171–176.