Health Equity & Community Engagement

32460

The Bench Tutorials Program: An Essential Educational Pivot in response to COVID-19

Chantele Singleton¹, MS, MBA, Sharon A. Croisant¹, MS, PhD, Lance Hallberg¹, PhD, John Prochaska¹, DrPH, Krista Bohn¹, MPH, Michelle Puig², MEd and Cornelis Elferink¹, PhD

¹The University of Texas Medical Branch and ²Galveston Independent School District

ABSTRACT IMPACT: The Bench Tutorials Program is an independent study course in biomedical research in which high school students are paired with graduate and post-doctoral students during the academic year. The purpose is to enhance the rigor of high school science education and build the pipeline of tomorrow's researchers. OBJECTIVES/GOALS: The Bench Tutorials Program: o Proficiency in research design, implementation, and presentation; o Acquisition of hands-on laboratory skills; o Increase in scientific literacy; o Increase in analytical skills and critical thinking; o Career in science; o Build the pipeline of tomorrow's biomedical researchers METHODS/STUDY POPULATION: High School seniors are paired with graduate and postdoc mentors through a matching process. Students spend approximately four hours/week in supervised instruction and research from a participating laboratory in addition to classroom experience at their High School. Mentors design research projects relating to the larger research framework of their laboratories. In light of COVID-19, approaches have been adjusted to maintain the program safely through a hybrid method of using the high school lab for hands-on learning and through the use of Go-Pros's to enable our mentors to video and narrate as they conduct experiments in their own labs to teach their mentees scientific methods and processes. RESULTS/ANTICIPATED RESULTS: Since inception, more than 400 students and mentors have participated in the Bench Tutorial's program. This year we found a way to continue the program under COVID-19 restraints without putting anyone in harms way. Go-Pros have been essential for our program to maintain continuity for high school students who receive academic credit for this course. This program is also one of few in which our graduate students have the opportunity to serve as mentors in the scientific setting. Using Go-Pro's will also enable us to provide teaching videos online for other academic institutions, so even in the absence of COVID-19 in the future, the continued use of these devices will still be of great value. DISCUSSION/SIGNIFICANCE OF FINDINGS: High school students are afforded the ability to work on cutting edge research projects alongside graduate students and postdocs, who are afforded the chance to mentor and teach. Due to the COVID-19 pandemic, we have successfully adjusted our methods for teaching through the use of Go-Pro technology.

Mechanistic Basic to Clinical

10399

DNA-PK(cs) Regulates Stability of Egr1 During T Cell Activation

Zachary Waldrip, David Harrison, Marie Burdine and Lyle Burdine University of Arkansas for Medical Sciences, Arkansas Children's Research Institute

ABSTRACT IMPACT: This work provides supporting evidence for the development of a novel immunosuppression therapy for transplant patients. OBJECTIVES/GOALS: Our laboratory reported that inhibition of the kinase DNA-PK(cs) in mice delays allogeneic graft rejection in part by mitigating the induction of certain cytokines. We hypothesized that this was due to an inhibition of intracellular signaling programs in T cells and designed studies to identify the mechanism(s) by which this occurs. METHODS/STUDY POPULATION: The immortalized Jurkat T cell line was used to evaluate the effect of the DNA-PK(cs) inhibitor NU7441 on T cell activation by PMA/Ionomycin or PMA/PHA. Mouse primary splenocytes also were used to demonstrate the universality and reproducibility of our observations. Initially, protein mass spectrometry of lysates from untreated and NU7441-treated Jurkat cells identified proteins of interest regulated by DNA-PK(cs) that play a role in T cell activation and cytokine production. CRISPR genome editing was used to validate a potential downstream target of DNA-PK(cs). Western blot, ELISA, and flow cytometry were used to document changes in protein levels with respect to treatments. RESULTS/ ANTICIPATED RESULTS: We observed that expression of the transcription factor Egr1 was highly induced after activation but attenuated after treatment with NU7441 in both Jurkat T cells and mouse splenocytes. Phosphorylated serine 301 of Egr1 was identified by mass spectrometry in stimulated cells and fits the kinase consensus sequence for DNA-PK(cs). Both an endogenous CRISPR-generated serine 301 to alanine mutant and expression of a plasmid-based S301A mutant resulted in an unstable form of Egr1 that was barely detectable. In contrast, expression of either a S301 to D or E phospho-mimetic mutant resulted in a stable form of the protein detectable by Western blot. Further evaluation of these mutants and Egr1 phosphorylation is underway to determine the mechanism by which DNA-PK(cs) kinase regulates protein stability. DISCUSSION/SIGNIFICANCE OF FINDINGS: We previously reported a role for DNA-PK(cs) in immunomodulation. We now have evidence that this occurs in part through stabilization of Egr1. We believe this novel finding will lead to uncovering a broader role for DNA-PK(cs) as a mediator of protein stability in T cells and provide support for targeting DNA-PK(cs) in immunosuppression therapy.

12382

Circadian Disruption in Pancreatic Cancer Carcinogenesis Patrick B. Schwartz¹, MD, Morgan T. Walcheck¹, BS, Kristina A. Matkowskyj², MD, PhD, Christopher A. Bradfield³, PhD and Sean M. Ronnekleiv-Kelly⁴, MD

¹Department of Surgery, University of Wisconsin School of Medicine and Public Health, ²Department of Pathology and Laboratory Medicine, University of Wisconsin School of Medicine and Public Health, ³Department of Oncology, University of Wisconsin School of Medicine and Public Health and ⁴Department of Surgery, Division of Surgical Oncology, University of Wisconsin School of Medicine and Public Health

ABSTRACT IMPACT: Circadian disruption is known to cause significant human pathology but has not been evaluated in pancreas cancer carcinogenesis; through understanding how disruption of circadian rhythms can lead to pancreas cancer development and spread, preventive and therapeutic strategies can be devised. OBJECTIVES/ GOALS: Pancreatic ductal adenocarcinoma (PDAC) is a lethal cancer due to early spread and poor response to therapy. Identifying factors driving PDAC growth could lead to new therapeutic strategies. Thus, we evaluated the extent to which circadian rhythm disruption, a factor strongly associated with cancer formation, contributes to PDAC pathogenesis. METHODS/STUDY