

6. Public Health Laboratory Service. Incidence of surgical wound infection in England and Wales. *Lancet* 1960;2:659-663.
7. Loewenthal J. Sources and sequelae of surgical sepsis. *BMJ* 1962;1:1437-1440.
8. Cruse PJE, Foord R. The epidemiology of wound infections: a 10-year prospective study of 62,939 wounds. *Surg Clin North Am* 1980;60:27-40.
9. Green JW, Wenzel RP. Postoperative wound infection: a controlled study of the increased duration of hospital stay and direct cost of hospitalization. *Ann Surg* 1977;85:264-268.
10. Freeman J, Rosner BA, McGowan JE Jr. Adverse effects of nosocomial infection. *J Infect Dis* 1979;140:732-740.
11. Haley RW, Schaberg DR, Von Allmen SD, McGowan JE Jr. Estimating the extra charges and prolongation of hospitalization due to nosocomial infections: a comparison of methods. *J Infect Dis* 1980;141:248-257.
12. Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG. CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Infect Control Hosp Epidemiol* 1992;13:606-608.
13. Emori TG, Culver DH, Horan TC, Jarvis WR, White JW, Olson DR, et al. National Nosocomial Infections Surveillance system (NNIS): description of surveillance methods. *Am J Infect Control* 1991;19:19-35.
14. Culver DH, Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG, et al. Surgical wound infection rates by wound class, operative procedure, and patient risk index. *Am J Med* 1991;91(suppl 3B):152S-157S.
15. Rothman KJ. *Modern Epidemiology*. Boston, MA: Little, Brown & Co; 1986:275-279.
16. Hollander M, Wolfe DA. *Nonparametric Statistical Methods*. New York, NY: John Wiley & Sons; 1973:33-38.
17. Candela PV, Wenzel RP. Postoperative wound infection after total abdominal hysterectomy: a controlled study of the increased duration of hospital stay and trends in postoperative wound infection. *Am J Infect Control* 1993;21:201-204.
18. Holtz TH, Wenzel RP. Postdischarge surveillance for nosocomial wound infection: a brief review and commentary. *Am J Infect Control* 1992;20:206-213.
19. Sands K, Vineyard G, Platt R. Surgical site infections occurring after hospital discharge. *J Infect Dis* 1996;173:963-970.
20. Brown RB, Bradley S, Opitz E, Cipriani D, Pieczarka R, Sands M. Surgical wound infections documented after hospital discharge. *Am J Infect Control* 1987;15:54-58.
21. Byrne DJ, Lynch W, Napier A, Davey P, Malek M, Cuschieri A. Wound infection rates: the importance of definition and post-discharge wound surveillance. *J Hosp Infect* 1994;26:37-43.
22. Haley RW. Measuring the costs of nosocomial infections: methods for estimating economic burden on the hospital. *Am J Med* 1991;91(suppl 3B):32S-38S.
23. Davey P, Hernanz C, Lynch W, Malek M, Byrne D. Human and non-financial costs of hospital-acquired infection. *J Hosp Infect* 1991;18(suppl A):79-84.
24. Haley RW, Culver DH, White JW, Morgan WM, Emori TG, Munn VP, et al. The efficacy of infection surveillance and control programs in preventing nosocomial infections in US hospitals. *Am J Epidemiol* 1985;121:182-205.
25. Wenzel RP. Nosocomial infections, diagnosis-related groups, and study on the efficacy of nosocomial infection control: economic implications for hospitals under the prospective payment system. *Am J Med* 1985;78(suppl B):3-7.
26. Haley RW, White JW, Culver DH, Hughes JM. The financial incentive for hospitals to prevent nosocomial infections under the prospective payment system: an empirical determination from a nationally representative sample. *JAMA* 1987;257:1611-1614.
27. Centers for Disease Control. Public health focus: surveillance, prevention, and control of nosocomial infections. *MMWR* 1992;41:783-787.

## Infection Control Measures Reduce Transmission of VRE

**Gina Pugliese, RN, MS**  
**Martin S. Favero, PhD**

Montecalvo and coinvestigators from New York Medical College, Westchester Medical Center, and the CDC's Hospital Infections Program studied whether enhanced infection control strategies reduced transmission of vancomycin-resistant enterococci (VRE) in an endemic setting (an adult oncology inpatient unit). Two hundred fifty-nine patients were evaluated during use of enhanced infection control strategies, and 184 patients were evaluated during use of standard infection con-

trol practices. For enhanced interventions, patient surveillance cultures were taken, patients were assigned to geographic cohorts, nurses were assigned to patient cohorts, gowns and gloves were worn on room entry, compliance with infection control procedures was monitored, patients were educated about VRE transmission, patients taking antimicrobial agents were evaluated by an infectious disease specialist, and environmental surveillance was performed.

During use of enhanced infection-control strategies, VRE bloodstream infections decreased significantly (from 2.1 to 0.45 patients per 1,000 patient-days,  $P=.04$ ), as did VRE colo-

nization (from 20.7 to 10.3 patients per 1,000 patient-days; relative rate ratio,  $P<.001$ ). Use of all antimicrobial agents except clindamycin and amikacin was significantly reduced.

The authors concluded that enhanced infection control strategies reduced VRE transmission in an oncology unit in which VRE infections were endemic.

**FROM:** Montecalvo MA, Jarvis WR, Uman J, Shay DK, Petrullo C, Rodney K, et al. Infection-control measures reduce transmission of vancomycin-resistant enterococci in an endemic setting. *Ann Intern Med* 1999;131:269-272.