

This is a “preproof” accepted article for Weed Science. This version may be subject to change in the production process, *and does not include access to supplementary material*.

DOI: 10.1017/wsc.2025.9

**Short title: *Setaria* interseed corn alfalfa**

**Yellow foxtail (*Setaria pumila*) reduces establishment of alfalfa interseeded into corn**

Md Rayhan Shaheb<sup>1</sup>, John H. Grabber<sup>2</sup>, Marta M. Kohmann<sup>3</sup> and Mark J. Renz<sup>4</sup>

<sup>1</sup>Research Associate, Department of Plant and Agroecosystem Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

<sup>2</sup>Research Agronomist, United States Department of Agriculture – Agriculture Research Service, Dairy Forage Research Center, Madison, Wisconsin, USA

<sup>3</sup>Assistant Professor, Department of Plant and Agroecosystem Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

<sup>4</sup>Professor (<https://orcid.org/0000-0002-7558-6494>), Department of Plant and Agroecosystem Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

**Author for correspondence:** Mark Renz; E-mail: [mrenz@wisc.edu](mailto:mrenz@wisc.edu)

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

## Abstract

Interseeding alfalfa (*Medicago sativa* L.) into corn (*Zea Mays* L.) is a novel approach that increases the production of high-quality forage and reduces the risk of nutrient and soil loss from cropland. Annual grass weeds like yellow foxtail [*Setaria pumila* (Poir.) Roem. & Schult.] can reduce the success of alfalfa establishment and are difficult to manage in the interseeding system. This study evaluated ground cover, fall biomass, and fall plant density of interseeded alfalfa in response to varying populations of *S. pumila*. Our goal was to identify a threshold for initiating control of annual grasses to ensure good establishment of alfalfa in this intercropping system. Groundcover of interseeded alfalfa growing under corn declined as *S. pumila* density increased from 0 to 125 plants m<sup>-2</sup> in July, August and October with the sharpest decline in August (up to a 70% reduction in alfalfa cover). This reduction in groundcover was associated with a decline in post-establishment shoot and root mass and a reduction in alfalfa plant density from 246 to 146 plants m<sup>-2</sup> in October. Results suggest that June *S. pumila* populations should be kept < 50 plants m<sup>-2</sup> to obtain recommended fall alfalfa densities of 200 plants m<sup>-2</sup> that are needed to maximize alfalfa yield the following year. This research provides crucial information to practitioners on when annual grass management is needed to ensure successful alfalfa establishment in this interseeded system.

**Keywords:** Alfalfa-corn interseeding, crop-weed competition, threshold.

## Introduction

Milk production by dairy cows is highly dependent on diets formulated with high quality forages. Historically, corn (*Zea mays* L.) silage and alfalfa (*Medicago sativa* L.) have served as the primary sources of forage for dairy cows in the Midwestern USA (Gillespie, 2023; Kellogg et al. 2001). In Wisconsin, corn silage and alfalfa were the top two forage crops harvested in 2023 occupying 25% of the state's cropland (1.6 and 1.3 million ha, respectively) (USDA NASS 2024a). Fields are often rotated between alfalfa and corn to provide multiple production benefits such as increased forage yield of both crops, reduced pest populations, improved soil health and nutrient retention, and reduced reliance on nitrogen fertilizers (Huggins et al. 2001; Kanwar et al. 2005; Olmstead and Brummer 2008; Russelle 2014; Sanford et al. 2021). Inclusion of alfalfa in corn silage-based diets also benefits cattle health and productivity (Brito and Broderick 2006; Lopes et al. 2015).

Despite these benefits, land devoted to alfalfa production has decreased within the USA. Nationally, the area of alfalfa harvested (hay and haylage) has decreased by 32% over the past 15 years, whereas harvested corn silage increased by 15% (USDA NASS 2024a). Reductions in alfalfa harvested in Wisconsin were even greater than the national average, declining by 44% between 2009 and 2023 (USDA NASS 2024a). Alfalfa plantings have typically been replaced with less diverse production systems based on annual row crops such as corn silage, corn grown for grain, or soybean (Blum 2020; Zulauf 2018). Increasing the area where alfalfa is grown would improve the sustainability of crop and dairy production, which is aligned with national and Wisconsin priorities (Innovation Center of U.S. Dairy 2023; Wisconsin DATCP and UW System 2019).

Establishment of alfalfa by interseeding into corn silage is a novel cropping system that could be used to increase alfalfa usage on farms while providing multiple environmental benefits. In this system, alfalfa is established under a corn silage companion crop that is capable of producing up to 4.4-fold more high quality forage than conventionally spring seeded alfalfa. During its establishment, interseeded alfalfa serves as a cover crop before and after corn silage harvest and then it is harvested in subsequent years, producing forage of comparable yield as alfalfa established by conventional methods (Grabber 2016; Grabber et al. 2021b; Grabber et al. 2024). As a result, the interseeding system bypasses the low-yielding establishment year typical of conventionally spring seeded alfalfa. In addition to improved yield, this interseeding approach

has been shown to improve the profitability of alfalfa-corn silage rotations (Berti et al. 2021; Osterholz et al. 2020). Environmental benefits from this approach include reduced soil erosion, nutrient runoff (Osterholz et al. 2019), and nitrate leaching (Osterholz et al. 2021b) compared to corn silage monocultures. Because of these benefits, there are multiple efforts underway aiming to increase adoption of this interseeding system in the Midwestern USA.

Although this intercropping system provides economic and environmental benefits, stands of interseeded alfalfa can fail due to excessive competition from the corn silage companion crop and from defoliation and necrosis caused by disease and insect pests. These issues can be managed by planting corn silage at moderate populations, early seeding of well-adapted alfalfa varieties, and by application of prohexadione-calcium growth retardant and foliar fungicide and insecticide (Osterholtz et al. 2021; Grabber et al. 2021b; Grabber et al. 2023). Observations across multiple studies by our group, however, suggest high populations of annual grasses such as yellow foxtail [*Setaria pumila* (Poir.)] might also contribute to poor establishment of interseeded alfalfa. Poor alfalfa establishment due to competition with high populations of annual grasses such as *S. pumila* has been reported as a concern by others (Norris and Ayres 1991). Maintaining control of weeds during establishment is important as Chu et al. (2022) found a weed-free period of 394 growing degree days (GDD) was needed to maximize alfalfa establishment and productivity. While herbicides (e.g. acetochlor) have been identified to control early-emerging weeds including annual grasses and resulted in the successful establishment of alfalfa (Osterholz et al. 2021a), they are applied near planting and are not effective in controlling annual grasses that emerge later in the season. Roundup Ready (RR) corn and alfalfa hybrids used in conjunction with glyphosate are an effective tool to provide annual grass control in the alfalfa-corn interseeding system when applied postemergence. However, concern exists in relying on the repeated use of a single active ingredient as it increases the risk of selecting for herbicide-resistant populations. Additionally, many of the alfalfa cultivars that are best adapted to the interseeding system are conventional, for which postemergence glyphosate is not an option (Grabber et al. 2021a).

Weed densities are frequently used as decision support tools for weed management (Larson et al. 2016). While the effects of annual grasses on the establishment of interseeded alfalfa is poorly understood, its impact is likely density-dependent, and identifying its threshold is crucial to inform management decisions. In this study, the effect of annual grass weed density

on the establishment of alfalfa interseeded into corn silage was evaluated during 2023 at two locations in southern Wisconsin. *S. pumila* was selected as the model annual grass, as this late-emerging species is abundant in agronomic fields in Wisconsin (Fickett et al. 2013a, 2013b). Specifically, the impact *S. pumila* density in June had on summer ground cover, fall plant density, and fall biomass of interseeded alfalfa during the establishment year was evaluated.

## **Materials and Methods**

### ***Site Description***

Field experiments were implemented in southern Wisconsin in 2023 at the Lancaster Agricultural Research Station (LARS; 42°49' N, 90°47' W) and the United States Department of Agriculture Dairy Forage Research Center at Prairie du Sac (PDS; 43°20' N, 89°45' W). In the two years prior to experiment initiation, sites were planted with corn (LARS) or soybean followed by corn (PDS). Weed community composition included giant foxtail [*Setaria faberi* Herrm.], common ragweed (*Ambrosia artemisiifolia* L.), common lambsquarters (*Chenopodium album* L.), dandelion (*Taraxacum officinale* F. H. Wigg.), and large crabgrass [*Digitaria sanguinalis* (L.) Scop.] at LARS. At the PDS site, *C. album* and red root pigweed (*Amaranthus retroflexus* L.) were the only common species.

Soil types were Fayette (fine-silty, mixed, superactive, mesic typic hapludalfs) at LARS and Richwood (fine-silty, mixed, superactive, mesic typic argiudolls) at PDS. Before experiment establishment, soil samples were collected for characterization. Soil at LARS had 6.6 soil pH, 2.1% organic matter, 25 mg P kg<sup>-1</sup>, 106 mg K kg<sup>-1</sup>, and 0.99 g cm<sup>-3</sup> soil bulk density. Soil at PDS had 6.4 pH, 2.8% organic matter, 33 mg P kg<sup>-1</sup>, 173 mg K kg<sup>-1</sup>, and 0.94 g cm<sup>-3</sup> soil bulk density. Total rainfall and average temperatures for 2023 were obtained from nearby weather stations at Lancaster and Sauk City. Growing degree days (GDD) were calculated summing the difference between average daily temperature and alfalfa's base temperature (5 C; Sharratt et al. 1989), starting on the first day after planting. Values were averaged across both locations, as data were pooled for analysis.

### ***Experiment Establishment and Management***

Field trials in both locations were planted without tillage (no-till). Plots received glyphosate at 1 kg a.e. ha<sup>-1</sup> (Roundup PowerMax®, 540 g ai L<sup>-1</sup>, Bayer) in the last week of April before planting to control any weeds that had emerged. Based on soil analysis results, fertilizer was broadcasted pre-plant at rates of 202 kg N, 52 kg P, and 344 kg K ha<sup>-1</sup> on 27 April at LAR and on 29 April at

PDS. Corn (ARL:P0529Q; Pioneer Seed Company; LARS:BO4RR11Q-N804Q; Brevant) was planted on 4 May at LARS and 9 May at PDS. Alfalfa (431RRLH; Farm Science Company) was interseeded in four rows between each corn row the following day (5 May at LARS and 10 May at PDS) as described by Grabber et al. (2021). The seed rate of corn and alfalfa was 74,100 seeds  $\text{ha}^{-1}$  and 18 kg PLS  $\text{ha}^{-1}$ , respectively, with a row spacing of 0.76 m for corn and 0.15 m for alfalfa.

### ***Treatments and Experimental Design***

Initially, target treatments were seven *S. pumila* density ranges (0, 5-15, 30-50, 70-90, 91-100, 101-120, and  $> 120$  plants  $\text{m}^{-2}$ ) established in 3 m  $\times$  7.6 m plots arranged in a randomized complete block design with four replicates. To achieve these treatments, *S. pumila* seeds (75% germination rate) were broadcasted on 5 May at LARS and on 10 May at PDS immediately prior to alfalfa planting. Weed control in weed-free treatments was achieved by applying glyphosate at 1 kg a.e. $\text{ha}^{-1}$  (Roundup PowerMax®, 540 g ai  $\text{L}^{-1}$ , Bayer) when alfalfa had four trifoliolate leaves. Twenty-five days after planting alfalfa (DAPa), three 0.76  $\times$  0.76  $\text{m}^2$  sampling areas were marked within the center interrow of each plot for subsequent measures. In each sampling area, *S. pumila* plants were counted and thinned by hand weeding to achieve target densities. This procedure started 30 DAPa and was repeated bi-weekly thereafter until the end of June. Annual grasses that were not *S. pumila* and broadleaf weeds were also removed during this timeframe. Two irrigation events (64 and 38 mm) were applied at 31 DAPa (11 June) and 77 DAPa (27 July) at PDS due to the unusually dry spring, but no irrigation was applied in LARS due to infrastructure constraints.

To control alfalfa foliar diseases, plots were treated with fluxapyroxad at 48.6 g a.i.  $\text{ha}^{-1}$  and pyraclostrobin, and 97.4g a.i. $\text{ha}^{-1}$  (Priaxor® BASF) at 75 DAPa in both locations when the corn canopy was beginning to close (V10-V12 growth stages). Lambda-cyhalothrin at 18.2 g ai  $\text{ha}^{-1}$  (Warrior II®, Syngenta) was mixed with fungicide and applied at PDS to control potato leaf hopper [*Empoasca fabae*], but no insecticides were needed at LARS as potato leaf hopper was not present.

### ***Measurements***

#### ***A) Corn Silage***

Silage biomass was harvested on 12 and 5 September at LARS and PDS, respectively, and yields were estimated at the field level at each location. Four representative areas (23  $\text{m}^2$ ) were

harvested within each location to determine silage dry matter (DM) yield. This was done by measuring fresh weights, oven-drying sample until constant weight at 60°C, and reweighing to calculate percentage moisture. Moisture was averaged over the four samples within each location (64% and 63% in LARS and PDS, respectively) and used to correct field-level silage yield to a DM basis.

### ***B) Alfalfa and *S. pumila****

All measurements for alfalfa and *S. pumila* were conducted within the 0.76 × 0.76 m<sup>2</sup> sampling area determined 25 DAPa. At 35 DAPa, *S. pumila* density (plants m<sup>-2</sup>) was estimated. At 40, 70, 105, and 160 DAPa, percentage alfalfa and *S. pumila* cover were visually estimated. To document initial alfalfa establishment, alfalfa plants (plants m<sup>-2</sup>) were counted in early summer (40 DAPa) within one sampling area within each plot at each site (n=28). To ensure individual plants were identified, all alfalfa plants within sampling areas were destructively harvested by uprooting plants to a depth of 20 cm. Alfalfa plant density was counted in all remaining sampling areas within each plot at each site after corn silage harvest (160 DAPa; n=56). At this time alfalfa crowns were collected, then separated into shoot and root biomass. These samples were then dried at 105°C until constant weight and then weighed to determine alfalfa shoot and root biomass accumulation at the end of the growing season.

### ***Modeling and Statistical Analysis***

*S. pumila* populations varied substantially within plots and did not conform to target treatment densities. Therefore, *S. pumila* populations were considered continuous rather than categorical treatments (Kohmann et al. 2018). Regression analysis was used with the nls() function to determine the relationships between alfalfa response variables and *S. pumila* density using the R Studio platform (R Development Core Team 2024). Data across sites were visually inspected and considered similar in their responses among sites therefore were pooled for analysis. *S. pumila* density was chosen as the determinant variable, as this is a common metric used to assess weed competition, and results are easily adopted by stakeholders. For all responses, three models were compared: linear, two-parameter concave, and linear plateau, all commonly used with establishment and yield experiments (Larson et al. 2016; McCartor and Rouquette 1977; Ratkowsky 1990). The best model was selected based on a combination of visual assessment of residuals, normality, and root mean square error (RMSE).

Percentage alfalfa cover at 40, 70, 105, and 160 DAPa were best described by a two-parameter concave model shown in Eq. [1] (Ratkowsky 1990):

$$f(x) = 1/(a + bx) \quad [1]$$

where  $1/a$  is the percent alfalfa cover when *S. pumila* population is zero,  $b$  is the rate of decline of alfalfa cover with increasing *S. pumila* density, and  $x$  is *S. pumila* density in June. Fall alfalfa plant density, and shoot and root biomass at 160 DAPa were fit to a linear plateau model (McCartor and Rouquette 1977) described in Eq. [2]:

$$f(x) = a - b(x - c) \text{ if } x \leq c; \text{ otherwise, } f(x) = c \quad [2]$$

where  $a$  is the point where alfalfa plant density, shoot, or root biomass reached a plateau,  $b$  is the rate of change in those responses as *S. pumila* density increased (before reaching the plateau),  $c$  is the *S. pumila* density at the join point of the linear and plateau response of alfalfa, and  $x$  is *S. pumila* density in June.

## Results and Discussion

Temperature and precipitation during the growing season (April-November) were atypical at LARS and PDS (Table 1). Average monthly temperature was 5-16% and 3-10% greater than the 30-year average at LARS and PDS, respectively, except July which had similar temperatures as the 30-year average. Monthly precipitation was below average at both locations, and the total for the growing season amounted to only about 57% of the normal expected at LARS and PDS. With irrigation applied at PDS in June and July, the precipitation deficit was partially offset and the total for the growing season reached 70% of the expected normal. Corn silage yield at LARS and PDS were 7.5 and 16.7 Mg ha<sup>-1</sup>, respectively. In prior studies, yields of corn silage grown with interseeded alfalfa typically approach or exceed 20 Mg ha<sup>-1</sup> at these sites under near normal precipitation (Grabber et al. 2016, 2021b, 2023, 2024; Osterholtz et al. 2018).

### *Relationship between June S. pumila density and alfalfa cover*

*S. pumila* density in June ranged from 0 to 460 plants m<sup>-2</sup>. This resulted in different amounts of *S. pumila* cover, which ranged from 0 to 95% throughout the growing season. A concave function best described the relationship between *S. pumila* density and alfalfa cover in July, August, and October (Figure 1), for which all parameters were significant (P<0.0001; RMSE =12, 17, and 13, respectively). However, the parameter that describes change in alfalfa cover relative to *S. pumila* density  $b$  in June was not significant (P=0.109) and approached zero, suggesting *S. pumila* competition was not limiting alfalfa growth at that time. Chu et al. (2022)



reported similar results as the critical period for weed control for alfalfa didn't begin until late June in a weed competition study using Japanese millet [*Echinochloa esculenta* (A. Braun) H. Scholtz] as a surrogate for annual weeds in alfalfa interseeded systems in Michigan.

Competition between *S. pumila* and alfalfa was evident between July and October as the *b* parameter that described change in alfalfa cover relative to *S. pumila* density was significant ( $P < 0.0001$ ; Figure 1). The effect of June *S. pumila* density on alfalfa cover was greatest in August, when the *b* parameter was nearly 3-fold greater than in July. The pronounced increase in the *b* parameter during August likely occurred because the adverse effects of abiotic and biotic stress on stand loss and defoliation of interseeded alfalfa are most pronounced in late July and August when plants are subjected maximal shading from corn and defoliated from foliar disease and insects (Grabber et al. 2021a; Grabber et al. 2021b; Grabber et al. 2023). These effects declined by October, because corn silage harvest in early September alleviated competition and *S. pumila* did not resprout after harvest. This allowed surviving alfalfa plants, especially those subjected to additional stress from moderate to high populations of *S. pumila*, to regrow and increase ground cover. While the relationship between foxtail density and alfalfa cover was significant, alfalfa plant density is recommended for evaluating successful alfalfa establishment as the degree of stand loss and defoliation of interseeded alfalfa during late July and August and regrowth following silage harvest, is highly dependent on other factors that can't always be controlled (Osterholtz et al. 2021; Grabber et al. 2021; Grabber et al. 2023). In particular alfalfa leaf defoliation (often caused by potato leaf hopper and alfalfa foliar diseases) and fall precipitation have been identified as important factors for alfalfa plant survival during this timeframe (Grabber et al. 2021b). These factors make it difficult to rely solely on cover as an assessment of establishment success.

### ***Relationship between June S. pumila density and fall alfalfa density***

June *S. pumila* density impacted alfalfa density in October. A linear plateau function best described the relationship of all three models evaluated (RMSE = 49), with all parameters significant ( $P < 0.0001$ ; Figure 2). Alfalfa density when no weeds were present was 242 plants  $m^{-2}$  and decreased linearly with increasing June *S. pumila* density by 0.8 alfalfa plants  $m^{-2}$  for every 1 *S. pumila* plant  $m^{-2}$ . However, when June *S. pumila* density was  $\geq 125$  plants  $m^{-2}$ , alfalfa population remained constant (145 plants  $m^{-2}$ ) under the relatively dry growing conditions of this experiment.

Adequate alfalfa plant density is critical to maximize productivity in alfalfa interseeded systems. In solo-seeded alfalfa fields, densities  $> 107$  plants  $m^{-2}$  are required to maximize forage harvested (Sheaffer et al. 2023). Several studies suggest that alfalfa plant densities  $> 200$  plants  $m^{-2}$  are required by the fall of the establishment year to maximize first cut yield the following year (Osterholtz et al. 2021; Grabber et al. 2021b; Grabber et al. 2024). In our study, fall plant density of alfalfa exceeded 200 plants  $m^{-2}$  at *S. pumila* densities  $< 50$  plants  $m^{-2}$ . This threshold is similar to Zhou et al. (1992) as they found *S. pumila* densities needed to be  $< 32$  plants  $m^{-2}$  to ensure successful establishment of solo-seeded alfalfa. As noted above, this study was conducted under relatively dry growing conditions, thus *S. pumila* effects on alfalfa establishment may differ if intercropping is carried out under normal to wet growing conditions. Weed densities are frequently used as decision support tools for weed management (Larson et al. 2016) due to their easy adoption by crop consultants and farmers. We recommend assessing *S. pumila* density at 35-45 DAPa (June in Wisconsin) to determine if additional weed management activities are needed to ensure successful establishment of alfalfa in this interseeded system.

***Relationship between S. pumila density and fall alfalfa shoot and root biomass.***

Similar to alfalfa plant density at October, the relationship between June *S. pumila* density and alfalfa shoot and root biomass were best described by a linear plateau function (RMSE = 47 and 26, respectively), in which all parameters were significant ( $P < 0.0001$ ; Figure 3). Shoot and root biomass both decreased by  $0.7$  g  $m^{-2}$  for each *S. pumila* plant  $m^{-2}$ . Shoot biomass decreased until *S. pumila* density reached  $66$  plants  $m^{-2}$ , while root biomass declined and plateaued at *S. pumila* densities of  $47$  plants  $m^{-2}$ . While removal of alfalfa shoots during corn silage harvest and its regrowth likely impacted these observations, these results suggest shoot biomass is more sensitive than root biomass at higher *S. pumila* densities in this system. The impact from *S. pumila* was on alfalfa plant survival as if a plant survived it had similar root or shoot biomass per plant regardless of the *S. pumila* density (data not shown).

Results provide strong evidence that annual grasses, estimated using *S. pumila*, can reduce establishment of alfalfa interseeded into corn silage. As *S. pumila* populations increased in June, alfalfa responded by sharply reducing groundcover during July and August when competition from corn and foliar damage from disease and insects are also typically most pronounced in this intercropping system. *S. pumila* densities above  $50$  plants  $m^{-2}$  in June reduced post establishment stand density of alfalfa in October to less than  $200$  plants  $m^{-2}$ , a level

previously established as a benchmark for maximizing first cut yield of alfalfa the following year. This suggests weed control efforts should be initiated if annual grass populations exceed 50 plants m<sup>-2</sup> in June. Several options exist for annual grass weed management in the interseeded system including preemergence applications of residual herbicides (alachlor, pendimethalin) or postemergence applications of glyphosate (in RR systems only) (Osterholz et al. 2021). Additional herbicide options need to be explored for to control annual grasses that emerge after alfalfa emergence as, if conventional alfalfa varieties are used, no options are currently registered for use for late season annual grass applications in this system. *S. pumila* populations need to be below the 50 plants m<sup>-2</sup> threshold before July as reductions to alfalfa cover were observed at this timeframe (Figure 1B). As environmental conditions can affect alfalfa establishment in the interseeded system, additional research is needed to confirm the validity of this threshold. Despite this limitation, our results provide information to crop consultants and farmers as to the level and timing of management of a difficult to control and impactful weed species in this interseeded system.

### **Acknowledgments**

The authors would like to thank Charlton Rodriguez and other the students and staff from the Renz Laboratory, as well as personnel at the Lancaster Agricultural Research Station, and Mr. Matthew Volenec, Prairie du Sac Research Station for support during the establishment, maintenance, and harvest of experiments.

### **Funding**

The research work was supported by the Dairy Innovation Hub, University of Wisconsin-Madison.

### **Competing Interests**

The authors declare none.

## References

- Berti MT, Lukaszewsky J, Samarappuli DP (2021) Intercropping alfalfa into silage maize can be more profitable than maize silage followed by spring-seeded alfalfa. *Agronomy* 11(6)
- Blum M (2020) U.S. alfalfa hay production declining: Hay prices remain stable – AgriNews (agrinews-pubs.com). Last accessed on 11 March 2024
- Brito AF, Broderick, GA (2006) Effect of varying dietary ratios of alfalfa silage to corn silage on production and nitrogen utilization in lactating dairy cows. *J Dairy Sci* 89(10):3924–3938
- Chu SAD, Cassida KA, Singh MP, Burns EE (2022) Critical period of weed control in an interseeded system of corn and alfalfa. *Weed Sci* 70:680–686
- Fickett ND, Boerboom CM, Stoltenberg DE (2013a) Predicted Corn Yield Loss Due to Weed Competition Prior to Postemergence Herbicide Application on Wisconsin Farms. *Weed Technol* 27:54–62
- Fickett ND, Boerboom CM, Stoltenberg DE (2013b) Soybean yield loss potential associated with early-season weed competition across 64 site-years. *Weed Sci* 61:500–507
- Gillespie, J (2023) Feed rations fed by US dairy producers. *J Dairy Sci* 106(11): 8152–8168.
- Grabber, JH (2016) Prohexadione–calcium improves stand density and yield of alfalfa interseeded into silage corn. *Agron J* 108:726–735
- Grabber JH, Bjerneberg DL, Rogers C. (2024) Intercropping in maize silage verses solo-seeding for alfalfa establishment in Wisconsin and Idaho. *Crop Sci* 64:1061–1071
- Grabber JH, Dias JLCS, Renz MJ (2023) Establishment of Alfalfa Intercropped under Corn in Response to Varying Rates of Prohexadione with or without Fungicide Plus Insecticide. *Agronomy* 13:2823
- Grabber, J. H., Osterholz, W. R., Riday, H., Cassida, K. A., Williamson, J. A., & Renz, M. (2021a). Differential survival of alfalfa varieties interseeded into corn silage. *Crop Sci* 61: 1797–1808
- Grabber JH, Smith DL, Osterholz WR, Renz MJ (2021b) Establishment and First Year Yield of Interseeded Alfalfa as Influenced by Corn Plant Density and Treatment with Prohexadione, Fungicide and Insecticide. *Agronomy* 11:2343
- Huggins DR, Randall GW, Russelle MP (2001) Subsurface drain losses of water and nitrate following conversion of perennials to row crops. *Agron J* 93:477–486


- Innovation Center of U.S. Dairy (2023) U.S. Dairy Sustainability Report. [https://2021-2022report.usdairy.com/images/IC\\_US\\_Dairy-Sustainability\\_Report\\_2021-2022.pdf](https://2021-2022report.usdairy.com/images/IC_US_Dairy-Sustainability_Report_2021-2022.pdf) Last accessed on 1 March 2024
- Kanwar RS, Cruse RM, Ghaffarzadeh M, Bakhsh A, Karlen DL, Bailey TB, Kanwar RS (2005) Corn–soybean and alternative cropping systems effects on NO<sub>3</sub>-leaching losses in subsurface drainage water. *Appl Eng Agric* 21:181–188
- Kellogg DW, Pennington JA, Johnson ZB, Panivivat R (2001) Survey of Management Practices Used for the Highest Producing DHI Herds in the United States. *J Dairy Sci* 84:E120–E127
- Kohmann MM, Sollenberger LE, Dubeux JCB, Silveira ML, Moreno LSB, da Silva L. S, Aryal P (2018) Nitrogen fertilization and proportion of legume affect litter decomposition and nutrient return in grass pastures. *Crop Sci* 58:2138–2148
- Larson AAD, Renz MJ, Stoltenberg DE (2016) Effects of Giant Foxtail (*Setaria faberi*) and Yellow Foxtail (*Setaria pumila*) Competition on establishment and productivity of switchgrass. *Weed Sci* 64:129–136
- Lopes F, Cook DE, Combs DK (2015) Effects of varying dietary ratios of corn silage to alfalfa silage on digestion of neutral detergent fiber in lactating dairy cows. *J Dairy Sci* 98: 6291–6303
- National Weather Service (2024) National Oceanic and Atmospheric Administration. NOWData - NOAA Online Weather Data. <https://www.weather.gov/wrh/Climate?wfo=arx> Last accessed on 5 March 2024
- Norris RF, Ayres D (1991) Cutting interval and irrigation timing in alfalfa: Yellow foxtail invasion and economic analysis. *Agron J* 83:552–558
- Olmstead J, Brummer EC (2008) Benefits and barriers to perennial forage crops in Iowa corn and soybean rotations. *Renew Agric Food Syst* 23:97–107
- Osterholz WR, Dias JLCS, Grabber JH, Renz MJ (2021a) PRE- and POST-applied herbicide options for alfalfa interseeded with corn silage. *Weed Technol* 35:263–270
- Osterholz WR, Ruark MD, Renz M, Grabber JH (2021b) Benefits of alfalfa interseeding include reduced residual soil nitrate following corn production. *Agric. environ. lett* 6(3)
- Osterholz WR, Renz MJ, Grabber JH (2020) Alfalfa establishment by interseeding with silage corn projected to increase profitability of corn silage–alfalfa rotations. *Agron J* 112:4120–4132

- Osterholz WR, Renz MJ, Jokela WE, Grabber JH (2019) Interseeded alfalfa reduces soil and nutrient runoff losses during and after corn silage production. *J Soil Water Conserv* 74:85–90
- Osterholz, W.R., Renz, M.J., Lauer, J.G., & Grabber, J.H. (2018). Prohexadione-calcium rate and timing effects on alfalfa interseeded into silage maize. *Agron J* 110: 85-94
- Ratkowsky DA (1990) Handbook of nonlinear regression models (Vol. 107)
- Russelle M (2014) Environmental benefits of growing perennial legumes in cropping systems. *Legume Perspectives* 4:11–12
- Sanford GR, Jackson RD, Booth EG, Hedtcke JL, Picasso V (2021) Perenniality and diversity drive output stability and resilience in a 26-year cropping systems experiment. *Field Crops Res* 263
- Sharratt BS, Sheaffer CC, Baker DG (1989) Base temperature for the application of the growing-degree-day model to field-grown alfalfa. *Field Crops Res* 21:95-102
- Sheaffer C, Drewitz C, Jungers J (2023). Stem and plant density as predictors of alfalfa yield. *Minnesota Crop News*. University of Minnesota Extension
- USDA NASS (2024a) USDA's National Agricultural Statistics Service Wisconsin Field Office. Last accessed on 11 March 2024
- Wisconet, Wisconsin's Environmental Mesonet. <https://wisconet.wisc.edu/>. Accessed 21 July 2024
- Wisconsin DATCP and UW System (2019) Wisconsin dairy task force 2.0. Towards a vibrant Wisconsin dairy industry: Issues, insights and recommendations. <https://datcp.wi.gov/Documents/DairyTaskForce20FinalReport.pdf> Last accessed on 11 March 2024
- Zhou SQ, Hou TJ, Liu YL, Huang ZJ (1992) A study on the ecologic-economic threshold model of *Setaria viridis* (L.) Beauv. in alfalfa. *Pratacultural Science* 9: 21–23
- Zulauf C (2018) U.S. Hay Market over the Last 100 Years. <https://farmdocdaily.illinois.edu/2018/09/us-hay-market-over-the-last-100-years.html> Last accessed on 11 March 2024

**Table 1.** Monthly 2023 weather and 30-year historical averages at Lancaster (LARS) and Prairie du Sac (PDS) Wisconsin during the growing season. Irrigation was available at PDS only; total volume of water, consisting of precipitation plus irrigation, is presented for that location. Sources: Wisconet, Wisconsin’s Environmental Mesonet (2024) and National Weather Service (2024).


Month	LARS				PDS					
	Temperature (°C)		Precipitation (mm)		Temperature (°C)		Precipitation and irrigation (mm)			
	Historical average	2023	Historical average	2023	Historical average	2023	Historical average	2023	Irrigation	Total
April	8.0	8.9	90	79	7.2	7.7	100	111		111
May	14.4	15.7	114	45	14.1	15.1	108	26		26
June	19.8	20.8	142	61	19.7	20.2	149	19	64	83
July	21.7	21.7	133	113	21.7	21.5	113	101	38	139
August	20.7	21.7	102	35	20.5	21.1	113	45		45
September	16.6	18.3	108	57	16.4	18.1	93	65		65
October	9.6	11.1	79	86	9.6	10.2	75	79		79
November	2.1	2.7	52	4	2.3	2.4	53	12		12
Total	-	-	820	480	-	-	804	457	102	559

## Figures




**Figure 1.** Effect of *S. pumila* density in June ( $x$ ) on alfalfa cover at 70, 105, and 160 days after planting alfalfa (DAPa) (July, August, and October, respectively). Points represent measured responses, continuous line represents the response estimated by the regression model, and dashed lines represent the 95% confidence interval. The relationship between alfalfa ground cover and June *S. pumila* density was established using data pooled across two locations ( $n=112$ ).





**Figure 2.** Relationship between *S. pumila* plant density in June (x) and alfalfa plant density in October (160 Days After Planting alfalfa; DAPa). Points represent measured responses, continuous line represents the response estimated by the regression model, and dashed lines represent the 95% confidence interval . Data was pooled across two locations (n=112).



**Figure 3.** The relationship between *S. pumila* plant density in June ( $x$ ) and alfalfa shoot and root biomass in October (160 Days After Planting alfalfa; DAPa). Points represent measured responses, continuous line represents the response estimated by the regression model, and dashed lines represent the 95% confidence interval. Data was pooled across two locations ( $n=112$ ).