



cambridge.org/pns

The Nutrition Society Congress 2024 was held at Queen's University Belfast on 2nd–5th July 2024

Conference on 'New data – focused approaches and challenges'

Silver Medal Lecture

Review Article

Cite this article: Needham C, Partridge SR, Alston L, Rawstorn JC, and Livingstone KM (2025). Co-designing interventions to improve diets in rural communities. *Proceedings of the Nutrition Society*, page 1 of 7. doi: [10.1017/S0029665125000060](https://doi.org/10.1017/S0029665125000060)

Received: 25 October 2024

Revised: 7 January 2025

Accepted: 20 January 2025


Keywords:

Rural; Co-design; Digital intervention; Community-based intervention; Diet

Corresponding author:

Katherine Mary Livingstone;

Email: k.livingstone@deakin.edu.au

Cindy Needham¹, Stephanie R Partridge^{2,3}, Laura Alston^{4,5}, Jonathan C Rawstorn⁶ and Katherine M Livingstone⁶ 

¹Global Centre for Preventive Health and Nutrition, Institute for Health Transformation, School of Health and Social Development, Faculty of Health, Deakin University, Geelong, VIC 3220, Australia; ²Susan Wakil School of Nursing and Midwifery, Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia; ³Charles Perkins Centre, The University of Sydney, Sydney, NSW, Australia; ⁴Deakin Rural Health, School of Medicine, Faculty of Health, Deakin University, Warrnambool, Australia; ⁵Research Unit, Colac Area Health, Colac, VIC 3251, Australia and ⁶Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Deakin University, Geelong, VIC 3220, Australia

Abstract

By involving stakeholders to identify issues, co-design facilitates the creation of solutions aligned with the community's unique needs and values. However, genuine co-design with consumers across all stages of nutrition intervention research remains uncommon. The aim of this review was to examine notable examples of interventions to improve diets in rural settings that have been co-designed by rural communities. Six studies were identified reporting on community-based and digital interventions to improve diets in rural settings that have been co-designed by rural communities. The level of co-design used varied, with two interventions describing co-design workshops and focus groups over a period of between 6 and 11 months, and others not reporting details on the co-design process. Collectively, most interventions demonstrated positive impacts on dietary markers, including an increase in purchase of fruit and vegetable, an increase in percentage energy from nutrient dense foods and a decrease in intake of high fat meats. While these interventions show promise for improving diets in these under-served communities, it is widely recognised that there is a lack of dietary interventions genuinely co-designed with and for rural communities. Future research should build on these studies to co-design dietary interventions that integrate the benefits of both community-based and digital interventions.

Co-design practices

Co-design is an active collaborative approach that involves stakeholders, including community members, in creating new knowledge and solutions⁽¹⁾. Co-design aligns with traditional participatory action research by involving stakeholders in a reflective inquiry process to generate actionable knowledge⁽²⁾. However, it goes further by enabling recipients to achieve practical outcomes, such as designing and implementing interventions. A recent scoping review led by Meloncelli and colleagues on the use and extent of co-design in nutrition interventions with consumers⁽³⁾ identified co-design as aligning with the 'collaborate' and 'empower' levels on the spectrum defined by The International Association of Public Participation⁽⁴⁾. Co-design operates on the principles of valuing the expertise and lived experiences of all participants, fostering equal and reciprocal relationships⁽⁵⁾. These principles emphasise shared decision-making, where stakeholders actively contribute to identifying issues, analysing needs, defining priorities and designing and implementing interventions. This process aims to promote power-sharing, trust and mutual dependence, ensuring that interventions are contextually relevant, culturally sensitive and sustainable.

By involving stakeholders to identify issues, co-design facilitates the creation of solutions aligned with the community's unique needs and values⁽⁶⁾. This participatory approach has been heralded in the research and health communities as a valuable approach, especially for priority populations, such as rural communities, where interventions must consider local knowledge, social dynamics and available resources⁽⁶⁾. While co-design for nutrition interventions has become more prevalent in recent years, genuine partnerships with consumers across all stages of nutrition intervention research remain uncommon⁽³⁾.

Aim

The aim of this review was to examine notable examples of interventions to improve diets in rural settings that have been co-designed by rural communities.

© The Author(s), 2025. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Table 1 Examples of co-designed community-based and digitally delivered interventions in rural communities to improve dietary intakes

Study, author, year	Population	Intervention			Description	Dietary outcome	Results
		Co-design method	Co-design participants	Study design			
Community-based							
<i>Neqa Elicarvigmun</i> or the Fish-to-School Program Bersamn <i>et al.</i> 2019	<i>n</i> 76 middle and high school students, remote Alaska, USA	Two phases: focus groups and work group meetings	An Elder, tribal council members, city government, teachers, parents, high school and university students, representatives from the fishing and business development sector	9 months, school-based, multilevel intervention pre-post comparison group design	Activities in the cafeteria, classroom and community	Diet quality, fish intake and attitudes and beliefs around traditional foods (specifically fish)	Significant improvements in diet quality (Beta = 4.57; $P < 0.05$) and fish intake (Beta = 0.16; $P < 0.05$) compared to the control
<i>SHOP@RIC Stores</i> Brimblecombe <i>et al.</i> 2017 ⁽³⁶⁾	Indigenous peoples living rural and remote NT indigenous communities, Australia	Not reported	Remote food retail and public health expertise, trained local community residents, store managers and public health nutritionists who worked in the communities	6-month stepped wedge randomised controlled trial	Intervention: 20 % price discount on fruit and vegetable purchases with and without consumer education Control: Waitlist intervention (8 months).	Fruit and vegetable purchases	Price discount alone was associated with a 12.7 % (4.1–22.1) increase in purchases of fruit and vegetables
<i>Healthy Foods North</i> intervention program Kolahdooz <i>et al.</i> 2014 ⁽⁵¹⁾	Six remote communities in Nunavut and the Northwest Territories, Canada	Interviews and workshops	Inuit and Inuvialuit community members	12-month quasi-experimental intervention	Intervention: healthy breakfasts, healthy meal planning and cooking and education sessions on consuming sufficient amounts of vitamins and minerals, among other activities to promote healthy diet. Implementation sites included food stores, health clinics, offices, as well as at community special events Control: delayed intervention	Consumption of de-promoted foods: high-fat meats; high-fat dairy; refined grains; high sugar drinks; unhealthy snacks; and unhealthy additions (such as high fat powdered creamer added to coffee).	Significant decrease in de-promoted foods, such as high fat meats (–27.9 g) and high fat dairy products (–19.8 g) among intervention communities (all $P \leq 0.05$).
Digitally delivered							
<i>OL@-OR@</i> mobile health programme Mhurchu <i>et al.</i> 2019 ⁽⁴¹⁾	<i>n</i> 1451 adults in Māori and Pasifika communities, New Zealand	Focus groups and group meetings over 11 months	Māori and Pasifika communities	12-week, two-arm, cluster-randomised controlled trial	Intervention: information on healthy eating; culturally relevant information; links to local activities and services, goal setting, lifestyle trackers; Control: control version of the app	Adherence to fruit and vegetable recommendations (5 or more serves/day)	No significant improvement
<i>Healthy rural hearts</i> telehealth program Herbert <i>et al.</i> 2024 ^(43,44)	<i>n</i> 105 adults at risk of heart disease living in rural and remote New South Wales, Australia	User-centred design	Adults and dietitians based in rural and remote New South Wales (Modified Monash 3–7)	12-month cluster randomised controlled trial	Intervention: Personalised nutrition report and two hours of medical nutrition therapy from a dietitian via telehealth and usual CVD care; Control: usual CVD care and automated feedback on diet-related CVD risk factors	Percentage total energy intake (%E) derived from core/non-core foods and fats/saturated fats; daily intake of fibre and added sodium	Significant increases in %E from nutrient-dense core foods compared to the control group ($P \leq 0.05$)
<i>Veg4Me</i> study Livingstone <i>et al.</i> 2024 ^(42,45)	<i>n</i> 116 adults in regional, rural and remote Victoria, Australia	Workshops over 6 months	Young adults (18–35 years) living in regional, remote and remote Victoria (Modified Monash 2–7), local rural governments, national health promotion organisation	12-week, two arm randomised controlled trial	Intervention: personalised recipes, information on healthy eating; personalised local food environment map, goal setting; Control: non-personalised recipes and information about local food initiatives	Vegetable intakes (serves/day) and eating habits (perceived change in vegetable intake, confidence to eat vegetables, confidence to eat healthy foods)	Evaluation ongoing

Diets in rural communities

Populations who live outside major cities and urban areas, such as rural communities experience, unequal health when compared to their urban/metropolitan counterparts, with many examples around the world^(7–10). For example, remoteness influences CVD mortality above and beyond the impact of socio-economic status for those outside of major cities in Australia⁽¹¹⁾.

Minimal progress has been made in addressing rural health inequities, despite calls for action^(12,13), and a focus on rural communities as priority populations in countries such as Australia, Canada and New Zealand^(13–15). Although the factors leading to these health inequities are complex, such as inadequate access to health care, nutrition and food environments have been shown to play a role in disparities in chronic disease in rural areas^(13,16,17). Of concern, nutrition research has been under-funded and undervalued, translating into a lack of knowledge around potential approaches to improve nutrition in rural areas⁽⁸⁾. Urgent action is needed to understand potential solutions, with involvement of rural communities⁽¹⁸⁾, to prevent rural health disparities from continuing to widen.

Addressing nutrition issues such as access to healthy foods in addition to individualised nutrition care, provides a key opportunity to progress health in rural communities for present and future generations⁽¹⁹⁾. Evidence suggests that interventions addressing rural nutrition do not often include co-design or involve local community members in the implementation of interventions⁽²⁰⁾.

Co-design of dietary interventions

Community-based interventions

Growth in community-based interventions reflects a paradigm shift away from individual-focused interventions to those that emphasise the merits of addressing the complex and dynamic drivers of unhealthy behaviours⁽²¹⁾. Community-based interventions take a systems approach by employing multiple interventions addressing both systemic, social and environmental drivers of behaviour in an effort to improve health of the community⁽²²⁾. The potential effectiveness of community-based interventions on supporting healthy weight^(23,24) and improving dietary outcome measures⁽²⁵⁾ through the use of various strategies (e.g. health promotion, education, community engagement and structural changes) has been demonstrated. Emphasis has been placed on the importance of community-based interventions being context-specific⁽²⁵⁾ with research incorporating participatory approaches to co-design strategies that meet the specific needs, preferences and challenges faced by the community⁽²⁶⁾. In rural communities the need for context specific multicomponent community-based interventions is profound given the unique challenges these communities face in relation to the availability, adequacy and accessibility of foods that can support a healthy diet^(27,28). Co-design of dietary interventions could assist in addressing the interconnected public health concerns in populations undergoing a nutrition transition by developing programs that support overall well-being by taking an indigenous worldview⁽²⁹⁾.

Digital health interventions

Digital health interventions leverage a variety of information and communication technologies to reach people outside of traditional healthcare pathways. Increasingly ubiquitous ownership or access to these technologies allows delivery of flexible, accessible and personalised support to communities in almost any location⁽³⁰⁾.

While digital health interventions should not replace in-person care, they can play important complementary roles by addressing unmet needs of people who cannot access in-person interventions and providing additional support between or beyond in-person visits⁽³¹⁾. Potential to improve dietary outcomes has been demonstrated by a meta-analysis that showed digital dietary interventions improved diet quality and fruit and vegetable intake among people with chronic health conditions including CVD, diabetes mellitus and/or renal disease; moreover, there was some evidence that higher frequency interventions were more effective⁽³⁰⁾. While digital dietary interventions show promise, it is important to consider how their design and use may contribute to health inequalities. Some evidence suggests digital interventions can help to improve dietary behaviours and knowledge among disadvantaged populations⁽³²⁾; however, mechanisms underlying digital health adoption, engagement and effectiveness are not well understood⁽³³⁾. Capability, opportunity and motivation for people living in rural areas to use digital health interventions may be influenced by factors such as varied broadband coverage, digital literacy and reduced access to healthcare professionals as a source of advice about digital health interventions⁽³³⁾. Co-designing with people living in rural areas could help to ensure interventions are socially as well as geographically accessible, although this has not been common to date⁽³⁴⁾.

Overview of selected interventions

This review includes selected examples of community-based and digitally delivered interventions conducted in Australia, New Zealand, Canada and the USA. These countries were selected as they are high-income countries with distinct geographic distributions of population groups, such as indigenous peoples, that impact on diet and health inequities. This review defines community-based interventions as those delivered primarily face-to-face and digital interventions as those delivered primarily online via digital devices.

Community-based interventions

Three studies^(29,35,36) have explored the effectiveness of community-based interventions aimed at improving dietary behaviours in underserved populations, including Inuit and Inuvialuit populations in Canada and remote communities in the USA and Australia (Table 1).

In Canada, Kolahdooz and colleagues⁽³⁵⁾ evaluated the 12-month Healthy Foods North intervention program tailored to the Inuit and Inuvialuit populations (Canadian Indigenous populations) in remote and semi-remote areas in Nunavut and Northwest Territories to reduce risk of chronic disease. The intervention participants had access to healthy breakfasts, healthy meal planning and cooking and education sessions, while the control received delayed access to the intervention. The multicomponent intervention was developed through community participatory approaches including extensive stakeholder collaboration enabling the development of a culturally appropriate program⁽³⁷⁾. Key elements of Healthy Foods North included promotion of healthier food preparation methods and the benefits of traditional foods and point of purchase activities to increase the availability, accessibility and visibility of healthiness foods in stores⁽³⁵⁾. In this quasi-experimental intervention evaluation four communities (two each from Nunavut and Healthy Foods North) received the intervention and two comparison control communities (one from each Nunavut and Northwest Territories) received the delayed intervention following data collection. Pre and post food frequency questionnaires measured consumption patterns

and Adult Impact Questionnaires determined food acquisition and preparation behaviours and sociodemographic factors (both using validated surveys)⁽³⁵⁾. A significant decrease in the consumption of high-fat meat and high-fat dairy, unhealthy drinks and refined grain products and an increase in healthier food preparation was observed in the intervention group compared to the control. Control groups had a significant increase in refined grain product and unhealthy drink consumption; increase in frequency of unhealthy additions intake was reported across both groups⁽³⁵⁾. The predominantly female sample (80–82 %), lower response rates from some communities, temporal factors that relate to data collection and limited data of potential confounders may have limited the strength of findings⁽³⁵⁾.

In a stepped-wedge trial, Brimblecombe and colleagues⁽³⁶⁾ tested the effect of price discount of 20% on purchasing of fruit, vegetables, bottled water and artificially sweetened drinks with and without a consumer education strategy. The intervention was applied at the population level in remote Indigenous communities in Australia across twenty stores (*n* 20) located 20 km or more from any other food outlet in communities of concentrated socio-economic disadvantage. Using a 6-month stepped wedged randomised controlled trial, the intervention group received 20 % price discount on fruit and vegetable purchases with and without consumer education, while the control group received a waitlist intervention (8 months). Following the 24-week intervention results (objectively measured store sales 24 weeks pre/during and post intervention) indicated discount alone stores resulted in a 12.7 % (95 % CI 4.1, 22.1) increase in fruit (mostly) and vegetable purchasing⁽³⁶⁾. There were inconsistencies in the delivery of the consumer education across combined (discount and education) intervention stores, but a small benefit was observed (7.6 %; 95 % CI -3.6, 20.2) on vegetable purchasing⁽³⁶⁾. Following (24 weeks post) the removal of the discount the increased purchasing of fruit and vegetables persisted somewhat. Findings also indicated an increase in bottled water purchasing because of the discount (no difference between stores) but not diet soft drinks. Total food and drink sales, total energy of foods and sodium and increased during the intervention period (discount alone) and after.

Bersamin and colleagues⁽²⁹⁾ reported on a multi-level school-based (middle and high) intervention (Fish-to-School Program) in two remote Alaska Native communities in southwestern Alaska, USA. The intervention was designed via participatory approach⁽³⁸⁾ and included activities delivered in the classroom (lessons demonstrating the environmental, nutrition and economic impacts of local and non-local supply chain usage), community (intergenerational cooking competition) and cafeteria. 'Farm-to-schools'⁽³⁹⁾ is an approach used in North America to reconnect adolescents with local agriculture by integrating locally grown foods into the school cafeteria to improve diet. Through the participatory intervention design salmon was selected as a focus due to it being an important traditional Yup'ik food and a commercial local product that is caught and processed by Native and independently owned and operated local fish business and could be served in the school cafeteria lunches weekly⁽²⁹⁾. Being a centralised service, all schools received the cafeteria intervention and only the indirect effects of the classroom and community components were examined between baseline and 4 months (time 1) and between 4 months and 9 months (time 2 and end of intervention). Diet measured from students (intervention *n* 38 and comparison *n* 38) through a single 24 hr recall and a validated biomarker of fish and marine mammal intake indicated improved overall diet quality of the intervention participants (after time 2)

with a decline in diet quality among comparison participants. The extent to which the students were engaging with the traditional norms and values (i.e. enculturation)⁽⁴⁰⁾ was measured via 15 item-survey with outcomes indicating an increase in positive beliefs in the benefits of salmon and Yup'ik food amongst other aspects of traditional life.

Across these three studies moderate benefits of community-based interventions are demonstrated from short-term (24 weeks to 12 months) interventions^(29,35,36). All interventions involved participatory approaches in the design phase which may have contributed to their success^(29,35,36). Of interest across two of the studies was the decline in dietary health of the control group^(29,35). One study suggested that the findings may reflect a trend in people eating less healthy in general it is possible that intervention may serve to keep intervention group's diets from getting unhealthier⁽³⁵⁾. Given this insight the longer-term impact and sustainability of interventions such as these would be of interest to policymakers internationally.

Digital interventions

Three recent studies^(41–43) have explored the effectiveness of digital interventions aimed at improving dietary behaviours in under-served populations, specifically focusing on Māori and Pasifika communities in New Zealand and rural populations in Australia (Table 1). These studies leveraged digital platforms to deliver personalised nutrition advice, with varying nutrition and feasibility outcomes^(41–43).

The OL@-OR@ mobile health (mHealth) programme was designed to support healthy behaviours among Māori and Pasifika communities in New Zealand⁽⁴¹⁾. This 12-week, cluster-randomised controlled trial provided co-designed culturally relevant health information, goal-setting tools and lifestyle trackers via a mobile app and website. Although this intervention was not exclusively conducted in rural areas, a large proportion of participants were rural-dwelling. Adherence to health-related guidelines, measured using a composite health behaviour score which included fruit and vegetable intake, improved over time, however, it was not significantly different to the control group, who received a control version of the app that collected baseline and outcome data only⁽⁴¹⁾. A notable feature of this smartphone intervention was personalised health behaviour goal setting and action planning. Adherence to health-related guidelines was higher among those who engaged with the goal-setting feature (meaning they set at least one behaviour change goal); however this only represented 26 % of participants. This highlights a key challenge for multi-component mHealth interventions – even if the technology is accessible, use of individual features can vary and this may impact programme effectiveness.

The Healthy Rural Hearts telehealth program, implemented in rural New South Wales, Australia, used telehealth to deliver personalised medical nutrition therapy to individuals at risk of CVD^(43,44). The 12-month cluster randomised controlled trial tested whether a personalised nutrition report and two hours of medical nutrition therapy from a dietitian via telehealth led to changes in diet compared with usual CVD care. This intervention demonstrated the potential of digital health in improving dietary behaviours in rural populations. Participants received five telehealth consultations from Accredited Practising Dietitians over a 6-month period, with dietary intake and progress tracked using digital tools such as the Australian Eating Survey Heart version. The intervention showed significant improvements in the percentage of energy intake from nutrient-dense core foods

compared to the control group. The high attendance rate (around 90 %) for telehealth consultations and positive acceptability ratings (mean score of 9.5/10) suggest this was a feasible and well-accepted model for delivering nutrition care in rural settings. However, intentions to use telehealth consultations outside the study, if available, were lower (6.0/10). This study shows the potential of digital health interventions to improve access to personalised nutrition support in remote communities but raises concerns about long-term adherence and translation into practice.

The Veg4Me study focused on increasing vegetable intake among young adults in rural Victoria, Australia, using a personalised web app that was co-designed with rural community members^(42,45). A 12-week randomised controlled trial was conducted from August 2023 until April 2024 to determine the feasibility of the intervention. Participants in the intervention group received personalised content including tailored National Heart Foundation of Australia recipe suggestions, goal-setting features and information on local food environments (e.g. farmers markets; community gardens), while the control group received non-personalised versions. This study highlights the importance of co-design in digital health interventions, ensuring the technology is relevant and engaging for users. The personalised features, such as goal setting and behaviour change support, were tailored to individual preferences, which has been shown to be an important component of effective digital interventions⁽³⁴⁾. However, the effectiveness and engagement rates are yet to be fully evaluated, as the data analysis for the study is ongoing.

Across these studies^(41–43), the digital elements, including mobile apps, telehealth or personalised web platforms played a central role in delivering health interventions. These studies illustrate the potential for digital health tools to provide accessible, tailored support in rural and underserved populations, though challenges with user engagement and effectiveness remain. Co-designing interventions and ensuring they are personalised to meet the preferences of the specific population may improve both engagement and health outcomes. As preferences may vary within specific populations, use of individual intervention features may also vary. Digital health interventions offer a promising way to address dietary health issues but continued evaluation and iteration are necessary to optimise their impact. Understanding how varying preferences among sub-groups of rural populations can influence intervention usage may provide context to better understand how engagement impacts on effectiveness.

Recommendations for research, policy and practice

This review highlights that utilising both community-based and digital interventions will undoubtedly be key to improving the health of rural communities into the future, with each study showing improvements in rural nutritional intakes. Small changes in rural diets can have a big impact⁽⁴⁶⁾, and given the limited funding resources available to fund nutrition interventions⁽⁸⁾ researchers must maximise every opportunity to engage with rural communities through co-design.

Although this review highlights the promise of community-based and digital nutrition interventions in rural areas, limited reporting on co-design makes it challenging to understand the best ways this can be done with and for rural communities. Only three out of these six key examples, provided detailed and replicable descriptions of the co-design processes utilised in their target rural communities^(29,41,45). In line with the literature, reporting on co-design has been highlighted as being inconsistent outside the context of nutrition intervention research, and broadly across health studies⁽⁴⁷⁾.

Nutrition researchers must take steps to ensure co-design methods are systematically reported, to enable learnings and replication of methodologies across communities. Transparent and detailed reporting will also assist with upscaling and sustainability of nutrition interventions in future. This review highlights the limited evidence to guide intervention co-design and delivery in the context of rural communities. Examples in this review focused mainly on indigenous populations and although these populations are of the highest priority due to disproportionate health disparities, it still leaves much unanswered about rural communities more broadly, and the best ways to engage them. Reporting on co-design methods and community participation at all stages of the research process is of clear importance when working with indigenous communities; to ensure researchers minimise any ongoing harms of colonisation and maximise the health benefits of research in these areas. However, evidence suggests that there are still large inadequacies in engagement by researchers with indigenous communities broadly across all stages of research including co-design⁽⁴⁸⁾. This must be a key consideration for future nutrition research in rural areas, and researchers have an ethical responsibility to undertake and transparently report their progress with engagement.

Finally, rural health bodies and experts have called for place-based research and the involvement of local community members and nutrition professionals in all stages of research to ensure successful outcomes^(18,49). Rural communities have intricate social fabrics, passionate health professionals and community members that can promote ideal conditions for impactful research⁽⁵⁰⁾; but this must be harnessed by researchers to ensure the greatest impact. Although this can be challenging for researchers in the context of low funding resources, it is the ethical responsibility of researchers to ensure they engage with the rural community resources on the ground. Based on this review we recommend researchers consider reporting on both co-design methods, engagement outcomes and the effectiveness of the overall intervention to enable the highest impact and empowerment of rural communities. This will ensure that every opportunity is maximised through co-design with rural communities, with promising and much needed health impacts to follow suit.

Conclusions

While interventions discussed in this review show promise for improving diets in these under-served communities, it's widely recognised that there is a lack of dietary interventions co-designed with and for rural communities and clear reporting on co-design methods. Future research should build on these studies to co-design dietary interventions that integrate the benefits of both community-based and digitally delivered interventions.

Acknowledgment. The authors would like to thank the organisers of the 2024 UK Nutrition Society Conference for their invitation to prepare this review.

Authorship. Substantial conception and contribution to the design of this work was made by K.M.L. and C.N. K.M.L., C.N., S.R.P., L.A. and J.C.R. drafted the manuscript with all co-authors contributing critical review to drafts of the manuscript. All authors approved the final manuscript.

Financial support. KML is supported by a National Health and Medical Research Council (NHMRC) Emerging Leadership Fellowship (APP1173803).

Competing interests. The authors declare none.

Ethical statement. Not Applicable.

References

- Vargas C, Whelan J, Brimblecombe J, et al. (2022) Co-creation, co-design, co-production for public health – a perspective on definitions and distinctions. *Public Health Res Pract* 32, e2022. <http://dx.doi.org/10.17061/phrp3222211>
- Jagosh J, Macaulay AC, Pluye P, et al. (2012) Uncovering the benefits of participatory research: implications of a realist review for health research and practice. *Milbank Q* 90, 311–346.
- Meloncelli N, Young A, Christoffersen A, et al. (2023) Co-designing nutrition interventions with consumers: a scoping review. *J Hum Nutr Diet* 36, 1045–1067.
- International Association for Public Participation IAP2 Public Participation Spectrum (2018). <https://iap2.org.au/resources/spectrum/> (accessed May 2022).
- Masterson D, Areskoug Josefsson K, Robert G, et al. (2022) Mapping definitions of co-production and co-design in health and social care: a systematic scoping review providing lessons for the future. *Health Expectations* 25, 902–913.
- Singh DR, Sah RK, Simkhada B, et al. (2023) Potentials and challenges of using co-design in health services research in low- and middle-income countries. *Glob Health Res Policy* 8, 5.
- Alston L, Allender S, Peterson K, et al. (2017) Rural inequalities in the Australian burden of ischaemic heart disease: a systematic review. *Heart Lung Circ* 26, 122–133.
- Alston L, Raeside R, Jia SS, et al. (2022) Underinvestment in nutrition research for at-risk populations: an analysis of research funding awarded in Australia from 2014 to 2021. *Nutr Diet* 79, 438–446.
- Logan RI & Castaneda H (2020) Addressing health disparities in the rural united states: advocacy as caregiving among community health workers and Promotores de Salud. *Int J Environ Res Public Health* 17, 9223.
- Australian Institute of Health and Welfare (2018) Australia's Health 2018 Canberra, Australia. <https://www.aihw.gov.au/reports/australias-health/australias-health-2018/contents/indicators-of-australias-health/fruit-and-vegetable-intake> (accessed May 2022).
- Jacobs J, Peterson KL, Allender S, et al. (2018) Regional variation in cardiovascular mortality in Australia 2009–2012: the impact of remoteness and socioeconomic status. *Aust N Z J Public Health* 42, 467–473.
- Porter J, Charlton K, Tapsell L, et al. (2020) Using the Delphi process to identify priorities for Dietetic research in Australia 2020–2030. *Nutr Diet* 77, 437–443.
- Wilson R & Rourke J (2023) Report card on access to rural health care in Canada. *Rural Remote Health* 23, 8108.
- MacKinnon NJ, Emery V, Waller J, et al. (2023) Mapping health disparities in 11 high-income nations. *JAMA Netw Open* 6, e2322310.
- Ministry of Health NZ (2023) Rural Health Strategy: Ministry of Health New Zealand. <https://www.health.govt.nz/strategies-initiatives/health-strategies/rural-health-strategy> (accessed May 2022).
- Alston L, Jacobs J, Allender S, et al. (2020) A comparison of the modelled impacts on CVD mortality if attainment of public health recommendations was achieved in metropolitan and rural Australia. *Public Health Nutr* 23, 339–347.
- Melaku YA, Renzaho A, Gill TK, et al. (2019) Burden and trend of diet-related non-communicable diseases in Australia and comparison with 34 OECD countries, 1990–2015: findings from the Global Burden of Disease Study 2015. *Eur J Nutr* 58, 1299–1313.
- Alston L, Heaney S, Kent K, et al. (2023) Rural nutrition and dietetics research-Future directions. *Aust J Rural Health* 31, 1027–1031.
- Alston L, Nichols M, Allender S, et al. (2023) Dietary patterns in rural and metropolitan Australia: a cross-sectional study exploring dietary patterns, inflammation and association with cardiovascular disease risk factors. *BMJ Open* 13, e069475.
- Osborne SR, Alston LV, Bolton KA, et al. (2020) Beyond the black stump: rapid reviews of health research issues affecting regional, rural and remote Australia. *Med J Aust* 213, S3–S32.e1.
- Swinburn BA, Sacks G, Hall KD, et al. (2011) The global obesity pandemic: shaped by global drivers and local environments. *Lancet (London, England)* 378, 804–814.
- The Centre for Applied Health Evidence (2009) *Community-Based Interventions: A Rapid Review*. Victoria, Australia: Department of Health, Victoria.
- Strugnell C, Orellana L, Crooks N, et al. (2024) Healthy together Victoria and childhood obesity study: effects of a large scale, community-based cluster randomised trial of a systems thinking approach for the prevention of childhood obesity among secondary school students 2014–2016. *BMC Public Health* 24, 1–11.
- Public Health England (2019) *Whole Systems Approach to Obesity: A Guide to Support Local Approaches to Promoting a Healthy Weight*. London: Public Health England.
- Hassen HY, Sisay BG, Van Geertruyden J-P, et al. (2023) Dietary outcomes of community-based CVD preventive interventions: a systematic review and meta-analysis. *Public Health Nutr* 26, 2480–2491.
- Dhanjani S, Allen H, Varman B, et al. (2024) Community-based participatory obesity prevention interventions in rural communities: a scoping review. *Nutrients* 16, 2201.
- Whelan J, Millar L, Bell C, et al. (2018) You can't find healthy food in the bush: poor accessibility, availability and adequacy of food in rural Australia. *Int J Environ Res Public Health* 15, 2316.
- Marjanovic S, Kent K, Morrison N, et al. (2023) The impact of community-based food access strategies in high-income countries: a systematic review of randomised controlled trials. *Public Health Res Pract* 33, 1–9.
- Bersamin A, Izumi BT, Nu J, et al. (2019) Strengthening adolescents' connection to their traditional food system improves diet quality in remote Alaska Native communities: results from the Neqa Elicarvigmun Pilot Study. *Transl Behav Med* 9, 952–961.
- Kelly JT, Reidlinger DP, Hoffmann TC, et al. (2016) Telehealth methods to deliver dietary interventions in adults with chronic disease: a systematic review and meta-analysis. *Am J Clin Nutr* 104, 1693–1702.
- Rawstorn JC, Gant N, Direito A, et al. (2016) Telehealth exercise-based cardiac rehabilitation: a systematic review and meta-analysis. *Heart* 102, 1183–1192.
- Karimi N, Opie R, Crawford D, et al. (2024) Digitally delivered interventions to improve nutrition behaviors among resource-poor and ethnic minority groups with type 2 diabetes: systematic review. *J Med Internet Res* 26, e42595.
- König LM, Krukowski RA, Kuntsche E, et al. (2023) Reducing intervention- and research-induced inequalities to tackle the digital divide in health promotion. *Int J Equity Health* 22, 249.
- Livingstone KM, Rawstorn JC, Partridge SR, et al. (2023) Digital behaviour change interventions to increase vegetable intake in adults: a systematic review. *Int J Behav Nutr Phys Act* 20, 36.
- Kolahdooz F, Pakseresht M, Mead E, et al. (2014) Impact of the Healthy Foods North nutrition intervention program on Inuit and Inuvialuit food consumption and preparation methods in Canadian Arctic communities. *Nutr J* 13, 68.
- Brimblecombe J, Ferguson M, Chatfield MD, et al. (2017) Effect of a price discount and consumer education strategy on food and beverage purchases in remote Indigenous Australia: a stepped-wedge randomised controlled trial. *Lancet Public Health* 2, e82–e95.
- Gittelsohn J, Roache C, Kratzmann M, et al. (2010) Participatory research for chronic disease prevention in Inuit communities. *Am J Health Behavior* 34, 453–464.
- Nu J & Bersamin A (2017) Collaborating with Alaska native communities to design a cultural food intervention to address nutrition transition. *Prog Community Health Partnerships: Res, Educ Action* 11, 71–80.
- Joshi A & Ratcliffe MM (2012) Causal pathways linking Farm to School to childhood obesity prevention. *Childhood Obes (Print)* 8, 305–314.
- Bersamin A, Zidenberg-Cherr S, Stern JS, et al. (2007) Nutrient intakes are associated with adherence to a traditional diet among Yup'ik Eskimos living in remote Alaska Native communities: the CANHR Study. *Int J Circumpolar Health* 66, 62–70.
- Ni Mhurchu C, Te Morenga L, Tupai-Firestone R, et al. (2019) A co-designed mHealth programme to support healthy lifestyles in Māori and Pasifika peoples in New Zealand (OL@-OR@): a cluster-randomised controlled trial. *Lancet Digit Health* 1, e298–e307.

42. Livingstone KM, Rawstorn JC, Partridge SR, *et al.* (2024) Determining the feasibility of a codesigned and personalised intervention (Veg4Me) to improve vegetable intake in young adults living in rural Australian communities: protocol for a randomised controlled trial. *BMJ Open* **14**, e078001.
43. Herbert J, Schumacher T, Brown LJ, *et al.* (2024) Healthy rural hearts: the feasibility of a telehealth nutrition randomised controlled trial for rural people at risk of cardiovascular disease. *J Telemed Telecare* 1357633X241247245.
44. Herbert J, Schumacher T, Brown LJ, *et al.* (2023) Developing a telehealth medical nutrition therapy (MNT) service for adults living in rural Australia at risk of cardiovascular disease: an intervention development study. *J Hum Nutr Diet* **36**, 1782–1794.
45. Livingstone KM, Rawstorn JC, Alston L, *et al.* (2024) Co-design of a personalised digital intervention to improve vegetable intake in adults living in Australian rural communities. *BMC Public Health* **24**, 146.
46. Alston L, Peterson KL, Jacobs JP, *et al.* (2017) Quantifying the role of modifiable risk factors in the differences in cardiovascular disease mortality rates between metropolitan and rural populations in Australia: a macro-simulation modelling study. *BMJ Open* **7**, e018307.
47. Vargas C, Whelan J, Brimblecombe J, *et al.* (2022) Co-creation, co-design, co-production for public health - a perspective on definition and distinctions. *Public Health Res Pract* **32**, 3222211.
48. Burchill LJ, Kotevski A, Duke DL, *et al.* (2023) Ethics guidelines use and Indigenous governance and participation in Aboriginal and Torres Strait Islander health research: a national survey. *Med J Aust* **218**, 89–93.
49. Hooker L, Burgemeister F & Mills J (2023) Investment in community-based rural health care innovation to address health inequities in Australia. *Lancet Reg Health West Pac* **40**, 100967.
50. Richman L, Pearson J, Beasley C, *et al.* (2019) Addressing health inequalities in diverse, rural communities: an unmet need. *SSM Popul Health* **7**, 100398.
51. Kolahdooz F, Pakseresht M, Mead E, *et al.* (2014) Impact of the Healthy Foods North nutrition intervention program on Inuit and Inuvialuit food consumption and preparation methods in Canadian Arctic communities. *Nutr J* **13**, 68.