

Breakfast skipping and its relation to BMI and health-compromising behaviours among Greek adolescents

Eftymios Kapantais, Eftychia Chala, Daphne Kaklamanou*, Leonidas Lanaras, Myrto Kaklamanou and Themistoklis Tzotzas

Hellenic Medical Association for Obesity (HMAO), 46 Kifisias Avenue, Athens 115 26, Greece

Submitted 23 July 2009; Accepted 28 February 2010; First published online 8 June 2010

Abstract

Objective: The increasing prevalence of obesity worldwide is a major health concern. Our study, which is part of the First National Epidemiological Study of Obesity in Greece, aimed to assess (i) breakfast habits and their relationship to BMI in Greek adolescents, as well as (ii) breakfast skipping in relation to other health behaviours.

Design: Epidemiological survey throughout Greece. Participants completed a questionnaire concerning breakfast habits and many lifestyle parameters.

Setting: The survey was conducted in schools throughout Greece.

Subjects: Anthropometric measurements were performed on 6500 boys and 7778 girls, aged 13–19 years, from schools throughout Greece.

Results: Among both boys and girls, breakfast consumers had a lower BMI than breakfast skippers. Moreover, breakfast skippers among both boys and girls were found to smoke more than breakfast consumers. The proportion of boys and girls who ate breakfast was found to be greater among those who had never been on a diet than among those who had already experienced dieting. Leisure-time activity was greater in breakfast consumers than breakfast skippers; among boys, 71·8% of breakfast consumers walked compared with 66·4% of breakfast skippers, whereas 38·4% and 35·0%, respectively, exercised. Among girls, these percentages were 73·1% *v.* 68·7% for walking and 36·7% *v.* 31·5% for exercising.

Conclusions: We found that breakfast skipping can be part of a constellation of several unhealthy lifestyle parameters and is related to higher BMI in Greek adolescents.

Keywords

Greek Epidemiological Study
Obesity
Adolescents
Breakfast skipping

The increasing prevalence of obesity worldwide is a major public health concern^(1,2). Longitudinal studies show a direct association between increase in BMI and changes in most of the established risk factors for CVD^(3,4). It has also been shown that weight gain, as a risk factor for other disease development, is a characteristic not only of adults but also of children and adolescents⁽²⁾. The tendency of overweight children and adolescents to remain overweight throughout adulthood renders them a population more vulnerable to earlier onset of cardiovascular problems.

To reverse the ongoing obesity epidemic, it is important to access lifestyle determinants of BMI and its increase over time. Some cross-sectional^(5–8) and a few longitudinal^(9,10) studies have addressed the association between BMI and lifestyle factors. Among them, breakfast consumption habits have been shown to be strongly related to overweight and obesity^(6,11–13). In addition, persons who consume low-energy breakfast or no breakfast at all have been shown to

have higher cholesterol levels, and this applies to both adults and children⁽¹⁴⁾. Moreover, a few studies have shown that breakfast skipping could be part of a constellation of other health-compromising behaviours that compound the many deleterious effects of obesity on health^(15–19).

Eating patterns, as well as smoking, physical activity and alcohol consumption habits, develop during adolescence and follow through to adulthood, when changing deeply ingrained habits becomes increasingly difficult. Adolescence, therefore, is a critical time to encourage the development of health-promoting behaviours.

On these grounds, our objective was to determine (i) the percentage of Greek adolescents who skip breakfast, (ii) the relationship between breakfast consumption and BMI in Greek adolescents and (iii) the interrelationship between breakfast skipping and other health-compromising behaviours, such as smoking, alcohol consumption and leading a sedentary life, in Greek adolescent boys and girls.

*Corresponding author: Email d.kaklamanou@sheffield.ac.uk

Materials and methods

The present study was part of a nationwide, cross-sectional, epidemiological study designed to estimate the prevalence of obesity in the entire Greek population. The survey was conducted from February to June 2003 by physical training instructors of the school who were trained in how to take body measurements by experienced doctors, all members of the Hellenic Medical Association of Obesity, with the approval and collaboration of the Greek Ministry of Education⁽²⁰⁾.

Sample selection

The methodology for the study was previously described^(20,21). The population studied consisted of adolescent pupils aged 13–19 years from public schools throughout Greece, including the islands. Secondary schools representing various social classes were included in the selected sample. The selection method used was proportionate stratified random sampling (SRS).

From among the 3514 public secondary schools, a sample of 332 (9.45%) was randomly selected. In each school, according to SRS, all the pupils from four out of six classes participated in the study. All adolescents in the survey were measured and completed a questionnaire at their respective schools⁽²¹⁾.

Procedure

Study approval was obtained from local and state school authorities. Student and parental consent was obtained based on whether the questionnaires were completed. Physical training instructors of the school were responsible for conducting the survey after being given training and standardised criteria by doctors of the Hellenic Medical Association for Obesity. Adolescents aged 13–19 years who were living in the same household but not attending the same schools as other participants were invited to their siblings' school to have measurements taken with the aid of gym instructors. This group comprised <3% of all adolescents.

The initial sample included 16 578 adolescents, and from this sample 14 454 (6676 boys and 7778 girls) entered the study. Reasons for exclusion from the study were missing values or reporting of aberrant values in measurements, incomplete questionnaires or unreliable answers.

Adolescents had direct measurements taken for height and weight and completed a self-reported paper questionnaire. The questionnaire consisted of questions regarding breakfast consumption, both qualitative and quantitative, as well as questions regarding their ethnic origin (Greek, non-Greek), their smoking and drinking habits and their physical activity during leisure time, which was estimated through the hours spent walking, exercising and watching television (TV) each week. The questionnaire was pre-tested in a large school in Athens.

Regarding drinking habits, grams of ethanol consumed per week were calculated from the reported number of glasses of wine, beer and spirits per week, according to the alcoholic degree of each beverage (12% for wine, 5% for beer and 40% for spirits).

Participants who reported that they did not eat anything for breakfast (or only reported occasionally eating breakfast, i.e. less than two times a week) constituted the 'breakfast skipping' group. All the remaining participants constituted the 'breakfast consumption' group. Participants were asked whether they classed themselves as smokers (yes/no). Adolescents who responded with 'no' to this question were defined as non-smokers, compared with those responding with 'yes' who were defined as smokers. They were also asked individually (i) whether they watched TV; (ii) whether they exercised; (iii) whether they walked; and (iv) whether they played computer games. All these questions could be answered with a 'yes' or 'no'. If participants answered 'yes' to any of the aforementioned activities, they were also required to report how many hours per week they engaged in that activity.

Measurements

Weight was measured to the nearest 0.1 kg using the same kind of portable weight scale (Terraillon T 715; Terraillon France, Chatou, France) and with the participants in minimal clothing. Height was measured to the nearest 0.1 cm, without shoes, using a stadiometer (Seca 220; Seca, Columbia, SC, USA).

Statistical analysis

Age and BMI are expressed as mean and sd. The χ^2 test was used to analyse the differences in behaviours. To compare quantitative characteristics, we used a *t* test, where applicable, or a non-parametric (Wilcoxon) test. In addition, a logistic regression was used to test which behaviours can predict breakfast skipping. All *P* values were two-tailed. Results were considered significant at $P < 0.05$. Statistical analyses were carried out using the Statistical Package for the Social Sciences statistical software package version 11.5 (SPSS Inc., Chicago, IL, USA).

Results

Mean BMI values for adolescent boys and girls were 22.1 (sd 3.9) kg/m² and 21.1 (sd 3.4) kg/m², respectively. Table 1 describes in more detail the demographic values for each group, as well as the ethnicity of the two groups (males *v.* females). A *t* test was conducted and significant differences in BMI were found between boys and girls, $t_{(2,14\,452)} = -16.653$ ($P < 0.001$), with boys having higher BMI.

Further findings showed significant differences between genders in the percentage of smokers ($\chi^2_{(2,1)} = 20.18$, $P < 0.001$; Fig. 1) and consumption of alcohol ($\chi^2_{(1,1)} = 205.05$,

$P < 0.001$; Fig. 1). Furthermore, some minor, not statistically significant, gender differences were found in the following behaviours: walking habitually (71.1% boys *v.* 72.3% girls, $P > 0.05$), exercising systematically (37.9% boys *v.* 35.8% girls, $P > 0.05$) and watching TV regularly (97.7% boys *v.* 97.8% girls, $P > 0.05$).

As shown in Table 2, breakfast consumers had a lower BMI than breakfast skippers in Greek adolescent boys ($t_{(2,6498)} = 9.273$, $P < 0.001$). Analogous significant results were found in female breakfast consumers; they had lower BMI than breakfast skippers ($t_{(2,7776)} = 9.369$, $P < 0.001$).

There were significant gender differences in breakfast skipping, with boys being more likely to consume breakfast than girls ($\chi^2 = 33.8$, $P < 0.001$). The rate of

breakfast skipping was also found to be higher in older ages in both sexes: 14.1% of girls at 13–14 years old skip breakfast *v.* 18.8% and 18.7% of girls at 15–16 and 17–19 years old, respectively ($\chi^2 = 27.74$, $P < 0.001$). In boys, the breakfast skipping rate was found to be 9.8% in the younger ones (13–14 years old), 14.3% in the middle age group (15–16 years old) and 17.2% in the older age group (17–19 years old; $\chi^2 = 50.29$, $P < 0.001$). The difference in the breakfast skipping rate between boys and girls was most obvious in the younger adolescents and decreased from the younger to the older ages: 14.1% of girls *v.* 9.8% of boys in the younger age group skipped breakfast ($\chi^2 = 21.78$, $P < 0.001$). This difference remained in the middle age group: 18.8% of girls *v.* 14.3% of boys at the age of 15–16 years skipped breakfast ($\chi^2 = 17.31$, $P < 0.001$). In the older adolescents, however, no statistically significant difference was found in the breakfast skipping rate between the sexes: 18.7% of girls *v.* 17.2% of boys were breakfast skippers.

Regarding the impact of ethnicity on BMI and breakfast habits (Table 3), it was found that Greek boys had significantly higher BMI than non-Greek boys ($t_{(2,6465)} = -3.317$, $P < 0.001$), although they did not differ in breakfast consumption ($\chi^2 = 0.186$, $P > 0.05$). Contrary to boys, no difference was found in BMI between Greek and

Table 1 Characteristics of the adolescents studied

	Male (n 6676)		Female (n 7778)	
	Mean	SD	Mean	SD
Age	15.4	1.8	15.4	1.8
	<i>n</i>	%	<i>n</i>	%
Greek	6261	96.8	7328	97.0
Non-Greek	206	3.2	223	3.0
Missing nationality	209	—	227	—




Fig. 1 Health-influencing habits among Greek adolescents (■, boys; □, girls)

Table 3 Nationality, BMI and breakfast habits of adolescent boys and girls

	Greek		Non-Greek		<i>P</i>
	Mean	SD	Mean	SD	
BMI					
Boys	22.1	3.9	21.2	3.4	0.001
Girls	21.1	3.4	20.8	3.2	0.279
	<i>n</i>	%	<i>n</i>	%	
Breakfast consumption					
Boys					
Breakfast consumers	5286/6107	86.6	171/200	85.5	0.674
Breakfast skippers	821/6107	13.4	29/200	14.5	
Girls					
Breakfast consumers	6081/7328	83.0	176/223	78.9	0.125
Breakfast skippers	1247/7328	17.0	47/223	21.1	

Table 2 BMI and lifestyle parameters of both breakfast consumers and skippers among Greek adolescent boys and girls

	Boys				<i>P</i>	Girls				<i>P</i>
	Breakfast skippers		Breakfast consumers			Breakfast skippers		Breakfast consumers		
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	
	<i>n</i>	%	<i>n</i>	%		<i>n</i>	%	<i>n</i>	%	
BMI	23.2	4.0	21.9	3.8	0.001	21.9	3.7	20.9	3.3	0.001
Smoking	125/756	16.5	443/4489	9.9	<0.001	134/1094	12.2	430/5023	8.6	<0.001
Alcohol drinking	374/808	46.3	1914/4767	40.2	<0.001	354/1134	31.2	1502/5266	28.5	>0.071
Walking	585/881	66.4	4037/5619	71.8	<0.001	913/1329	68.7	4714/6449	73.1	<0.001
Exercising	308/881	35.0	2157/5619	38.4	>0.052	419/1329	31.5	2367/6449	36.7	<0.001
TV watching	836/863	96.9	5341/5458	97.9	>0.084	1275/1296	98.4	6124/6266	97.7	>0.171

activity as this could be estimated through walking, exercising and TV watching habits, and the results were as follows. Regarding habitual walking, boys who ate breakfast were significantly more likely to give a positive answer to walking than breakfast skippers ($\chi^2 = 10.93$, $P = 0.001$; Table 2). Hours spent walking per week, however, did not differ significantly between breakfast consumers and breakfast skippers in the subgroup of boys who answered positively in this respect (Table 4). Adolescent girls who ate breakfast were significantly more likely to report a positive answer to habitual walking in leisure time than those who skipped breakfast ($\chi^2 = 10.66$, $P < 0.001$; Table 2). As was the case with boys, hours spent walking per week did not differ significantly between breakfast eaters and breakfast skippers in the subgroup of girls who answered positively to walking (Table 4).

Regarding the habit of regular exercise, although there were no significant differences between boys who ate breakfast *v.* breakfast skippers, boys who ate breakfast reported that they exercised more than those who skipped breakfast ($\chi^2 = 3.79$, $P < 0.05$; Table 2). Furthermore, there was no significant difference in the hours spent per week on exercise between breakfast consumers and breakfast skippers in boys who answered positively to exercising (Table 4). In contrast, adolescent girls who ate breakfast regularly exercised more than did breakfast skippers ($\chi^2 = 12.84$, $P < 0.001$; Table 2). However, there was no significant difference in hours of gym attendance per week between breakfast consumers and breakfast skippers in the subgroup of girls who answered positively to exercising (Table 4).

When it came to TV watching habits, there was no significant difference with respect to breakfast consumption in adolescent boys (Table 2). Likewise, a majority of adolescent girls watched TV regularly irrespective of breakfast habits ($\chi^2 = 2.12$, $P > 0.05$; Table 2). However, in the subgroup of girls who watched TV, the number of hours spent in watching TV per week differed significantly between breakfast eaters and breakfast skippers ($t = 4.506$, $P < 0.001$; Table 4).

Many Greek adolescent boys and even more girls felt the need to control their weight through dieting; 21% of adolescent boys and 44% of adolescent girls had already experienced at least one dieting effort in the past, while 11% of boys and 23% of girls were on a diet at the time of the study. Regarding the relationship between dieting and breakfast consumption, it was found that 87% of adolescent boys who had never been on a diet in the past ate breakfast significantly more regularly compared with 82% of those who had gone on a diet previously ($\chi^2 = 22.75$, $P < 0.001$; Fig. 3). The results for adolescent girls were analogous, since 85% of those who had never been on a diet ate breakfast more regularly than only 79% of those who had been on a diet ($\chi^2 = 47.99$, $P < 0.001$; Fig. 3).




Fig. 3 Greek adolescents' breakfast habits according to weight loss (WL) diets in the past (■, breakfast consuming; □, breakfast skipping)

Finally, a logistic regression was conducted for both boys and girls; a cluster of behaviours that were examined earlier (e.g. smoking, TV, walking, exercise, alcohol consumption, diet attempts) was entered to predict breakfast skipping. In girls, after an initial analysis was performed, all the non-significant predictors were removed from the model. Four predictors remained in the second analysis. These were the hours spent in the gym (Wald = 5.16, $P < 0.05$), the hours of TV watched (Wald = 21.563, $P < 0.001$), the number of diets attempted in the past (Wald = 45.928, $P < 0.001$) and weekly ethanol consumption (Wald = 6.8, $P < 0.05$). The model was significant with $\chi^2 = 82.19$, $df = 4$, $P < 0.001$. The $-2 \log$ likelihood (LL) was 6742.72 and the Cox and Snell R^2 was 0.011, which means that 1.1% of the variance was explained. This means that spending more hours in the gym, fewer hours of watching TV, fewer previous dieting attempts and lower alcohol consumption are significant predictors of breakfast eaters in adolescent girls.

In boys, a similar approach was followed. After the initial analysis, all the non-significant predictors were removed from the model and the logistic regression was performed again. This time the significant predictors of breakfast skippers were the hours spent in the gym (Wald = 4.225, $P < 0.05$) and the number of diets attempted in the past (Wald = 21.377, $P < 0.001$). The model was statistically significant ($\chi^2 = 24.73$, $P < 0.001$) and the $-2 \log$ LL = 5074.43 (the Cox and Snell $R^2 = 0.004$). This means that it explained 0.4% of the variance, which is really low. In adolescent boys, spending more hours in the gym and fewer previous dieting attempts are significant predictors of breakfast eaters. Both logistic regressions provide some evidence that breakfast skipping is associated with unhealthy behaviours such as watching TV for a long time, not exercising, a lot of attempts in dieting and drinking a lot of alcohol.

Discussion

Many studies have shown a decline in the rate of breakfast consumption over the years, which is the case mainly with American adolescents^(22,23). In Europe, the extent of breakfast skipping varies from 12% to 28%^(24,25) (12% in North Italian pubertal boys, 28% in North Italian pubertal girls, 13.8% in 15–16-year-old Dutch boys, 18.2% in 15–16-year-old Dutch girls). Another Greek study⁽²⁶⁾, which was conducted in a limited sample of Greek students from the city of Piraeus, came up with analogous results on breakfast skipping (13.7% of boys ate very rarely and 2.9% never ate breakfast on weekdays, while 14.8% of girls ate very rarely and 5.2% never ate breakfast on weekdays). The present study, however, has limitations in that it was conducted in older adolescents and young adults within the limits of a certain city.

Our results are more consistent with the findings of an Australian survey conducted among 13-year-olds concerning the extent of skipping breakfast. Moreover, as was the case in Australia and in the above-mentioned studies, Greek adolescent girls seem to skip breakfast more often than do boys⁽²⁷⁾.

In our study, breakfast skipping rates were found to be higher in older than in younger ages in both sexes, which could have two possible explanations: (i) the parental influence is more obvious in younger adolescents who are either forced or persuaded by their parents to have breakfast, or simply tend to mimic their parents, while older adolescents tend to differentiate themselves from the rest of the family and to follow the habits of their friends of waking up late and leaving for school on an empty stomach. (ii) As the adolescents get older, they are more and more concerned about their image and falsely believe that by skipping breakfast they may reduce their energy intake. This hypothesis may also explain why young women, even from very early adolescence, skip breakfast more often than do boys, since it is generally known that girls tend to be more obsessed about their weight at an earlier age.

Furthermore, several studies among children and adults have shown an inverse association between breakfast consumption and BMI^(5–10). To our knowledge, this is the first national analysis of breakfast habits in adolescents in Greece and its findings are consistent with studies conducted in other countries.

Several mechanisms to explain this putative effect of breakfast consumption on BMI have been postulated. Breakfast consumption could contribute to a greater meal frequency per day, which could lead to less efficient energy use by increasing dietary-induced thermogenesis, and hence result in a lower BMI⁽²⁸⁾. Moreover, breakfast consumption may prevent overeating during the day. Ghrelin levels rise⁽²⁹⁾ and insulin levels decline during prolonged fasting⁽³⁰⁾, which may trigger hunger and stimulate eating. It has also been shown that persons who

consume a high-energy breakfast derive more daily energy from carbohydrates and less from fat than those who consume either a low-energy breakfast or no breakfast at all. This could lead to better weight control for breakfast consumers⁽³¹⁾.

On the other hand, one could propose a different point of view concerning breakfast-eating habits. It could be hypothesised that adolescents who already have problems controlling their weight skip breakfast in order to limit their total daily energy intake.

On these grounds, one could propose two different explanations for our finding concerning the relationship between breakfast consumption habits and past dieting experience. Specifically, in both boys and girls who have been in need of a diet in the past, breakfast consumption was lower than in those who had never been on a diet. This could reflect either (i) the positive impact of breakfast consumption on weight control, thus reducing the need for dieting, or (ii) the conscious effort of adolescents with weight problems to limit their total daily energy intake by skipping breakfast in an attempt to lose weight.

In our study, we also found that skipping breakfast can be part of a constellation of several unhealthy lifestyle parameters, namely smoking, alcohol drinking and a more sedentary lifestyle, as can be estimated from leisure-time activities. This finding is consistent with other studies in which breakfast skipping in adolescents has also been associated with various health-compromising behaviours and unhealthy lifestyles^(20–23). Other studies have also shown that breakfast skippers, even after controlling for age, gender and BMI, have significantly higher total cholesterol levels than breakfast consumers. If that is verified by further study, then the constellation of higher BMI together with higher total cholesterol levels and the above-mentioned health-compromising behaviours in breakfast skippers could be detrimental to the health of adolescents⁽²⁸⁾.

Thus, it seems that skipping breakfast reflects more than just meal timing preferences; it appears to be, in fact, one component of frequently co-occurring health-compromising behaviours. Adolescents who skip breakfast may care less about their health than those who eat breakfast regularly. On these grounds, we could propose a different approach to the detection of dangerous behaviours in adolescents through questioning about their breakfast habits. Breakfast skipping is easily admitted, while alcohol consumption or smoking could be more difficult to assess since adolescents may feel ashamed about them. However, further evidence is needed to show that this is possible, as the present study's results cannot be generalised due to the low R^2 in the logistic regressions.

Moreover, smoking is sometimes used to augment dieting, since it has been shown to increase energy expenditure slightly. Furthermore, smoking, which was more common in our study among breakfast skippers than breakfast consumers, may either suppress one's appetite in the morning

or reduce the time available for a healthy first meal. Even if smoking were to be used in that sense, it would not prove fruitful according to our findings. As was shown by multiple regression analysis, there was no influence of smoking on BMI among boys, while among girls, smokers actually had a greater BMI than non-smokers.

Simple nutritional interventions concerning breakfast eating may fail to address these more complex issues. The most effective strategy for influencing the breakfast habits of adolescents is probably to create family atmospheres that endorse generally health-conscious lifestyles. This healthy family environment should be present even during the early years of life, since, as we have previously proposed, parental influence on the eating habits of offspring is likely to be age-specific and the eating habits of teenagers are less likely to be under parental control than those of young children.

Acknowledgements

The study was funded by the pharmaceutical companies Abbott Laboratories Hellas and Roche Hellas. There is no conflict of interest to declare. E.K. wrote the introduction and discussion sections. E.C. wrote the introduction and methods sections. D.K. wrote the results section and helped with the discussion. The rest of the authors helped in data collection, distributing questionnaires, etc. The authors would like to extend their sincere gratitude to the Hellenic Ministry of Education for approval of the project and their collaboration, as well as to the directors and gym instructors of the various schools for their active participation in the survey.

References

- World Health Organization (1998) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation on Obesity, Geneva, 3–5 June 1997*. Geneva: WHO.
- World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series no. 916*. Geneva: WHO.
- Jousilahti P, Tuomilehto J, Vartiainen E *et al.* (1996) Body weight, cardiovascular risk factors and coronary mortality: a 15-year follow-up of middle aged men and women in eastern Finland. *Circulation* **93**, 1372–1379.
- Lamon-Fava S, Wilson PW & Schaefer EJ (1996) Impact of body mass index on coronary heart disease risk factors in men and women. The Framingham Offspring Study. *Arterioscler Thromb Vasc Biol* **16**, 1509–1515.
- Jeffery RW, Forster JL, Folsom AR *et al.* (1989) The relationship between social status and body mass index in the Minnesota Heart Health Program. *Int J Obes* **13**, 59–67.
- Cho S, Dietrich M, Brown CJ *et al.* (2003) The effect of breakfast type on total daily energy intake and body mass index: results from the Third National Health and Nutrition Examination Survey (NHANES III). *J Am Coll Nutr* **22**, 296–302.
- Rasky E, Stronegger WJ & Freidl W (1996) The relationship between body weight and patterns of smoking in women and men. *Int J Epidemiol* **25**, 1208–1212.
- Jacobsen BK & Thelle DS (1987) The Tromsø Heart-Study – the relationship between food-habits and the body mass index. *J Chronic Dis* **40**, 795–800.
- Sundquist J & Johansson SE (1998) The influence of socioeconomic status, ethnicity and lifestyle on body mass index in a longitudinal study. *Int J Epidemiol* **27**, 57–63.
- Wilsgaard T, Jacobsen B & Arnesen E (2005) Determining lifestyle correlates of body mass index using multilevel analyses: The Tromsø Study 1979–2001. *Am J Epidemiol* **162**, 1179–1188.
- Ma Y, Bertone ER, Stanek 3rd EJ *et al.* (2003) Association between eating patterns and obesity in a free-living US adult population. *Am J Epidemiol* **158**, 85–92.
- Song WO, Chun OK, Obayashi S *et al.* (2005) Is consumption of breakfast associated with body mass index in US adults? *J Am Diet Assoc* **105**, 1373–1382.
- Van der Heijden AA, Hu FB, Rimm EB *et al.* (2007) A prospective study of breakfast consumption and weight gain among US men. *Obesity (Silver Spring)* **15**, 2463–2469.
- Resnicow K (1991) The relationship between breakfast habits and plasma cholesterol levels in schoolchildren. *J Sch Health* **61**, 81–85.
- Revicki D, Sobal J & Deforge B (1991) Smoking status and the practice of other unhealthy behaviors. *Fam Med* **23**, 361–364.
- Isralowitz RE & Trostler N (1996) Substance use: toward an understanding of its relation to nutrition-related attitudes and behavior among Israeli high school youth. *J Adolesc Health* **19**, 184–189.
- Höglund D, Samuelson G & Mark A (1998) Food habits in Swedish adolescents in relation to socioeconomic conditions. *Eur J Clin Nutr* **52**, 784–789.
- Keski-Rahkonen A, Kaprio J, Rissanen A *et al.* (2003) Breakfast skipping and health-compromising behaviors in adolescents and adults. *Eur J Clin Nutr* **57**, 842–853.
- Burke V, Milligan R, Beilin LJ *et al.* (1997) Clustering of health-related behaviours among 18-year old Australians. *Prev Med* **26**, 724–733.
- Kapantais E, Tzotzas T, Ioannidis I *et al.* (2006) First national epidemiological survey on the prevalence of obesity and abdominal fat distribution in Greek adults. *Ann Nutr Metab* **50**, 330–338.
- Tzotzas T, Kapantais E, Tziomalos K *et al.* (2008) Epidemiological survey for the prevalence of overweight and abdominal obesity in Greek adolescents. *Obesity (Silver Spring)* **16**, 1718–1722.
- Siege-Riz A, Popkin B & Carson T (1998) Trends in breakfast consumption for children in the United States from 1965 to 1991. *Am J Clin Nutr* **76**, Suppl. 4, S748–S756.
- Cohen B, Evers S, Manske S *et al.* (2003) Smoking, physical activity and breakfast consumption among secondary school students in a Southwestern Ontario community. *Can J Public Health* **94**, 41–44.
- Vanelli M, Iovane B, Bernardini A *et al.*; Students of the Post-Graduate School of Paediatrics, University of Parma (2005). Breakfast habits of 1202 Northern Italian children admitted to a summer sport school. Breakfast skipping is associated with overweight and obesity. *Acta Biomed* **76**, 79–85.
- Croezen S, Visscher TLS, Ter Bogt NCW *et al.* (2009) Skipping breakfast, alcohol consumption and physical inactivity as risk factors for overweight and obesity in adolescents: results of the E-MOVO project. *Eur J Clin Nutr* **63**, 405–412.

26. Dialektakou KD & Vranas PB (2008) Breakfast skipping and body mass index among adolescents in Greece: whether an association exists depends on how breakfast skipping is defined. *J Am Diet Assoc* **108**, 1517–1525.
27. Shaw ME (1998) Adolescent breakfast skipping: an Australian study. *Adolescence* **132**, 851–861.
28. Drummond S, Crombie N & Kirk T (1996) A critique of the effects of snacking on body weight status. *Eur J Clin Nutr* **50**, 779–783.
29. Cummings DE, Purnell JQ, Frayo RS *et al.* (2001) A preprandial rise in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes* **50**, 1714–1719.
30. Boyle PJ, Shah SD & Cryer PE (1989) Insulin, glucagon and catecholamines in prevention of hypoglycemia during fasting. *Am J Physiol* **256**, E651–E661.
31. Preziosi P, Galan P, Deheeger M *et al.* (1999) Breakfast type, daily nutrient intakes and vitamin and mineral status of French children, adolescents and adults. *J Am Coll Nutr* **18**, 171–178.