ticularly in *S* agalactiae infections. In the early postpartum period, fever in the mothers was significantly less likely in the patients offered vaginal disinfection, a reduction from 7% in those douched using saline compared with 3% in those disinfected using chlorhexidine. A lower occurrence of urinary tract infections also was observed: 6% in the saline group as compared with 3% in the chlorhexidine group (P<.01).

This prospective controlled trial demonstrated that vaginal douching with 0.2% chlorhexidine during labor can significantly reduce both maternal and early neonatal infectious morbidity. The squeeze bottle procedure was simple, quick, and well-tolerated.

FROM: Stray-Pedersen B, Bergan T, Hafstad A, Normann E, Grogaard J, Vangdal M. Vaginal disinfection with chlorhexidine during childbirth. *Int J Antimicrob Agents* 1999;12:245-251.

Effectiveness of Live, Attenuated Intranasal Influenza Virus Vaccine

A recent study by Nichol and colleagues concluded that, among healthy adults, a live, attenuated influenza vaccine delivered intranasally not only helps prevent serious illness but also saves money.

In a randomized, double-blinded, placebo-controlled trial of 4,561 healthy adults aged 18 to 64, investigators found that recipients of intranasally administered trivalent, live, attenuated influenza virus (LAIV) vaccine were as likely to experience one or more febrile illnesses as placebo recipients during peak outbreak periods (13.2% for vaccine vs 14.6% for placebo). However, vaccination significantly reduced the numbers of severe febrile illnesses (18.8% reduction) and febrile upper respiratory tract illnesses (23.6% reduction). Vaccination also led to fewer days of illness across all illness syndromes (22.9% reduction for febrile illnesses; 27.3% reduction for severe febrile illnesses), fewer days of work lost (17.9% reduction for severe febrile illnesses; 28.4% reduction for febrile upper respiratory tract illnesses), and fewer days with healthcareprovider visits (24.8% reduction for severe febrile illnesses; 40.9% reduction for febrile upper respiratory tract illnesses). Use of prescription antibiotics and over-the-counter medications was also reduced across all illness syndromes. Vaccine recipients were more likely to experience runny nose or sore throat during the first 7 days after vaccination, but serious adverse events between the groups were not significantly different.

The match between the type A(H3N2) vaccine strain and the predominant circulating virus strain (A/Sydney/05/97[H3N2]) for the 1997/98 season was poor, suggesting that LAIV provided substantial crossprotection against this variant influenza A virus strain. The authors concluded that intranasal trivalent LAIV vaccine was safe and effective in healthy, working adults in a year in which a drifted influenza A virus predominated.

FROM: Nichol KL, Mendelman PM, Mallon KP, Jackson LA, Gorse GJ, Belshe RB, et al. Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial. *JAMA* 1999;282:137-144.

Gastrointestinal Endoscopic Reprocessing Practices in the United States

Patient infection from contaminated gastrointestinal (GI) endoscopes generally can be attributed to failure to follow appropriate reprocessing guidelines. Recently, the Food and Drug Administration recommended a 45-minute exposure of GI endoscopes to 2.4% glutaraldehyde solutions heated to 25°C. Simultaneously, the American Society for Gastrointestinal Endoscopy (ASGE), the American Gastroenterological Association, and the Society of Gastroenterology Nurses and Associates endorsed a reprocessing guideline that emphasized manual precleaning and recommended a 20-minute exposure to a 2.4% glutaraldehyde solution at room temperature. Since then, little information has become available regarding actual reprocessing practices in the United States.

Cheung and colleagues mailed a questionnaire regarding endoscopic disinfection practices to 730 randomly selected members of the ASGE; 294 (40%) responded. Appropriate manual cleaning (suctioning detergent through the accessory channel and brushing the channel and valves) was reported by 91% of respondents; 70% then used automated reprocessors for disinfection or sterilization. Glutaraldehyde was the most widely used chemical disinfectant; 85% used glutaraldehyde as one of their primary disinfectants. The most commonly used disinfection time with 2.4% glutaraldehyde was 20 minutes (83.9%) followed by 45 minutes (11.4%). Only 24% of users of 2.4% glutaraldehyde heated their solution; 60% of centers tested disinfectant concentration daily or more frequently; 74% sterilized nondisposable forceps before use; 29% of centers reused disposable endoscopic accessories (which are more frequently disinfected rather than sterilized). Twelve respondents reported cases of endoscopic cross-infection.

The authors note that a significant minority of endoscopy centers still do not completely conform to recent ASGE, American Gastroenterological Association, and the Society of Gastroenterology Nurses and Associates guidelines on disinfection, and they may not be appropriately disinfecting GI endoscopes. Rigid adherence to recommended guidelines is strongly encouraged to ensure patient safety.

FROM: Cheung RJ, Ortiz D, DiMarino AJ Jr. GI endoscopic reprocessing practices in the United States. *Gastrointest Endosc* 1999;50:362-368.

The School of Medicine at the University of Washington

is recruiting a full-time faculty member at the Assistant or Associate Professor level to lead the program in Infection Control at the University of Washington Medical Center. The closing date for applications is December 10, 1999.

Although the School and Medical Center are seeking a person at the Assistant or Associate Professor rank, under unusual circumstances and commensurate with the qualifications of the individual, appointment may be made at the rank of Professor.

Desirable credentials include national board certification in the candidate's specialty and ABIM eligibility in infectious diseases or advanced training in epidemiology, microbiology or related fields. Must have a strong clinical background with training in hospital epidemiology with emphasis on infection control. The selected individual will direct infection control activities of both ambulatory and inpatient care at the University of Washington Medical Center (including aspects of the new Seattle Cancer Care Alliance) and will participate in the clinical and teaching activities of their academic department. Development or continuation of an infection control or hospital epidemiology research program is desirable.

Send C.V. to:

Henry Rosen, M.D., Associate Chair Department of Medicine, Box 356420 University of Washington, Seattle, WA 98195-6420

The University of Washington is building a culturally diverse faculty and strongly encourages applications from female and minority candidates. The University is an Equal Opportunity/Affirmative Action employer.

Expand Your Horizons in Epidemiology Take An Inside Look At Cedars

For resources and challenges that will keep you on the leading edge of your field, take an inside look at Cedars-Sinai Medical Center. We're a world-class tertiary, acute care teaching medical center offering supportive management, a variety of continuing education opportunities and the assistance of a career counselor.

NURSE EPIDEMIOLOGIST

Join our team and provide expert leadership in infection control and epidemiology. You will develop and implement didactic programs as well as provide direction, consultation and development in relation to infection control and epidemiology. In addition, you will plan, coordinate and implement a system for prevention, surveillance, investigation and control of infection. Requires a Bachelor's degree (Master's preferred), CA RN license and CIC. Must have 3-5 years acute hospital experience, (preferably including one year as a nurse epidemiologist). Strong verbal/written English, basic computer and database management skills are essential.

We offer a competitive salary and benefits package. Please fax or mail your resume to: Cedars-Sinai Medical Center, Attn: Harry KaKliKian (include Job #1981956), 8723 Alden Dr., HR Dept., Room 110, Los Angeles, CA 90048, FAX: (310) 423-0374. AA/EEO.



B. Alterative Till -	2 But Keatten Munhas	9 Effect Date
t, Puzikolizon tenj	2. Publication Number	IS. Faing Date
Interiori Control and Hospital Epidemiology	0899-823X	October 1, 1999
Lase Frequency	5. Number of Issues Published Annually	6. Annual Subscription Price
<i>Monthly</i>	12	\$106.00
7. Complete Mailing Address of Known Office of Publica	tion (Not Printer) (Street, city, county, state, and ZIP+4)	Contact Person
900 Grove Road		Lester Robeson
Thoronare, Gloucester County, New Jersey 08085-	9447	Talephone
Complete Mailing Address of Headquarters or General	Business Office of Publisher (Not printer)	1(856) 548-1000 axtension 245
lick, incorporated 6900 Grove Road Thorofare, New	Jereay 08086-9447	
Ful Names and Complete Mailing Addresses of Publ	sher, Editor, and Managing Editor (Do not leave blank)
Williher (Name and complete mailing address)		
Richard N. Roash. Sleck Incorporated. 6900 Grove Ro	ad Thorofare. New Jersey 08086-9447	
ballor (name and complete mailing address)		
ilichael D. Decker, MD,. MPH	Same as above	
Managing Editor (Name and complete mailing address)		
Shirley Strunk	Same as above	
hill Hame	Complete Mailing Address	
PNS & W8S, Incorporated dba Slack, Incorporated	6900 Grove Road, Thorotere, New Jerse	/ 08086-9447
Peter N. Slack	Same as Above	
N. Semuel Slack	Same as Above	
11. Know Bondholders, Mortgagers, and Other Security		
	Holders Owning or	
Inding 1 Percent or More of Total Amount of Bonds, Ma Other Recurities, If none, check box	Holders Owning or orlgages, or	
loiding 1 Percent or More of Total Amount of Bonds, M The Becurities, If none, check box	Holders Owning or stigages, or Complete Mailing Address	
Holding 1 Percent or More of Total Amount of Bonde, M Other Bacunties. If none, check box Fall Neme	Holders Owning or orlgages, or Complete Mailing Address	
Holding 1 Percent of More of Total Amount of Bonde, M Diller Beautities, if none, check box Int Neme	Holdres Owning or stigsges, or <u>Vision (Complete Malling Address</u>	
Idaling I Persent of More of Total Amount of Bonde, Ma Dier Beanties, If none, check box Mill Nene	Holders Owning or stigages, or Complete Mailing Address	
klefig i Parcent of More of Total Amount of Bonds, M Beer Recurdles, If none, check box full Nene	Notdes coming or riguages, or Complete Malling Address	
Helding I Process of Monor of Food Amount of Bonds, Mo Media Beaufies, II none, check box M Nene	Notices Conving or Trases, or Complete Malting Address	
Heling I recent of More of Total Anount of Bonds, More Recently, If none, check box	Notdes Coming or Tratages, or Complete Matting Address	
Helling I Process of Noro of Total Amount of Bonds, Mo Media Bautiles, II none, check box Ma Nene San San San San San San San San San San	Notides Completes Malling Address Completes Malling Address Completes Malling Address address authorized to mail at special intensi (Check one)	
Helling I Process or Moor of Total Amount of Bonds, Mo Media Securities, II none, check box M Nene 2 Tex Bellue For completion by nonprofit agentization The partice, function, and nonprofit agentization Helling Outring Processing (2) Months	Notides Conving or Tripases, or Complete Malling Address Complete Malling Address authorized to mail at spacial rates) (Check one) suthorized to mail at spacial rates) (Check one) suthorized to mail at spacial rates) (Check one)	bruboset:
Net Status For completion by nonport operatorials Net Status For completion by nonport operatorials The Net Status For completion by nonport operatorials The perpose, function, and nonprofit status of the org- Man Alc Changed During Preceding 12 Monther Hea Changed During Preceding 12 Monther Hea Changed During Preceding 12 Monther	Notices conting or rightses, or Complete Malling Address Complete Malling Address Complete Malling Address	purposer:

13. Publication Title	14. Issue Date for Circulation Date Below	
Infection Control and Hospital Epidemiology	September 1999	
15. Extent and Nature of Circulation	Average No. Copies Each leave During Preceding 12 Months	Actual No. Copies of Single issue Published Nearest to Filing Date
a. Total Number of Copies (Net Print Run)	6,487	8,633
 b. Paid and/or Requested Circulation Sales Through Dealers and Carriers, Street Vendors, and Counter Sales (Not Malled) 	o	0
(2) Paid or Requested Mail Subscriptions (include advertiser's proof copies and exchange copies)	5,312	5,362
c. Total Paid and/or Requested Circulation (Sum of 15b(1) and 15c(2))	5,312	5,362
 Free Distribution by Mall (Samples, complimentary, and other free) 	433	442
e. Free Distribution Outside the Mail (Carriers or other means)	272	352
I. Total Free Distribution (Sum of 15d and 15e)	705	794
g. Total Distribution (Sum of 15c and 15f)	6,017	6,158
h. Copies not Distributed (1) Office Use, Lettovers, Spolled	470	477
(2) Returns from News Agents	0	0
I. Total (Sum of 15g, 15h(1), and 15h(2))	6,487	6,633
Percent Paid and/or Requested Circulation (15a/15g x 100)	88%	87%
16. Publication Statement of Ownership <u>x</u> Publication required. Will be printed in the <u>Novemb</u> Publication not required.	ar 1999 issue of this publication.	
17. Signature and Title of Editor, Publisher, Business Manager, or Owner		Date
Richard N. Rossh, VP/Group Publisher Richard N. Coash		October 1, 1999

Richard N. Rossh, VP/Group Publisher Richard N. Coash	October 1, 1999
I certify that all information furnished on this form is true and complete. I understand that anyone who furnishes	s false or misseading information on this form
or who omits material or information requested on the form may be subject to criminal sanctions (including fine	a and imprisonment) and/or civil sanctions
(including multiple damages and civil penalties).	