Regular Article

Shift-&-Persist and discrimination predicting depression across the life course: An accelerated longitudinal design using MIDUSI-III

N. Keita Christophe 💿 and Gabriela L. Stein

Department of Psychology, The University of North Carolina at Greensboro, Greensboro, NC, USA

Abstract

Life course theorists posit that sensitive periods exist during life span development where risk and protective factors may be particularly predictive of psychological outcomes relative to other periods in life. While there have been between-cohort studies trying to examine differences in discrimination and depressive symptoms, these studies have not been designed to identify these sensitive periods, which are best modeled by examining intra-individual change across time. To identify sensitive periods where discrimination and shift-&-persist (S&P) – a coping strategy that may protect against the negative impact of discrimination – are most strongly predictive of depressive symptoms, we employed latent growth curve modeling using an accelerated longitudinal design to track intra-individual change in depressive symptoms from ages 20–69. Participants were 3,685 adults measured at three time points ~10 years apart from the Midlife in the United States study ($M_{age} = 37.93$, SD = 6.948 at Wave I). Results identified two sensitive periods in development where high levels of S&P interacted with discrimination to protect against depressive symptoms; during the 30s and a lagged effect where 40's S&P protected against depressive symptoms for the life course study of discrimination, coping, and depression are discussed.

Keywords: depression, discrimination, life course, shift-&-persist, trajectories

(First published online 29 April 2021)

Discrimination, or the behaviors and unfair treatment reflecting negative attitudes and judgments about individuals in a group (Pascoe & Richman, 2009), has been consistently found through meta-analysis to be associated with poor mental health across the life span (Schmitt, Branscombe, Postmes, & Garcia, 2014; Yip, Wang, Mootoo, & Mirpuri, 2019). While the greatest amount of attention has been paid to ethnic-racial discrimination (Yip et al., 2019), where individuals are discriminated against as a function of their racial or ethnic group membership, discrimination has negative implications for psychological well-being regardless of attribution and across racial/ethnic groups, including White populations (Kessler, Mickelson, & Williams, 1999; Schmitt et al., 2014). Indeed, the inherently negative and uncontrollable nature of discrimination may make it particularly harmful to health and well-being (Pascoe & Richman, 2009; Williams & Mohammed, 2009). Shift-&-persist (S&P) is a coping strategy that has been shown to lead to positive health outcomes for vouth (Chen & Miller, 2012) and adults (Chen, Miller, Lachman, Gruenewald, & Seeman, 2012) who face uncontrollable stress. More recent work has also found S&P to be protective in the context of general discrimination (Lam et al., 2018) and

Author for Correspondence: N. Keita Christophe, Department of Psychology, University of North Carolina at Greensboro, 296 Eberhart Building, PO Box 26170, Greensboro, NC, 27402-6170; E-mail: nnchrist@uncg.edu

Cite this article: Christophe NK, Stein GL (2022). Shift-&-Persist and discrimination predicting depression across the life course: An accelerated longitudinal design using MIDUSI-III. Development and Psychopathology 34: 1544–1559. https://doi.org/10.1017/S0954579421000146

ethnic-racial discrimination (Christophe et al., 2019). To better understand how S&P may protect against negative mental health in the face of discrimination, we utilize the Midlife in the United States (MIDUS) dataset, and adopt a life course approach to examine how these factors impact the trajectory of depressive symptoms from age 20 to age 69.

Discrimination and Depression from a Life Course Perspective

Discrimination is a chronic and pernicious stressor that has been associated with greater concurrent and prospective depressive symptomatology (Luo, Xu, Granberg, & Wentworth, 2012; Schmitt et al., 2014; Williams, Neighbors, & Jackson, 2003). Despite scholarly work outlining discrimination's ability to concurrently and prospectively predict depressive symptoms, there is much less known from a life course perspective about whether there are particular periods in life course development where discrimination may operate as a particularly harmful stressor in predicting depressive symptoms and depression.

Life course theorists have proposed that the timing of exposure to discrimination may be particularly important in understanding its impact across the life span (Gee, Hing, Mohammed, Tabor, & Williams, 2019). To date, however, only a small number of studies are available that allude to possible changes in discrimination at different stages throughout the life span, possibly due to the scarcity of such wide-ranging data. For instance, using Wave I of MIDUS, Kessler and colleagues (1999) observed that, while overall reports of daily discrimination are lower in older cohorts

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (http:// creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

compared to younger cohorts, discrimination had an equal and negative impact on mental health across age cohorts. While this may, initially, imply is that the impact of discrimination on mental health is equivalent across age, this claim cannot be made, as Kessler and colleagues' study does not measure the intraindividual changes in associations between discrimination and depression across time. This study, and all studies focused solely on inter-individual change, cannot examine sensitive periods in individuals' lives where risks such as discrimination pose a particularly strong risk for their negative mental health. While not examining links to psychological functioning, Gee and colleagues' (2007) study of age discrimination among working women followed for 17 years provides a window into potential intraindividual change in exposure to discrimination. Specifically, the authors observed nonlinear trajectories of discrimination frequency, where discrimination starts high in the 20s, drops in the 30s, then increases and peaks when women were in their 50s (Gee, Pavalko, & Long, 2007). These meaningful changes in exposure to discrimination across decades may impact individuals' concurrent and future endorsement of depressive symptoms. In other words, as exposure to discrimination changes across the life course, its impact on concurrent and later psychological functioning may be dependent on the specific developmental stage in the life course.

The life stress literature provides a particularly useful perspective when thinking of sensitive periods where discrimination is more strongly associated with the occurrence or recurrence of depressive symptoms. In an attempt to highlight how stress exposure impacts the likelihood of depression as a function of age, Mazure and Maciejewski (2003) created a model, derived from data, illustrating that risk for depression due to life stressors increases in adulthood, peaks between ages 40 and 50, before declining to low levels of risk in the 60s and beyond. This would imply that discrimination may pose a particular risk for depression in midlife relative to early adulthood or old age. While the overall trajectory of depression risk due to life stressors does not fully map onto the trajectory of discrimination exposure outlined by Gee and colleagues (2007), it is important to note that frequency of exposure to discrimination may not be a driver of depressive symptoms as much as the impact of discrimination during different decades/developmental stages of life. For instance, the stress sensitization model of depression (see Monroe & Harkness, 2005 for review) asserts that, while initial onset of depression is more likely to be triggered by major life stressors, which typically first occur earlier in development, we become progressively more sensitized to stress and more chronic daily stressors - including discrimination - become more capable in inciting recurrences of depressive symptomatology than they did earlier in life. The authors contend that chronic daily life stressors, relative to major life stressors such as job loss and relationship dissolution, are in fact the modal stressors that precede increases in depressive symptoms later in adulthood.

Taken together, these studies provide evidence that midlife may be a particularly sensitive period where discrimination may, more strongly than is typical across the life course, predict greater levels of depression. Those in midlife are at a crossroads where their life stressors, both major and chronic, intersect leading to greater symptoms (Mazure & Maciejewski, 2003), and at the same time, they may also be more sensitized to stressors and more likely to become depressed in response to chronic, daily stressors, such as discrimination (Monroe & Harkness, 2005). In the current study, we attempt to test the claim that there is a sensitive period during midlife where discrimination is a particularly strong predictor of depression by employing an accelerated longitudinal design (Bollen & Curran, 2005) to assess intra-individual change across an almost 50-year time period using Waves I–III of the MIDUS study.

Shift-&-Persist

In addition to considering timing effects of discrimination, a life course perspective also posits that psychological resources may be differentially protective at specific sensitive periods in development (Gee et al., 2019). S&P coping may be such a resource that can mitigate the longitudinal negative impact of discrimination on depressive symptoms at specific points in development. Outlined by Chen and Miller (2012), S&P is a class of coping strategies that involve a combination of accepting and reappraising uncontrollable stressors (shift) coupled with maintaining optimism, purpose in life, and an orientation towards the future in spite of this uncontrollable stress exposure (persist). This coping strategy is theorized to be particularly well suited for uncontrollable stressors relative to direct attempts to eliminate or stop uncontrollable stressors (e.g., poverty, discrimination, etc.) and the combination of shifting and persisting in the context of uncontrollable stress that has been asserted (Chen, 2012) and empirically demonstrated (Chen et al., 2012) to be more effective than either strategy in isolation. In this sense, S&P coping may be conceptualized as what Lazarus and Folkman (1984) would refer to as a secondary-control coping strategy, or a strategy where one changes themselves to better fit the demands of the stressful environment. This is contrasted to primary-control coping strategies (Lazarus & Folkman, 1984), where individuals cope by actively altering the conditions of their environment. Attempts to actively and immediately "alter" or changes one's impoverished or discriminatory environment are likely to be unsuccessful, and these unsuccessful attempts to cope with uncontrollable stress may, themselves, elicit subsequent and exaggerated maladaptive stress responses (Chen, 2012).

When individuals come from disadvantaged backgrounds, S&P, has been associated with a number of positive physiological health outcomes. For instance, when low-socioeconomic status (SES) individuals utilize high levels of S&P coping, they have been shown to be protected against exhibiting greater body mass indexes (BMIs) (Kallem et al., 2013), experiencing greater asthma-related impairment (Chen et al., 2011), and demonstrating greater inflammation (Chen, McLean, & Miller, 2015). Particularly relevant to the current study, which draws from Waves I through III of the MIDUS study, Chen and colleagues (2012) conducted a study using a subset of 1,255 MIDUS II participants who provided biological samples to examine how S&P impacted allostatic load (i.e., physiological "wear and tear") for those from disadvantaged backgrounds. The authors observed that S&P protected against greater allostatic load for those from low-SES backgrounds. Adults who grew up in low-SES conditions but used high levels of S&P coping displayed the lowest levels of allostatic load (Chen et al., 2012). In addition, among youth who faced high levels of general discrimination, another form of uncontrollable stress, individuals who utilized S&P coping in the face of discrimination reported better asthma control and asthma quality of life compared to those who less frequently utilized S&P coping (Lam et al., 2018). Similar to what has been found with economic stress, the protective effects of S&P coping were not observed among those who faced little general discrimination.

Recent scholarship has attempted to extend the work on S&P to examine whether S&P coping may be similarly protective depressive symptoms, an important mental health outcome that is, itself, influenced in part by inflammatory processes (Slavich & Irwin, 2014). While originally conceptualized as being protective for physiological health (Chen & Miller, 2012), secondarycontrol coping strategies, which encompass strategies such as S&P, have been meta-analytically associated with fewer internalizing symptoms (Compas et al., 2017). Meaning and purpose in life, key components of persisting, have also been directly associated with fewer depressive symptoms (Sumner, Burrow, & Hill, 2018). Successful S&P coping with uncontrollable stressors may not only lead to adaptive downregulations in the stress response system (Chen, 2012), but may also reduce the impact that these stressors have on mental functioning. This conception of S&P as potentially adaptive for physiological and mental well-being stands, at face value, in contrast to recent work on "skin-deep resilience" (see Brody, Yu, Chen, & Miller, 2020 for brief summary of this work), which suggests that upwardly mobile individuals, especially those from minority groups, may demonstrate good mental health and poor physical health. However, the factors suggested to lead to a skin-deep resilience pattern, such as self-control (Brody et al., 2020; Miller, Yu, Chen, & Brody, 2015) and striving, which includes components such as high educational aspirations, an intense commitment to hard work, and a "single-minded" focus on success (Brody, Yu, Miller, & Chen, 2016), are not the same as factors characteristic of S&P coping. Stated differently, while striving and very high levels of selfcontrol have been associated with positive mental health but poor physiological health, it remains to be seen whether S&P coping in the context of uncontrollable stress is or is not protective of mental health in the same way that it has been shown to be protective of poor physical health.

Initial work examining the impact of S&P in the context of depressive symptoms has shown that S&P is protective against greater depressive symptoms depending on the nature of the uncontrollable stressor. For instance, in a sample of Latinx youth, Christophe et al. (2019) found that S&P coping was protective against the harmful effects of economic hardship on youth depressive symptoms. Further, when discrimination was examined as an uncontrollable stressor, discrimination was not associated with depressive symptoms when youth with relatively weak ethnic racial identities (i.e., meaning and importance of race/ethnicity; Umaña-Taylor et al., 2014) used high levels of S&P coping (Christophe et al., 2019). This protective effect of S&P against greater depressive symptoms has been found in an independent sample of 674 Mexican-origin youth (Stein et al., under review). Specifically, low ethnic-racial identity youth who faced high levels of discrimination endorsed relatively lower levels depressive symptoms in 9th grade and had marginally more steeply declining trajectories of symptoms throughout the high school years (Stein et al., under review). These two studies, thus, provide initial evidence that S&P coping may, contrary to striving (Brody et al., 2020), be protective against poor mental (Christophe et al., 2019) and physical health (Chen et al., 2015; Lam et al., 2018).

While these studies provide the basis for the notion that S&P might protect against discrimination, and specifically protect against depressive symptoms due to discrimination, this concept has not yet been tested in an adult sample, nor has it been tested across a large swath of the life span. Furthermore, while sensitive periods are most commonly explored for risk factors, Gee and colleagues (2019) assert that sensitive periods may also exist for

protective factors. Specifically, there may be times throughout development where protective factors such as S&P display particularly potent effects. Theoretically, these periods may overlap with those for stressors such as discrimination and also occur during midlife. If individuals are particularly sensitized to the effects of daily stressors in midlife (Monroe & Harkness, 2005), and most likely to experience depression at that time period due to those stressors (Mazure & Maciejewski, 2003), coping methods used during those times may be particularly efficacious. Therefore, in the current study we examine whether main effects of S&P coping and interactions between S&P and discrimination predict trajectories of depression across an almost 50-year window, as well as examining whether sensitive periods exist where S&P coping interacts with discrimination in predicting less concurrent and future depressive symptoms.

Across many of these studies, and in line with Chen and Miller's (2012) original theoretical assertion, S&P coping is often not directly associated with positive physiological outcomes as a main effect (Chen, Lee, Cavey, & Ho, 2013; Chen et al., 2015; Kallem et al., 2013; Lam et al., 2018), but instead moderates the effect of the uncontrollable stressor such that those who face greater levels of uncontrollable stress – be it economic stress or discrimination – are the ones who benefit from S&P coping. Individuals who do not face pervasive uncontrollable stressors likely do not benefit from employing coping strategies such as S&P coping that are especially suited to help people cope with uncontrollable stress (Chen, 2012; Chen & Miller, 2012); the effectiveness of S&P, thus, may be observable only for those facing frequent uncontrollable stress (i.e., a Stressor×S&P interaction).

Trajectories of Depressive Symptoms

While there exist a multitude of studies examining trajectories of depression from adolescence into emerging adulthood, far fewer studies have examined trajectories of depressive symptoms across and through the midlife period. Regardless of the population or trajectory modeling techniques employed, these studies point to meaningful, intra-individual change in depressive symptoms across time. For instance, employing multigroup growth curve modeling using the National Longitudinal Survey of Youth, Walsemann and colleagues (2009) found decreasing linear trajectories of depression from ages 27 to 43 among White, Black, and Latinx participants. By contrast, in a study utilizing Waves I-V of the National Longitudinal Study of Adolescent to Adult Health, Chen and Harris (2019) observed nonlinear trajectories of depression where unconditional depression scores, stratified by sex, tended to decrease throughout the 20s, and increase throughout the 30s until the end of the study window at age 40. In a study of 8,801 women studied across a 20-year period, Gillis et al. (2019) observed four different trajectories of depressive symptoms from ages 60 to 80 using latent growth curve mixture modeling: a small persistently high depression group (5%), and increasing depression group (14%), a decreasing trajectory group (19%), and a low and stable group that also appeared to be slightly decreasing over time (62%). It is important to note that 81% of this sample across the decreasing and low stable trajectories displayed at least some decline in symptomatology over time. In a study examining trajectories of negative affect in a large (N = 1992 at Wave I, 85% White), multigenerational sample of individuals, Charles, Reynolds, and Gatz (2001) observed a linear decrease in negative affect from ages 15 to 55 that then changed trajectory and began to decrease linearly at a slower rate from

age 60 through age 85. Finally, and most relevant to our current study that also uses the MIDUS sample, Blöchl and Nestler (2019) used second-order latent growth curve modeling on Waves I through III of the MIDUS data (see Isiordia & Ferrer, 2018 for explanation of this model) to examine how vascular risk factors (e.g., hypertension, diabetes, and smoking) impacted the linear trajectory of depression across the three study waves. Largely similar to past studies (Charles et al., 2001; Gillis et al., 2019; Walsemann, Gee, & Geronimus, 2009) the authors also found evidence for a slightly decreasing unconditional linear slope of depressive symptoms.¹

To summarize across these findings from varying samples and using varying modeling techniques, these studies point towards meaningful differences in the intra-individual trajectory of depression during and beyond the midlife period as a function of age. Furthermore, several of these studies (with Chen & Harris, 2019 as a notable exception) broadly point to an on-average decrease in depressive symptoms across the life course. Because intra-individual change across development has been observed for depressive symptoms (Charles et al., 2001; Chen & Harris, 2019; Gillis et al., 2019; Kessler, Foster, Webster, & House, 1992; Walsemann et al., 2009), it is crucial to employ modeling designs that account for developmental stage when examining both predictors of depression across the life course and sensitive periods during development when factors such as discrimination and S&P coping are particularly predictive of symptomatology relative to other developmental stages. This study builds upon these previous examinations of depressive trajectories by transforming the MIDUS study into an accelerated longitudinal design to both examine trajectory of depression across the life course as well as how exposure to discrimination and use of S&P coping influences depressive symptomatology.

Current Study

Based on life course theory, the current study attempted to conduct a developmentally informed examination of the trajectory of depression from ages 20 to 69 and examine how discrimination, S&P, and interactions between the two may concurrently and prospectively predict depressive symptoms during sensitive periods in development - specifically decades - controlling for participants' expected trajectory of depressive symptoms across the life course. While it is, theoretically, possible to examine the effect of discrimination and S&P on depressive symptoms from year to year, we assert that, when attempting to identify specific sensitive periods during development across the life course, using a decade-based approach to approximate each period in development is more appropriate. Given past studies finding differences in initial levels and the slope of depression from young adulthood through midlife (Walsemann et al., 2009), we also examine how initial levels of depression and the trajectory of depression may vary based on race and relevant covariates. Specifically, this study uses multigroup latent growth curve modeling with data from Waves I through III of the MIDUS study to estimate the trajectory of depression across each decade of this nearly 50-year timespan. We hypothesized that (a) depressive symptoms will show a linear decline throughout the life course and (b) controlling for covariates and the downward trajectory

of depressive symptoms, discrimination will concurrently predict greater than expected levels of depression during each decade of life. We also predicted, based on the work of Mazure and Maciejewski (2003) that (c) a sensitive period will exist during midlife (40s/50s) where discrimination will be an especially strong and predictor of concurrent and later depressive symptoms. Finally, we hypothesized that (d) while S&P will not be associated with depression as a main effect, it will interact with discrimination during midlife such that discrimination will not be associated with depressive symptoms during midlife for those using high levels of S&P coping.

Method

Participants & procedure

Participants were 3,685 adults ($M_{age} = 37.93$, SD = 6.948 range = 20-49 at Wave I) from the MIDUS dataset, a national study of health and well-being conducted by the MacArthur Foundation Research Network on Successful Midlife Development where participants were first selected through random-digit phone dialing (see Radler, 2014 for brief description). The participants in this study represent a subsample of MIDUS participants from Project 1, the original project launched in 1995 assessing psychosocial and health-related outcomes, who were not missing on important demographic information such as race and gender. Participants included those from the national probability sample (N=1716), the sibling sample (N=456), the twin sample (N=1143), and the city oversample (N = 370). Our sample was 89.2% monoracial White, with 5.7% of participants identifying as Black/African American and 2.2% identifying as Hispanic/ Latina(o). Given the skewed race/ethnicity distribution, sampling weights were used to increase the influence of responses from racially minoritized participants, thus improving generalizability and partially correcting for the increased attrition observed among racially minoritized participants (see "Sample weights for race" section for greater detail). The sample was fairly evenly split with regards to biological sex (51.6% female) (only binary sex collected in study). A small proportion of our sample (5.3%) was born outside of the United States. At Wave 1, 25.6% of the sample had obtained a high school diploma, an additional 22.2% had obtained at least a bachelor's degree, and an additional 10.7% had obtained a masters of other graduate degree. (e.g., MD, JD, PhD, etc.).

Wave I was collected during 1995 and 1996, while Wave II was collected about 10 years later in 2004 and 2005. Wave III was collected during 2013 and 2014. Missingness was 25.3% at Wave II of data collection (N = 2751 present), and 46.4% at Wave III (N = 1974 present). To facilitate our comparison of trajectories of depression across development, participants were separated into three separate birth cohorts based on age at Wave I of data collection. Participants from Cohort 1 were born between 1965 and 1974 and were thus between the ages of 20 and 29 at Wave I in 1995. Participants in Cohort 2 were between 30 and 39 (birth years 1955–1966) at Wave I and Cohort 3 was between 40 and 49 (birth years 1945-1956). This study focused on understanding depression risk and resilience during sensitive developmental periods in adulthood, versus the impact of discrimination and S&P coping on depression in advanced ages (70-90+ years old), where significant physical declines, cognitive declines, and declines of social connectedness may disrupt one's previously observed trajectory of depressive symptoms. Given

¹This study solely examines change over time, not change over time based on an individual's age as do the previously cited studies. Further explanation of this is provided in the Discussion section.

our focus on the former question, individuals above age 49 at Wave I were excluded from our sample, meaning that the most elderly participants in our sample were between ages 60 and 69 at Wave III of data collection. Additional demographic information for the whole sample and each cohort may be seen in Table 1.

Measures

Depressive symptoms

Depressive symptoms were measured at each time point using the five items from the Kessler-6 (Kessler et al., 2002). Participants were asked how often they felt, for example, "hopeless", "worthless", and "that everything was an effort" in the last 30 days for 1 (all of the time) to 5 (none of the time). One item asking participants how often they felt "nervous" was dropped based on a confirmatory factor analysis with the full MIDUS I - MIDUS III data that indicated that adequate fit of a unidimensional model for depressive symptoms were obtained only when this item was dropped (Blöchl & Nestler, 2019). The remaining five items were reverse-coded such that greater scores indicated more depressive symptoms. The full Kessler-6 has displayed reliability of .89 in the nationally representative telephone-based pilot survey for MIDUS (Kessler et al., 2002). At Waves I, II, and III of our sample, reliability was .857, .853, and .849, respectively, for the five-item measure. Test-retest reliability between Waves 1 and 2 was .479, and test-retest reliability between Waves 2 and 3 was .567.

Daily discrimination

Daily discrimination was assessed at each time point using the nine-item daily scale (Williams, Yan, Jackson, & Anderson, 1997). Participants were asked to respond to items such as how often "people ask as if you are dishonest" and "you are treated with less respect than other people" on a 1 (*often*) to 4 (*never*) scale. Items were reverse-coded and summed together such that higher scores indicated more frequent daily discrimination. This measure has displayed a reliability of .88 in a diverse sample of 1,139 participants from the Detroit Area Study (Williams et al., 1997). Reliability in our sample at Wave 1, 2, and 3 was .855, .918, and .916, respectively. Test-retest reliability between Waves I and II was .52, and test-retest reliability between Waves II and III was .569.

Participants who indicated that they have faced some form of daily discrimination were then asked to select the main reason they were subjected to discrimination. Response options available across all three waves of data were: age, gender, race, ethnicity/ nationality, religion, height/weight, physical appearance, physical disability. Participants were allowed to select multiple main reasons for facing daily discrimination. Asking about the prevalence of discrimination before asking about the attribution has been theorized to help reduce underestimation of discrimination – a phenomenon that can occur when victims struggle to clearly identify the reason for which they were discriminated against (Williams et al., 2003).

Shift-&-Persist

Shift was measured at each of the three time points using the fouritem positive reappraisal subscale of the Primary and Secondary Control Questionnaire (Wrosch, Heckhausen, & Lachman, 2000). Participants responded to sample items such as "I can find something positive, even in the worst situations" and "when I am faced with a bad situation, it helps to find a different way of looking at things" on a 4-point scale from 1 (*A lot*) to 4 (*Not at all*). Items were reverse-coded and averaged such that greater values indicated greater shifting, or positive reappraisal. Reliability in our sample at Wave I, II, and III was .796, .801, and .788, respectively. Test-retest reliability between Waves I and II was .562, and test-retest reliability between Waves II and III was .646.

Persist was measured at each of the three time points using the four-item "live for today" subscale of the "Planning and Making Sense of the Past Questionnaire" (Prenda & Lachman, 2001). Participants responded to sample items such as "I have too many things to think about today to think about tomorrow" and "I live one day at a time" on a 4-point scale from 1 (*a lot*) to 4 (*not at all*). Items were reverse-coded and averaged, where greater scores indicated a higher propensity to *persist* through life. Reliability in our sample at Wave I, II, and III was .641, .637, and .609, respectively. Test–retest reliability between Waves I and II was .521.

Consistent with past work with the S&P construct (Chen et al., 2015; Kallem et al., 2013), and the theoretical assertion that S&P exerts a joint influence greater than the sum of its component parts (Chen, 2012; Lam et al., 2018), we computed a single "S&P" score by creating separate mean scores for shift and for persist, standardizing these mean scores, then averaging these standardized scores together. Higher scores on this composite variable indicate greater use of S&P coping.² Test–retest reliability between Waves I and II was .530, and test–retest reliability between Waves II and IIIwas .619.

Covariates

Covariates in our study include biological sex, nativity status, race, education as a measure of SES (higher values = greater education/ SES), and neuroticism. Sample was also added through three dummy-coded variables for sibling sample, twin sample, and city oversample, with the main probability sample as the reference group. Due to low Ns for non-White participants, race was dichotomized such that those who identify as monoracial white were coded "0", while those who are multiracial or of other racial minority groups were coded as "1". Similar to our measures of discrimination and S&P, SES was entered as a time varying covariate, as education/SES may change across time. Neuroticism, which has been proposed as a factor that may influence relations between stressors such as discrimination, and psychological adjustment (Stokes, 2019), was measured at all three time points using a four-item neuroticism subscale generated from previous "Big-Five" inventories for the MIDUS study (Keyes, Shmotkin, & Ryff, 2002; Rossi, 2001). On a 1 (a lot) to 4 (not at all) scale, participants rated how well the words, moody, worrying, nervous, and calm described them. The moody, worrying, and nervous items were reverse coded before creating a mean score, where higher numbers indicate greater trait neuroticism. Reliability was .747 in our sample. Finally, dummy-coded variables indicating the sample a participant was drawn from (i.e., city oversample,

²This measure of S&P differs slightly from that used by Chen et al.'s (2012) study examining S&P during Wave II of MIDUS. To assess shift, Chen et al. (2012) use the positive reappraisals subscale and the emotional reactivity subscale from the Multidimensional Personality Questionnaire (Patrick, Curtin, & Tellegen, 2002). Because the emotional reactivity subscale is not available at all three waves of data, we, unlike Chen et al., do not use it as part of our measure of shift. For persist, we use the same measure as Chen et al. (2012), the live for today subscale (Prenda & Lachman, 2001).

Table 1	L.	Sample	demograph	nics f	for	final	analytical	sample
---------	----	--------	-----------	--------	-----	-------	------------	--------

Variable	Total sample (N = 3685)	Cohort 1 (<i>N</i> = 538)	Cohort 2 (<i>N</i> = 1518)	Cohort 3 (<i>N</i> = 1629)
Age at Wave I (SD)	37.93 (6.948)	26.98 (1.476)	34.72 (2.806)	44.55 (2.849)
Birth years	1945–1974	1965–1974	1955–1966	1945–1956
Biological sex				
Male	48.4%	47.2%	47.5%	49.5%
Female	51.6%	52.8%	52.5%	50.5%
Race				
White	89.2%	84.0%	88.1%	91.9%
Black/African American	5.7%	7.4%	6.1%	4.7%
Native American/Aleutian Islander/ Eskimo	.7%	.9%	.6%	.6%
Asian/ Pacific Islander	1.2%	2.6%	1.6%	.4%
Multiracial	.7%	1.3%	.5%	.6%
Other	2.6%	3.3%	3.0%	1.7%
Latinx origin*				
Yes	2.2%	2.1%	2.7%	2.0%
No	72.2%	66.4%	70.6%	75.6%
Missing	25.6%	31.6%	26.8%	22.4%
Born in United States				
Yes	94.5%	93.7%	93.9%	95.5%
No	5.3%	5.9%	5.9%	4.5%
Missing	.2%	.4%	.3%	.1%
Parental Education				
Below high school	15.1%	6.7%	12.3%	18.5%
Above high school	80.1%	89.0%	82.4%	75.0%
Missing	5.6%	4.3%	5.3%	6.4%
Sample				
Main	46.6% (1716)	52.4% (282)	45.7% (694)	45.4% (740)
Sibling	12.4% (456)	6.9% (37)	11.7% (178)	14.8% (241)
Twin	31.0% (1143)	27.7% (149)	32.1% (487)	31.1% (507)
City oversample	10.0% (370)	13.0% (70)	10.5% (159)	8.7% (141)

Note. Questions asking about Latinx/Hispanic origin were first asked at Wave II.

sibling sample, or twin sample) were entered as covariates, with the national probability sample serving as the reference group. groups being in our MIDUS I sample and partially corrects for the observed differential attrition by race in MIDUS II and III.

Sample weights for race

Because racial minority groups were underrepresented in our sample relative to the population, post-stratification sampling weights were used adjusting the weight of responses by race. Using the same convention as is done with the full MIDUS sample, weights for our subsample were created by comparing the percentages of White, Black, and "Other" (i.e., non-Black minority and multiracial participants) of our sample against those from the 2013 Current Population survey from the US Census Bureau. Dividing the demographic percentages in the Current Population Survey by the percentages in our sample produced weights of .90 for Whites, 2.76 for Blacks, and 2.13 for "Others". Using sampling weights partially corrects for the underrepresentation of minority

Results

Descriptive statistics

Before running our multigroup latent growth curve model, we calculated descriptive statistics for the whole sample and each sample separately (see Table 2). To summarize, depressive symptoms were positively associated with discrimination at each time point, and discrimination was also positively associated with S&P across time points. Neuroticism was also moderately associated with depressive symptoms at every time point ($rs \ge .328$), whereas minority status and immigrant status were only weakly associated with depressive symptoms (rs .037-.136). Across cohorts and waves of data, gender was the most commonly

 Table 2. Means and correlations among key study variables

	Variable	Cohort 1	Cohort 2	Cohort 3	Full Sample	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
(1)	Dep 1	1.616 (.675)	1.526 (.640)	1.511 (.645)	1.534 (.649)	1								
(2)	Dep 2	1.612 (.766)	1.538 (.677)	1.502 (.609)	1.531 (.660)	.456	1							
(3)	Dep 3	1.561 (.701)	1.459 (.610)	1.428 (.619)	1.455 (.627)	.397	.599	1						
(4)	Disc 1	14.981 (5.634)	14.442 (5.543)	13.657 (5.268)	14.192 (5.465)	.200	.197	.184	1					
(5)	Disc 2	14.383 (5.397)	13.946 (5.001)	13.493 (4.977)	13.787 (5.051)	.199	.315	.287	.606	1				
(6)	Disc 3	13.765 (5.266)	13.420 (5.085)	12.695 (4.579)	13.096 (4.878)	.153	.219	.275	.590	.641	1			
(7)	S&P 1	012 (.731)	.003 (.702)	.066 (.725)	.028 (.718)	072	.022	.041	.066	.077	.107	1		
(8)	S&P 2	.022 (.646)	.009 (.717)	.052 (.727)	.031 (.713)	.021	.050	.022	.092	.131	.107	.545	1	
(9)	S&P 3	.037 (.694)	031 (.704)	.038 (.705)	.011 (.704)	.039	.037	.019	.081	.080	.116	.510	.628	1

Note. Correlation are for the whole sample. Correlations significant at p < .05 are in **bold**. Dep = depressive symptoms. Disc = discrimination. S&P = shift-&-persist. The number after variable name corresponds to the Wave of data.

cited reason for discrimination (36.2%), followed by race (20.4%) and height/weight (19.6%). Discrimination based on physical disability (2.8%) was the most uncommon perceived reason for discrimination.

Data analytic plan

After running descriptives on observed study variables, inferential data analysis advanced in four steps. In the first step we estimated a multigroup latent intercept model and compared its fit relative to an unconditional linear growth model, where depressive symptoms were entered as indicators of the latent intercept and latent linear slope factors. Time was rescaled such that the intercept for the latent curve model corresponded to participants' 20s (ages 20-29). Scaling of the slope factor for each cohort therefore corresponded to the deviation in decades from participants' 20s (see Table 2 for visual representation). This allows the loadings for the slope factor to be identical at overlapping ages and for us to thus model growth in depressive symptoms from age 20 through age 69 (Mehta & West, 2000). Second, we tested whether homoscedasticity, or constraining the residual variances in our depressive symptom factors across time to equality, fit as well as a latent intercept and slope model where residual variances were freely estimated. This allowed us to test whether the impact of depressive symptoms on the intercept and slope varied across time.

In Step 3, we added time-invariant covariates (TICs), namely race, biological sex, immigrant status, sample, neuroticism, and parental education predicting the intercept and linear growth factors of depressive symptoms. We also added our main variables of interest, daily discrimination and S&P, as well as their product term, to the model as time-varying covariates (TVCs). In this model, which included both TICs and TVCs, TVCs predicted depressive symptoms at each time point controlling for the impact of the latent intercept and slope factor, while TICs predicted the growth factor of depressive symptoms over and above the effects of the TVCs (Bollen & Curran, 2005). We tested for contemporaneous and lagged effects of our TVCs (e.g., Wave I discrimination predicting Wave IIdepression) given theoretical work proposing that sensitive periods may exist throughout development where exposure to discrimination at that time is particularly harmful for later health and well-being (Gee et al., 2019). These sensitive periods are also theorized to exist for protective factors. Finally, in Step 4, we compare this conditional model to a model where the covariance between the latent intercept and slope is constrained to equality across our three cohorts of participants. This tests whether initial depression in the 20s is equally related to the slope of depression across all three cohorts. Throughout these steps, model fit was determined based on a combination of indices, where a nonsignificant chi-square, a comparative fit index $(CFI) \ge .95$, an root mean square error of approximation $(RMSEA) \leq .05$, and standardized root mean square residual $(SRMR) \le .08$ indicate "good" fit (Hu & Bentler, 1999). When appropriate, competing models are compared using the Satorra-Bentler scaled chi-square likelihood ratio test (Satorra, 2000). Model parameters were not interpreted in depth until we had decided upon our well-fitting, final model. Because we used sampling weights, model parameters were estimated using a maximum likelihood estimator that was robust to nonnormality (MLR). Because of our accelerated longitudinal design, some data were missing by design³ and, thus, intentionally not handled using missing data procedures. Missing data that were not missing by design were estimated using full information maximum

Competing latent curve models

We first fit an intercept-only model using summary scores for depression as indicators of the latent intercept (Model 1). This model provided an adequate fit to the data but did not fit better than a subsequent unconditional model (see Table 3 for model fit statistics from competing models) including a random intercept and slope factor (Scaled ΔX^2 (1) = 31.007, *p* < .0001) where residual variances for depression were freely estimated (i.e., heteroscedastic). This unconditional heteroscedastic model (Model 2) provided a good fit to the data $(X^2 (11) = 17.359, p = .0077,$ RMSEA = .022, CFI = .975, SRMR = .045). There was a significant intercept (M = 1.575, p < .001) and negative slope (M = -.026, p = .001), indicating that depressive symptoms across the sample is expected to go down by .026 points per ~10-year period from ages 20 to 69. The intercept was not associated with the slope for any cohort, meaning that initial levels of depressive symptoms during the 20s was not associated with the downward trajectory of depressive symptoms across time. In this unconditional model, there was significant individual variability around the intercept of depressive symptoms (.200, p = .007) but not the slope (.021, p = .118). The unconditional heteroscedastic model was then tested against a homoscedastic model (Model 3), where residual variances for the depression factors were constrained to equality. The more parsimonious homoscedastic model provided a significantly worse fit to the data (Scaled ΔX^2 (8) = 134.782, p < .0001). Therefore, a heteroscedastic structure was retained and used in subsequent model evaluation.

Next, we added TICs (biological sex, immigrant status, parental education, neuroticism, and race) and TVCs (discrimination, S&P, and their interactions at each time point) to our model (Model 4). Contemporaneous and lagged effects (e.g., discrimination in 20s predicting depressive symptoms in one decade later) were assessed for our TVCs. Again, at points where our three cohorts are assessed at overlapping ages (see Supplementary Table 1), parameters were constrained to equality to facilitate the study of quasi-intra-individual change through our accelerated longitudinal design. Sex, immigrant status, and our dummycoded MIDUS sample variables (i.e., sibling, twin, or city oversample - national probability sample is reference group) were not associated with the slope and intercept of depressive symptoms and were thus dropped for greater model parsimony. This trimmed model provided a good fit to the data $(X^2 (121) =$ 185.896, *p* = .0001, RMSEA = .021, CFI = .958, SRMR = .031). Given nonsignificant covariances between the intercept and slope between each of the three cohorts, we tested whether a more parsimonious model where these covariances were set to equality would fit the data equally well. This model (Model 5) provided an adequate fit to the data $(X^2 (123) = 214.059, p <$.0001, RMSEA = .025, CFI = .940, SRMR = .035), but fit the data significantly worse than the Model 4, where these covariances were freely estimated by cohort (Scaled ΔX^2 (2) = 19.020, p < .0001). Given the good model fit and good fit relative to an

 $^{^{3}}$ By creating cohorts to facilitate our accelerated longitudinal design, we have introduced missingness by design. For example, Cohort 2, who were between ages 30 and 39 during Wave I and between 50 and 59 at Wave III, do not have data during the 20s or 60s by design (see Supplemental Table 1 for visual depiction for which decades cohorts are missing data by design).

Model	X^2 (df)	p	RMSEA	CFI	TLI	SRMR	MLR scaling factor	Satorra– Bentler Scaled X ² (df)	p	Decision
Unconditional models										
(1) Intercept Only	28.306 (12)	.0050	.033	.936	.952	.072	2.9159	-	-	-
(2) Unconditional heteroscedastic	17.359 (11)	.0977	.022	.975	.980	.045	2.2510	31.007 (1)	<.0001	Reject model 1 Retain model 2
(3) Unconditional homoscedastic	46.364 (19)	.0004	.034	.893	.949	.088	3.1041	134.782 (8)	<.0001	Reject model 3 Retain model 2
Conditional models										
*(4) Conditional model with TICs & TVCs	185.896 (121)	.0001	.021	.958	.953	.031	1.3901	-	-	-
(5) Conditional model with TICs & TVCs - intercept & slope covariance set to equality	214.059 (123)	<.0001	.025	.940	.935	.035	1.4034	19.020 (2)	<.0001	Reject model 5 Retain model 4

Table 3. Model fit statistics from principled evaluation of competing curve of factors models

Note. * = final analytical model. Maximum likelihood robust (MLR) scaling factor is needed in the calculation of the Satorra-Bentler Scaled χ^2 difference test (Satorra, 2000). Depressive symptoms are the dependent variable for all models. Unconditional models do not contain any covariates, while conditional models include the following covariates: socioeconomic status (SES) (time-varying), neuroticism (time-invariant), and race (time-invariant).

CFI = comparative fit index; RMSEA = root mean square error of approximation; SRMR = standardized root mean square residual; TLI = Tucker-Lewis Index.

alternative model, we used Model 4, our conditional model with TICs and TVCs where intercept slope covariance is freely estimated across cohort, as our final analytical model.

Interpreting final analytical model

In our final conditional analytical model (see Table 4), there continued to be a significant intercept (intercept = 1.578, p < .001), and a significant downward linear slope (slope = -.026, p < .001) of depressive symptoms. Again, the intercept is interpreted for all cohorts as the level of depression when participants were in their 20s. There was significant residual variation in the value of depressive symptoms in the 20s (residual variance = .107, p = .030) and the slope of depression across time (residual variance = .023, p = .018) that was not accounted for by our covariates.

Overall, the time-invariant and time-varying predictors in the model explained a significant amount of variance in the intercept $(R^2s = .447 - .633, ps < .001)$ but not in the linear slope $(R^{23}s = .079)$ -.136, ps = .062 - .186). The intercept of depression was not associated with the slope for any of the three cohorts. Greater neuroticism predicted a higher latent intercept of depression in the 20s (B = .446, p < .001) and a more steeply decreasing trajectory of depressive symptoms across the life course (B = -.060, p = .001), again controlling for our TVCs. Moving to our TVCs, whose levels and effects may vary by decade of life, we first examined the effects of SES (measured by education level) on depressive symptoms. During each decade of life, greater SES was associated with fewer same-decade depressive symptoms. Controlling for the negative slope of depression and all TICs, there were no contemporaneous nor lagged effects of S&P on depressive symptoms during any decade of life. During all decades of life, discrimination was concurrently associated with greater depressive symptoms controlling for the TICs and the negative linear slope of depressive symptoms across time (see Figure 1 for visual representation). There was one significant lagged effect where discrimination experienced in participants' 40s led to greater depression 10 years later when participants were in their 50s (B = .010, p = .019), implying a sensitive period where discrimination during the 40s was particularly harmful in predicting later depression. This lagged effect occurred over and above the significant contemporaneous effect of discrimination in the 50s.

Examining interactions between S&P and discrimination, when participants were in their 30s, S&P and discrimination significantly interacted with each other and predicted less concurrent depression than would be expected given the negative trajectory of depression (B = -.011, p = .015). Specifically, at mean levels of S&P, discrimination predicted greater concurrent depression when participants were in their 30s (B = .019, p < .001); this effect was intensified for participants who endorsed low S&P coping in their 30s (B = .027, p < .001), but was attenuated for those who were high in S&P coping at this time (B = .011, p = .018). This means that, when participants were in their 30s, high levels of S&P coping partially buffered against the concurrent association between discrimination and depressive symptoms (see Figure 2).

In addition, we found evidence of a lagged interactive effect where the interaction between S&P and discrimination during the 40s was predictive of depression during the 50s than what would be expected given the negative slope of depression (B = -.012, p = .022). Probing this interaction (see Figure 3) revealed that, while discrimination when participants were in their 40s predicted greater depression 10 years later if participants endorsed low (B = .019, p < .001) and mean (B = .010, p = .019)S&P coping at that earlier time, this association did not hold for those who had displayed high S&P coping while in their 40s

Table 4. Standardized parameter estimates for final analytical model (model 4)

Path	Cohort 1 (N = 538)	р	Cohort 2 (N = 1518)	р	Cohort 3 (N = 1629)	р
Intercept	1.578	<.001	1.578	<.001	1.578	<.001
Slope	026	<.001	026	<.001	026	<.001
R ² Intercept	.469	<.001	.447	<.001	.633	<.001
R ² Slope	.079	.186	.070	.180	.136	.062
Intercept ↔ Slope	.007	.792	012	.542	033	.120
Effects of time-invariant covariates						
Neuroticism → Intercept	.446 (.030)	<.001	.446 (.030)	<.001	.446 (.030)	<.001
Neuroticism → Slope	060 (.018)	.001	060 (.018)	.001	060 (.018)	.001
Race (1 = non-White) \rightarrow Intercept	.073 (.054)	.174	.073 (.054)	.174	.073 (.054)	.174
Race \rightarrow Slope	039 (.039)	.317	039 (.039)	.317	039 (.039)	.317
Concurrent effects by age						
SES						
20's SES	031 (.013)	.019	-	-	-	-
30's SES	022 (.007)	.001	022 (.007)	.001	-	-
40's SES	011 (.005)	.042	011 (.005)	.042	011 (.005)	.042
50's SES	-	-	019 (.006)	.001	019 (.006)	.001
60's SES	-	-	-	-	035 (.009)	<.001
S&P						
20's Depression	102 (.061)	.094	-	-	-	-
30's Depression	029 (.022)	.200	029 (.022)	.200	-	-
40's Depression	010 (.019)	.603	010 (.019)	.603	010 (.019)	.603
50's Depression	-	-	013 (.025)	.592	013 (.025)	.592
60's Depression	-	-	-	-	032 (.038)	.399
Discrimination						
20's Depression	.016 (.007)	.023	-	-	-	-
30's Depression	.019 (.004)	<.001	.019 (.004)	<.001	-	-
40's Depression	.021 (.003)	<.001	.021 (.003)	<.001	.021 (.003)	<.001
50's Depression	-	-	.020 (.004)	<.001	.020 (.004)	<.001
60's Depression	-	-	-	-	.020 (.010)	.044
S&P×Discrimination interaction						
20's Depression	031 (.016)	.055	-	-	-	-
30's Depression	011 (.005)	.015	011 (.005)	.015	-	-
40's Depression	.001 (.004)	.886	.001 (.004)	.886	.001 (.004)	.886
50's Depression	-	-	.002 (.006)	.756	.002 (.006)	.756
60's Depression	-	-	-	-	.004 (.009)	.649
10-year lagged effects						
S&P						
20's S&P \rightarrow 30's Depression	.152 (.105)	.149	-	-	-	-
30's S&P \rightarrow 40's Depression	.014 (.031)	.644	.014 (.031)	.644	-	-
40's S&P \rightarrow 50's Depression	-	-	003 (.025)	.901	003 (.025)	.901
50's S&P \rightarrow 60's Depression	-	-	-	-	021 (.039)	.581
Discrimination						
20's Discrimination \rightarrow 30's Depression	.009 (.013)	.500	-	-	-	-

Table 4. (Continued.)

Path	Cohort 1 (N = 538)	р	Cohort 2 (N = 1518)	р	Cohort 3 (N = 1629)	р
	· · ·	•			· · ·	•
30's Discrimination \rightarrow 40's Depression	.006 (.004)	.186	.006 (.004)	.186	-	-
40's Discrimination \rightarrow 50's Depression	-	-	.010 (.004)	.019	.010 (.004)	.019
50's Discrimination \rightarrow 60's Depression	-	-	-	-	.014 (.009)	.104
S&P × Discrimination interaction						
20's interaction \rightarrow 30's Depression	035 (.019)	.063	-	-	-	-
30's Interaction \rightarrow 40's Depression	009 (.007)	.161	009 (.007)	.161	-	-
40's Interaction \rightarrow 50's Depression	-	-	012 (.005)	.022	012 (.005)	.022
50's Interaction \rightarrow 60's Depression	-	-	-	-	.005 (.011)	.617

Note. Significant values are bolded for increased readability. - = estimate is missing by design. At points where the same parameter is assessed across multiple cohorts, that parameter has been constrained to equality across those cohorts. \leftrightarrow indicates a correlation. \rightarrow indicates a regression path.





Figure 1. Graph of depressive trajectory across the life course and the same-decade impact of discrimination on depressive symptoms. *Note.* The symptom trajectory value should be interpreted as the level of depressive symptoms predicted based on one's symptom trajectory across time. The effect of same decade discrimination value should be interpreted as the predicted level of depressive symptoms based on one's exposure to discrimination during that decade of life. The difference between these values is equal to the significant (p < .05) beta weight for discrimination. The gray bar indicates the predicted level of G&P coping.



Figure 2. Simple slopes plot of the Discrimination×Shift-&-Persist (S&P) interaction predicting same-decade depressive symptoms in the 30s.

(B = .001, p = .902). This means that while discrimination was associated with concurrent greater depression in the 40s regardless of S&P, high levels of S&P prevented discrimination from having an impact on depression 10 years later; this fully protective

Figure 3. Simple slopes plot of the 10-year lagged interaction effect predicting depressive symptoms in the fifties.

effect may only be observed during this specific sensitive period of development.

Discussion

This study took a life course perspective to examine how exposure to discrimination and S&P coping impacted depressive trajectories from ages 20 to 69. The accelerated longitudinal design improves our ability both to track the trajectory of depressive symptoms throughout a large swath of the life span by modeling intra-individual change over time while also testing whether sensitive periods in development exist where discrimination, S&P, and their interactions are especially strong predictors of concurrent and prospective deviations from individuals expected trajectory of depressive symptoms. Broadly, this study aligns with past work demonstrating change across the life span in the expression of depressive symptoms (Charles et al., 2001; Gillis et al., 2019; Walsemann et al., 2009) and affirms that discrimination is a significant predictor of depressive symptoms throughout much of the life span (Schmitt et al., 2014). Furthermore, our findings expand what is known regarding the efficacy of S&P coping beyond the realm of physiological health outcomes by contributing to a small but growing body of literature demonstrating the protective effects of S&P coping for depressive symptoms. Specifically, our findings provide additional evidence that S&P coping may protect

individuals against depressive symptomatology during certain sensitive periods in life course development.

We first found evidence for a significant downward trajectory of depression from ages 20 to 69. This finding builds off of previous work by Blöchl and Nestler (2019) examining the trajectory of depression in the MIDUS sample. By transforming the MIDUS data into an accelerated longitudinal design, we, contrary to Blöchl and Nestler (2019), were not only able to model trajectories of depressive symptoms as a function of time (i.e., the 20 years that pass between Wave I and Wave III), but also as a function of development (i.e., estimating intra-individual change from ages 20 to 69). This emphasis on intra-individual change is especially needed given previously documented within-person changes in symptomatology across the life course (Charles et al., 2001; Chen & Harris, 2019; Gillis et al., 2019; Kessler et al., 1992; Walsemann et al., 2009). Our findings coincide with much of this previous work and suggest that, on average, individuals endorse fewer depressive symptoms as they age up until age 69.

Examining predictors of slope of depression, neuroticism emerged as a significant predictor. Across the life course, neuroticism has been strongly associated with greater internalizing symptoms (Tackett & Lahey, 2016) and trajectories of negative affect (Charles et al., 2001). This corresponds with our finding that neuroticism in the 20s is related to initial depressive symptoms in the 20s and the slope of depressive symptoms through the 60s. Given that 20's neuroticism predicted a steeper decline in depressive symptoms over time, our finding may represent a "regression to the mean" where individuals who are highly neurotic in their 20s endorse fewer depressive symptoms across development – effectively placing these individuals back on the modal downward trajectory of depressive symptoms.

In terms of our time-varying effects, discrimination, largely consistent with Hypothesis 1, predicted greater concurrent levels of depression than would be expected based on the downward trajectory of depressive symptoms. The findings that discrimination predicts greater depressive symptoms at each decade of development is unsurprising and aligns with a large body of work (see Paradies et al., 2015; Pascoe & Richman, 2009; Schmitt et al., 2014; Williams & Mohammed, 2009 for reviews and metaanalyses) research attesting to cross sectional positive associations between discrimination and depression. In addition, we did not observe any significant main effects of S&P on depressive symptoms. While this may cast doubt on the efficacy of S&P as a coping strategy, these findings also align with previous work finding interactive, but not main effects of S&P (Chen et al., 2015; Christophe et al., 2019; Kallem et al., 2013). Indeed, S&P is conceptualized as a set of coping strategies that is particularly effective in the context of uncontrollable stressors (Chen & Miller, 2012). In the absence of uncontrollable stressors such as low-SES or discrimination, S&P is not theorized and has not been shown to be associated either with improved physical (Chen et al., 2012; Lam et al., 2018) or mental health (Stein et al., under review; Christophe et al., 2019). "Goodness of fit" is an important factor to consider when examining coping; other active coping strategies that are aimed at altering the environment itself may be more suited to coping with controllable stressors, while changing oneself to the demands of the environment (e.g., shifting and persisting) may be more adaptive when stressors are uncontrollable (Lazarus & Folkman, 1984; Stanisławski, 2019). Therefore, theory would assert that S&P should be principally adaptive in the context of high levels of uncontrollable stress, which is what we found

when examining time-specific interactions between S&P and discrimination.

In terms of context-specific effects, which imply especially sensitive decades in development where the effects risk and protective factors may be particularly pronounced, we found evidence for Hypothesis 2 in the form of two time-specific interactions between S&P and discrimination. The first was a concurrent interactive effect observed during the 30s. Specifically, the positive relation between discrimination and depressive symptoms in the 30s were weakened at high levels of S&P coping. We also observed a lagged interactive effect where discrimination when participants were in their 40s predicted greater depression than would be expected in the 50s. This lagged effect, however, was attenuated by S&P coping during the 40s and nonsignificant when S&P was at high levels. This lends credence to the hypothesis proposed by Gee and colleagues (2019) that sensitive periods exist both for risk factors (i.e., discrimination), but also for protective factors (i.e., S&P).

Relative to other points along the life course, there may be something unique about facing discrimination and using S&P coping in the 30s and especially in the 40s where these two factors interact to predict depressive symptoms a full 10 years later. Historically, psychologists such G. Stanley Hall conceptualized midlife as spanning from 25 or 30 to 40 or 45 years of age and described this as when we are, generally, at the peak of our abilities (1922). While other theorists commonly place "midlife" slightly later in the life cycle, Lachman and colleagues (2015) highlight that Jung and Erikson also saw midlife as a critical time where individuals are "negotiating and regulating growth and decline" (p. 21). Given that midlife likely constitutes the intersection of various areas of growth and decline (see Lachman, Teshale, & Agrigoroaei, 2015 for visual representation), uncontrollable stressors such as discrimination when individuals are supposed to be approaching and at this peak of life may be particularly detrimental to their psychological well-being. Specifically, midlife may be theorized as a time of relative social and financial stability (Alwin, McCammon, Mortimer, & Shanahan, 2003); to the extent that discrimination in different social and occupational contexts has the ability to disrupt this life stability, one may expect meaningful concurrent and longlasting impacts on well-being and on depressive symptoms.

These life course perspectives explaining the justification for our finding that sensitive periods for risk and protective factors occur when entering and in midlife align with those from the life stress field. For instance, stress sensitization theory would predict that, during midlife, daily and chronic stressors such as discrimination may more easily trigger bouts of depressive symptoms than at points earlier in development (Monroe & Harkness, 2005). Interestingly, the partially protective, buffering interactive effect of S&P occurs during the 30s a time where risk for depression due to different life stressors is gradually increasing (Mazure & Maciejewski, 2003). However, it is only at the peak of risk for depression due to life stressors, the 40s, where S&P demonstrated a fully protective effect in minimizing the typically harmful effect of discrimination on depressive symptoms. Our results, therefore, may illustrate the gradual start of this sensitive period in the 30s that then reaches its zenith during the 40s; S&P coping may be, thus, increasingly important in the 30s and maximally important and effective in the 40s relative to other times during adult development.

While theoretical scholarship has posited the existence and highlighted the potential importance of identifying sensitive periods during adulthood (Gee, Walsemann, & Brondolo, 2012; Gee et al., 2019), the broader field of psychology has studied sensitive periods almost exclusively in the context of early childhood development. Indeed, much work has focused on increased capacity for how factors such as early infant-mother separation and child maltreatment influences epigenetic changes implicated in the development of mental disorders (Roth & Sweatt, 2011), while other work has broadly pointed to the benefits of positive parenting "the earlier the better" to take advantage of sensitive periods in neurodevelopment occurring during the first months of life (Zeanah, Gunnar, McCall, Kreppner, & Fox, 2011). Our findings, in concert with previously cited empirical work proposing sensitive periods in adulthood for discrimination (e.g., Gee et al., 2012) and life-stress work pointing to variable risk for depressive episodes at different points in the life course (Mazure & Maciejewski, 2003), suggest that sensitive periods are not, however, exclusively found during early development. In studying sensitive periods during other periods of development, scholars' conceptions of the potential "window of opportunity" during which sensitive period may occur may need to be adjusted based on the developmental stage and phenomenon being studied. In our study, for instance, we operationalize a sensitive period for depression during the life course as a 10-year window, which stands in stark contrast to the month-long to week-long sensitive periods observed in human and animal models of early development (see Knudsen, 2004 for examples). Ultimately, much more empirical work is needed to uncover and understand the environmental, social, neural, genetic, and epigenetic mechanisms behind potential sensitive periods in midlife. These nascent efforts may be strengthened by adapting and synthesizing the schemas, models, and methods used when studying early childhood sensitive periods with those employed by those studying life course development.

Controlling for the effect of discrimination and S&P at each wave of data, we failed to find significant differences in the intercept and slope of depression between White and non-White participants. It is important to note that tests for racial differences in initial levels of depression and its trajectory were done controlling for discrimination, a robust predictor of greater depressive symptoms among ethnic/racial minorities at different stages of development (Benner et al., 2018; Paradies et al., 2015). Therefore, by controlling for discrimination, there may be no remaining difference in depressive trajectories between Whites and racially minoritized individuals, as the construct of race may have no longer served as a proxy variable for social disadvantage. Indeed, only 10.8% of our sample were from ethnically/racially minoritized groups, and despite our use of sampling weights to partially correct for this low level of representation, our sample sizes are too imbalanced to draw definitive conclusions as to the impact of race on the trajectory of depressive symptoms across the life course. In addition, given our small percentage of non-White participants, we made the decision to dichotomize race and preserve statistical power despite the ready admission that the lived experiences of various marginalized groups may differ widely and that this approach may not be sensitive to important between (i.e., between minoritized groups) and within-group variability (Hall, Yip, & Zárate, 2016). Future culturally sensitive longitudinal work on discrimination may be better suited to tease apart potential differences in the course of depression throughout the life span if they include balanced samples of Whites and individuals from multiple ethnically/ racially minoritized groups.

Limitations and Conclusions

Despite its contribution, this study has several limitations that set the stage for future work informed by a life course perspective. First, while our cohort sequential, or accelerated longitudinal design, allowed us to examine the trajectory of depression across a 49-year period, each cohort included individuals as far as 9 years apart in age. Our inferences are, thus, limited to the level of the decade, whereas the effects of discrimination and S&P coping likely also have more short-term effects on depressive trajectories. While difficult to extend over such a time period, future work should test for both short-term (e.g., 1-, 3-, or 5-year-) and longterm (10+-year) effects of these factors on the course of depression throughout the life span. These types of short-term longitudinal tests of S&P may be particularly useful in the study of child and adolescent development, where there are already wellestablished, albeit shorts sensitive periods that spur rapid developmental change.

Secondly, we chose to measure depressive symptoms and S&P as TVCs which assumes that these factors themselves do not change over time in a systematic way. While cognitively based coping strategies, such as S&P, may come online during adolescence (Compas et al., 2017), we are aware of no scholarly work that maps changes in the frequency of S&P coping across time. A limited amount of work has begun to examine how discrimination unfolds across long periods of time. For instance, using 3,032 participants from Wave I of MIDUS, Kessler and colleagues (1999) found that younger cohorts (age 25-44) reported greater everyday discrimination than did those over 65. However, because these differences are between individuals, inferences for how discrimination unfolds across the life span (i.e., age effects) cannot be inferred. Furthermore, it is unclear whether those observed differences in discrimination reflect what Gee et al. (2012) refer to as "age-patterned responses", or differences in prevalence due to stage in life, or period effects, or changes brought on by historical events and/or social transition. Indeed, participants who were over the age of 65 during MIDUS I data collection in 1995/ 1996 were young adults during the civil rights movement; living through this time may have both influenced what these individuals perceive (and report) as discrimination relative to those younger individuals in Kessler et al.'s (1999) MIDUS I sample who were not alive during this period of social change.

By contrast, in a sample of 7,225 working women followed for 15 years, Gee and colleagues (2007) found that age-based discrimination peaked in the 20s and 50s, was low during the 30s, and declined from the 50s peak around age 55 age discrimination, which implies that an individual's exposure to discrimination changes over as a function of their age. While Gee et al.'s (2007) study design is particularly useful in its ability to illustrate within-person change over time, such a design is not well suited to account for significant legal or social changes during this 15-year time frame (i.e., any period effects) that may have impacted women's exposure to workplace discrimination. While we have no reason to believe specific period effects significantly impacted our findings related to individual's trajectory depressive symptoms over time or age-based changes in the effects of discrimination and S&P on symptomatology, it is impossible to rule out the potential influence of small and unmeasured influences due to time and context. To better disaggregate potential period effects, which may be observed between cohorts, from time patterned responses, which may only be observed via intraindividual change over time, scholars should endeavor to examine

and compare intra-individual changes in factors such as discrimination between samples exposed and not exposed to potentially meaningful period effects (e.g., pre-post civil rights legislature, pre-post federal legalization of same-sex marriage, etc.).

Finally, a notable limitation examining the effects of discrimination in the MIDUS sample is the relative racial/ethnic homogeneity of the MIDUS sample. Although we utilized sampling weights to increase the influence of non-White participants in the estimation of our parameters, the fact remains that White participants made up almost 90% of our sample. This sample homogeneity limited both the amount of reported racial/ethnic discrimination we observed across the life course and limited our power to detect reliable differences in our estimates based on race. Future work would benefit from examining large, targeted samples of racially/ethnically minoritized individuals to better and more reliably observe the impact of racial discrimination on depressive symptoms throughout the life course. Indeed, if systematic change over time exists in discrimination, the pattern of change over time may vary based on the type of discrimination. While used as control variables, Gee and colleagues (2007) observed that patterns of racial and genderbased discrimination did not mirror patterns of age-based discrimination. Future work may, therefore, benefit from examining multiple types of discrimination concurrently and from using more restricted samples if the substantive interest is on understanding specific types of discrimination (e.g., using a Muslim-American sample to study religious and ethnic-racial discrimination).

Despite its limitations, the current study adds to our understanding of how depressive symptoms change over the life course, as well as how discrimination and S&P coping impacts depressive symptomatology both contemporaneously and up to 10 years later during sensitive periods in adult development. Our findings demonstrated that, among a representative sample of adults in the US, depression follows a slightly downward linear trajectory from ages 20 through 69. Our use of an accelerated longitudinal design allows us to approximate intra-individual change in depressive symptoms over time and move beyond population-based epidemiological studies that compare rates of depression between individuals at different ages. In addition, our findings add to the bodies of work that have independently examined the impact of discrimination at different points along the life course by showing that discrimination is predictive of greater depressive symptoms throughout human development over and above the natural negative trajectory of depression observed with aging. Finally, our study provides further evidence that, similar to what has been found with physiological health, S&P may also be protective for mental health. Chen et al. (2012), using MIDUS Waves I and II, found that adults who grew up in poverty but displayed high levels of S&P coping endorsed the lowest levels of physiological wear and tear, whereas our findings using MIDUS Waves I, II, and III suggest that S&P buffers against the harmful impact of discrimination on depressive symptoms in the 30s and is crucial in the 40s at protecting against the long-term impact of discrimination 10 years later. These findings point to potential sensitive periods of human development (Gee et al., 2019) where the impact of S&P may be particularly important in promoting psychological well-being. In conclusion, this study adds to the field by adopting developmental and life course perspective to the study of how discrimination and S&P coping impact depressive symptoms from ages 20 to 69.

Supplementary material. The supplementary material for this article can be found at https://doi.org/10.1017/S0954579421000146

Financial Support. This work was supported in part by a predoctoral fellowship provided by the National Institute of Child Health and Human Development (T32-HD07376) through the Carolina Consortium on Human Development, University of North Carolina at Chapel Hill, to N. Keita Christophe

Conflicts of Interest. None.

References

- Alwin, D. F., & McCammon, R. J. (2003). Generations, cohorts, and social change. In J. T. Mortimer & M. J. Shanahan (Eds.), *Handbook of the life course* (pp. 23–49). US: Springer. doi:10.1007/978-0-306-48247-2_2
- Benner, A. D., Wang, Y., Shen, Y., Boyle, A. E., Polk, R., & Cheng, Y.-P. (2018). Racial/ethnic discrimination and well-being during adolescence: A metaanalytic review. *The American Psychologist*, 73, 855–883. doi:10.1037/ amp0000204
- Blöchl, M., & Nestler, S. (2019). Vascular risk factors and long-term changes of depressed mood: Findings from two population-based longitudinal studies [Preprint]. Open Science Framework. doi:10.31219/osf.io/u2j8h.
- Bollen, K. A., & Curran, P. J. (2005). Latent curve models: A structural equation perspective (1st ed.). Hoboken, NJ: John Wiley & Sons, Ltd. doi:10.1002/ 0471746096
- Brody, G. H., Yu, T., Chen, E., & Miller, G. E. (2020). Persistence of skin-deep resilience in African American adults. *Health Psychology*, 39, 921–926. doi:10.1037/hea0000945
- Brody, G. H., Yu, T., Miller, G. E., & Chen, E. (2016). Resilience in adolescence, health, and psychosocial outcomes. *Pediatrics*, 138, doi:10.1542/ peds.2016-1042
- Charles, S. T., Reynolds, C. A., & Gatz, M. (2001). Age-related differences and change in positive and negative affect over 23 years. *Journal of Personality* and Social Psychology, 80, 136–151.
- Chen, E. (2012). Protective factors for health among low-socioeconomic-status individuals. *Current Directions in Psychological Science*, *21*, 189–193. doi:10.1177/0963721412438710
- Chen, P., & Harris, K. M. (2019). Association of positive family relationships with mental health trajectories from adolescence to midlife. JAMA Pediatrics, 173, e193336–e193336. doi:10.1001/jamapediatrics.2019.3336
- Chen, E., Lee, W. K., Cavey, L., & Ho, A. (2013). Role models and the psychological characteristics that buffer low-socioeconomic-status youth from cardiovascular risk. *Child Development*, 84, 1241–1252. doi:10.1111/cdev.12037
- Chen, E., McLean, K. C., & Miller, G. E. (2015). Shift-and-persist strategies: Associations with socioeconomic status and the regulation of inflammation among adolescents and their parents. *Psychosomatic Medicine*, 77, 371–382.
- Chen, E., & Miller, G. E. (2012). "Shift-and-persist" strategies: Why low socioeconomic status isn't always bad for health. *Perspectives on Psychological Science*, 7, 135–158. doi:10.1177/1745691612436694
- Chen, E., Miller, G. E., Lachman, M. E., Gruenewald, T. L., & Seeman, T. E. (2012). Protective factors for adults from low childhood socioeconomic circumstances: The benefits of shift-and-persist for allostatic load. *Psychosomatic Medicine*, 74, 178–186. doi:10.1097/PSY.0b013e31824206fd
- Chen, E., Strunk, R. C., Trethewey, A., Schreier, H. M. C., Maharaj, N., & Miller, G. E. (2011). Resilience in low-socioeconomic-status children with asthma: Adaptations to stress. *Journal of Allergy and Clinical Immunology*, 128, 970–976. doi:10.1016/j.jaci.2011.06.040
- Christophe, N. K., Stein, G. L., Martin Romero, M. Y., Chan, M., Jensen, M., Gonzalez, L. M., & Kiang, L. (2019). Coping and culture: The protective effects of shift-&-persist and ethnic-racial identity on depressive symptoms in Latinx youth. *Journal of Youth and Adolescence*, 48, 1592–1604. doi:10.1007/s10964-019-01037-8
- Compas, B. E., Jaser, S. S., Bettis, A. H., Watson, K. H., Gruhn, M. A., Dunbar, J. P., ... Thigpen, J. C. (2017). Coping, emotion regulation, and psychopathology in childhood and adolescence: A meta-analysis and narrative review. *Psychological Bulletin*, 143, 939–991. doi:10.1037/bul0000110

- Gee, G. C., Hing, A., Mohammed, S., Tabor, D. C., & Williams, D. R. (2019). Racism and the life course: Taking time seriously. *American Journal of Public Health*, 109, S43–S47. doi:10.2105/AJPH.2018.304766
- Gee, G. C., Pavalko, E. K., & Long, J. S. (2007). Age, cohort and perceived age discrimination: Using the life course to assess self-reported age discrimination. *Social Forces*, 86, 265–290. doi:10.1353/sof.2007.0098
- Gee, G. C., Walsemann, K. M., & Brondolo, E. (2012). A life course perspective on how racism may be related to health inequities. *American Journal of Public Health*, 102, 967–974. doi:10.2105/AJPH.2012.300666
- Gillis, J. C., Chang, S.-C., Wang, W., Simon, N. M., Normand, S.-L., Rosner, B. A., ... Okereke, O. I. (2019). The relation of telomere length at midlife to subsequent 20-year depression trajectories among women. *Depression and Anxiety*, 36, 565–575. doi:10.1002/da.22892

Hall, G. S. (1922). Senescence, the last half of life. D. Appleton.

- Hall, G. C. N., Yip, T., & Zárate, M. A. (2016). On becoming multicultural in a monocultural research world: A conceptual approach to studying ethnocultural diversity. *American Psychologist*, 71, 40–51. doi:10.1037/ a0039734
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6, 1–55. doi:10.1080/ 10705519909540118
- Isiordia, M., & Ferrer, E. (2018). Curve of factors model: A latent growth modeling approach for educational research. *Educational and Psychological Measurement*, 78, 203–231. doi:10.1177/0013164416677143
- Kallem, S., Carroll-Scott, A., Rosenthal, L., Chen, E., Peters, S. M., McCaslin, C., & Ickovics, J. R. (2013). Shift-and-persist: A protective factor for elevated BMI among low-socioeconomic-status children. *Obesity*, 21, 1759–1763. doi:10.1002/oby.20195
- Kessler, R. C., Andrews, G., Colpe, L. J., Hiripi, E., Mroczek, D. K., Normand, S.-L. T., ... Zaslavsky, A. M. (2002). Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological Medicine*, 32, 959–976. doi:10.1017/ S0033291702006074
- Kessler, R. C., Foster, C., Webster, P. S., & House, J. S. (1992). The relationship between age and depressive symptoms in two national surveys. *Psychology* and Aging, 7, 119–126. doi:10.1037/0882-7974.7.1.119
- Kessler, R. C., Mickelson, K. D., & Williams, D. R. (1999). The prevalence, distribution, and mental health correlates of perceived discrimination in the United States. *Journal of Health and Social Behavior*, 40, 208–230. doi:10.2307/2676349
- Keyes, C. L. M., Shmotkin, D., & Ryff, C. D. (2002). Optimizing well-being: The empirical encounter of two traditions. *Journal of Personality and Social Psychology*, 82, 1007–1022. doi:10.1037/0022-3514.82.6.1007
- Knudsen, E. I. (2004). Sensitive periods in the development of the brain and behavior. *Journal of Cognitive Neuroscience*, 16, 1412–1425. doi:10.1162/ 0898929042304796
- Lachman, M. E., Teshale, S., & Agrigoroaei, S. (2015). Midlife as a pivotal period in the life course: Balancing growth and decline at the crossroads of youth and old age. *International Journal of Behavioral Development*, 39, 20–31. doi:10.1177/0165025414533223
- Lam, P. H., Miller, G. E., Chiang, J. J., Levine, C. S., Le, V., Shalowitz, M. U., ... Chen, E. (2018). One size does not fit all: Links between shift-and-persist and asthma in youth are moderated by perceived social status and experience of unfair treatment. *Development and Psychopathology*, 30, 1699–1714. doi:10.1017/S0954579418000913
- Lazarus, R. S., & Folkman, S. (1984). *Stress, appraisal, and coping* (1st ed.). New York, NY: Springer Publishing Company.
- Luo, Y., Xu, J., Granberg, E., & Wentworth, W. M. (2012). A longitudinal study of social status, perceived discrimination, and physical and emotional health among older adults. *Research on Aging*, 34, 275–301. doi:10.1177/ 0164027511426151
- Mazure, C. M., & Maciejewski, P. K. (2003). A model of risk for major depression: Effects of life stress and cognitive style vary by age. *Depression and Anxiety*, 17, 26–33. doi:10.1002/da.10081
- Mehta, P. D., & West, S. G. (2000). Putting the individual back into individual growth curves. *Psychological Methods*, 5(1), 23–43. doi:10.1037/1082-989X.5.1.23

- Miller, G. E., Yu, T., Chen, E., & Brody, G. H. (2015). Self-control forecasts better psychosocial outcomes but faster epigenetic aging in low-SES youth. *Proceedings of the National Academy of Sciences*, 112, 10325– 10330. doi:10.1073/pnas.1505063112
- Monroe, S. M., & Harkness, K. L. (2005). Life stress, the "kindling" hypothesis, and the recurrence of depression: Considerations from a life stress perspective. *Psychological Review*, 112, 417–445. doi:10.1037/ 0033-295X.112.2.417
- Paradies, Y., Ben, J., Denson, N., Elias, A., Priest, N., Pieterse, A., ... Gee, G. (2015). Racism as a determinant of health: A systematic review and metaanalysis. *PLoS One*, 10, e0138511. doi:10.1371/journal.pone.0138511
- Pascoe, E. A., & Richman, L. S. (2009). Perceived discrimination and health: A meta-analytic review. *Psychological Bulletin*, 135, 531–554. doi:10.1037/ a0016059
- Patrick, C., Curtin, J., & Tellegen, A. (2002). Development and validation of a brief form of the multidimensional personality questionnaire. *Psychological Assessment*, 14, 150–163. doi:10.1037/1040-3590.14.3.262
- Prenda, K. M., & Lachman, M. E. (2001). Planning for the future: A life management strategy for increasing control and life satisfaction in adulthood. *Psychology and Aging*, 16, 206–216. doi:10.1037/0882-7974.16.2.206
- Radler, B. T. (2014). The midlife in the United States (MIDUS) series: A national longitudinal study of health and well-being. *Open Health Data*, 2), doi:10.5334/ohd.ai
- Rossi, A. S. (2001). Caring and doing for others: Social responsibility in the domains of family, work, and community. Chicago, IL: University of Chicago Press.
- Roth, T. L., & Sweatt, J. D. (2011). Annual research review: Epigenetic mechanisms and environmental shaping of the brain during sensitive periods of development. *Journal of Child Psychology and Psychiatry*, 52, 398–408. doi:10.1111/j.1469-7610.2010.02282.x
- Satorra, A. (2000). Scaled and adjusted restricted tests in multi-sample analysis of moment structures. In R. D. H. Heijmans, D. S. G. Pollock & A. Satorra (Eds.), *Innovations in multivariate statistical analysis: A festschrift for heinz neudecker* (pp. 233–247). US: Springer. doi:10.1007/ 978-1-4615-4603-0_17.
- Schmitt, M. T., Branscombe, N. R., Postmes, T., & Garcia, A. (2014). The consequences of perceived discrimination for psychological well-being: A metaanalytic review. *Psychological Bulletin*, 140, 921–948. doi:10.1037/a0035754
- Slavich, G. M., & Irwin, M. R. (2014). From stress to inflammation and major depressive disorder: A social signal transduction theory of depression. *Psychological Bulletin*, 140, 774–815. doi:10.1037/a0035302
- Stanisławski, K. (2019). The coping circumplex model: An integrative model of the structure of coping with stress. *Frontiers in Psychology*, 10, doi:10.3389/ fpsyg.2019.00694
- Stein, G. L., Jensen, M., Cruz, R. A., Christophe, N. K., Martin Romero, M., & Robbins, R. (under review). Shift and persist in Mexican-origin youth: A longitudinal test.
- Stokes, J. E. (2019). Social integration, perceived discrimination, and self-esteem in mid- and later life: Intersections with age and neuroticism. Aging & Mental Health, 23, 727-735. doi:10.1080/13607863.2018. 1450834
- Sumner, R., Burrow, A. L., & Hill, P. L. (2018). The development of purpose in life among adolescents who experience marginalization: Potential opportunities and obstacles. *American Psychologist*, 73, 740–752. doi:10.1037/ amp0000249
- Tackett, J. L., & Lahey, B. B. (2016). Neuroticism. In T. A. Widiger (Ed.), *The Oxford handbook of the five factor model* (Vol. 1, pp. 39–56). Oxford, England: Oxford University Press. doi:10.1093/oxfordhb/9780199352487. 013.14
- Umaña-Taylor, A. J., Quintana, S. M., Lee, R. M., Cross, W. E., Rivas-Drake, D., Schwartz, S. J., ... Seaton, E. & Ethnic and Racial Identity in the 21st Century Study Group (2014). Ethnic and racial identity during adolescence and into young adulthood: An integrated conceptualization. *Child Development*, 85, 21–39. doi:10.1111/cdev.12196
- Walsemann, K. M., Gee, G. C., & Geronimus, A. T. (2009). Ethnic differences in trajectories of depressive symptoms: Disadvantage in family background, high school experiences, and adult characteristics. *Journal of Health and Social Behavior*, 50, 82–98. doi:10.1177/002214650905000106

- Williams, D. R., & Mohammed, S. A. (2009). Discrimination and racial disparities in health: Evidence and needed research. *Journal of Behavioral Medicine*, 32, 20–47. doi:10.1007/s10865-008-9185-0
- Williams, D. R., Neighbors, H. W., & Jackson, J. S. (2003). Racial/ethnic discrimination and health: Findings from community studies. *American Journal of Public Health*, 93, 200–208. doi:10.2105/AJPH.93.2.200
- Williams, D. R., Yan, Y., Jackson, J. S., & Anderson, J. S. (1997). Racial differences in physical and mental health: Socio-economic status, stress and discrimination. *Journal of Health Psychology*, 2, 335–351. doi:10.1177/ 135910539700200305
- Wrosch, C., Heckhausen, J., & Lachman, M. E. (2000). Primary and secondary control strategies for managing health and financial stress across adulthood. *Psychology and Aging*, 15, 387–399. doi:10.1037/0882-7974.15.3.387
- Yip, T., Wang, Y., Mootoo, C., & Mirpuri, S. (2019). Moderating the association between discrimination and adjustment: A meta-analysis of ethnic/ racial identity. *Developmental Psychology*, 55, 1274–1298. doi:10.1037/ dev0000708
- Zeanah, C. H., Gunnar, M. R., McCall, R. B., Kreppner, J. M., & Fox, N. A. (2011). Sensitive periods. *Monographs of the Society for Research in Child Development*, 76, 147–162. doi:10.1111/j.1540-5834.2011.00631.x