

pyramidal reaction, which may need to be countered by antiparkinsonian drugs.

M. H. ABENSON.

*Director of Psychiatry,
Kaplan Hospital,
Rehovoth,
Israel.*

PSYCHOTHERAPY WITH FAILURES OF PSYCHOANALYSIS

DEAR SIR,

I should like to comment on two patients who were recently referred to me, having received psychoanalysis with no improvement.

Case 1

Male aged 50, single, suffered from an obsessional neurosis, in particular agoraphobia, over 20 years. He had had five years of psychoanalysis with no improvement, and was sent to me by an analyst because his condition was worsening and he was no longer suitable for analysis.

I attempted to treat him by behaviour therapy, and tried to persuade him that a new approach to his treatment could be helpful. I was impressed by his continued reference to his previous therapist, whom he regarded as a supreme being. He continually stated that only analysis could help him, despite his worsening condition. After four visits he informed me that he could not accept behaviour therapy, because he *knew* it would not help him.

Case 2

Female aged 44, divorced. A new immigrant with adaptation problems complicated by an aggressive dependent personality. She had ten years of continuous psychotherapy with two well-known analysts with no improvement, before she decided to emigrate (running away from her therapist)? I treated her with encouragement and supportive therapy, purposively avoiding analytic explanations. She rebelled against this and discontinued treatment after two months. Two months later she contacted me, requesting further advice; she appeared self-sufficient and well on the way to integration in a new 'society' but still spoke of her analysts with considerable awe.

These cases illustrate some of the points made by Dr. Schmidberg (1) and some of the difficulties psycho-analysis creates for patients. Analysts are generally very vague in their criteria for the selection of patients. Moreover, the process of treatment appears to 'brainwash' the patients, and thus make other methods of treatment unacceptable. The first case, an obsessional neurotic, is an extreme example—a person totally unsuited for psychoanalysis who was unfortunately unable to accept a course of treatment that could help, if not cure his disablement. The second patient seemed to fit some of the main criteria for psychoanalysis (intelligent patient with personality problems) but did not improve; and later although

she responded to a simpler and shorter form of treatment she continued to find non-analytic reasoning difficult to accept. This case also illustrates that 'failed' cases of psychoanalysis can be helped by other forms of psychotherapy.

I have always found it difficult to discover what are the criteria of successfully treated cases of psychotherapy. Here again, analysts are vague in their answers. I personally find that many psychoanalytic concepts of thought formation are useful in the understanding of patients, but consider rigid adherence to one particular method of treatment is destructive. This is also true in other spheres of medicine.

M. H. ABENSON.

*Director of Psychiatry,
Kaplan Hospital,
Rehovot,
Israel.*

REFERENCE

- SCHMIDBERG, M. (1970). Psychotherapy with failures of psychoanalysis. *Brit. J. Psychiat.*, **116**, 195–200.

THE MAUDSLEY HOSPITAL CLINICAL SENSORIUM TEST

DEAR SIR,

We have recently undertaken a revision and adaptation to American conditions of the battery of questions described under the headings of clinical sensorium test (Part I), clinical questionnaire, and memory questionnaire, used in the Institute of Psychiatry at the Maudsley Hospital for assessment of mental function in elderly patients (Ball *et al.*, 1967; Shapiro *et al.*, 1956). We have not yet undertaken tests of validity and reliability, and before proceeding with these we should be grateful for the comments of any persons likely to be interested in using such a test. Copies may be obtained by writing to Mr. Bruce Boltuch at the Geriatric-Psychiatric Unit, Building #12, Bergen Pines County Hospital, Paramus, N.J. 07652.

BRUCE BOLTUCH.
DAVID PETER BIRKETT.

*Bergen Pines,
County Hospital,
Paramus, N.J. 07652,
U.S.A.*

REFERENCES

- BALL, J. A. C., and TAYLOR, A. R. (1967). 'Effects of cyclandelate on mental function and cerebral blood flow in elderly patients.' *British Medical Journal*, **iii**, 525–8.
- SHAPIRO, M. D., POST, F., LOFVING, B., and INGLIS, J. (1956). 'Memory function in psychiatric patients over sixty, some methodological and diagnostic implications.' *Journal of Mental Science*, **102**, 233–46.