



Original Article

Cite this article: Wikner A, Johansson K, Enocson E, Sthen Bergdahl M, Hansson L, Rydberg A, and Sandberg C (2024) Lower bone strength in young patients with Fontan circulation compared to controls. *Cardiology in the Young* **34**: 1487–1492. doi: [10.1017/S1047951124000404](https://doi.org/10.1017/S1047951124000404)

Received: 4 October 2023

Revised: 22 January 2024

Accepted: 5 February 2024




First published online: 7 March 2024

Keywords:

Strength-strain index; bone mineral content; bone mineral density; total cavo-pulmonary connection; CHD; children; adolescents

Corresponding author:

A. Wikner; Email: anna.wikner@umu.se

Anna Wikner¹ , Karna Johansson¹, Elin Enocson¹, Magne Sthen Bergdahl², Lena Hansson² , Annika Rydberg¹ and Camilla Sandberg^{1,3} 

¹Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden; ²Department of Clinical Sciences, Pediatrics, Umeå University, Umeå, Sweden and ³Department of Community Medicine and Rehabilitation, Physiotherapy, Umeå University, Umeå, Sweden

Abstract

Objectives: Previous reports indicate bone deficits in patients with Fontan circulation. However, the consequences of these deficits on bone strength and when these changes occur are unclear. **Aim:** To compare the tibial bone strength-strain index between young patients (6–19 years) with Fontan circulation and age- and sex-matched controls, and to determine strength-strain index in subgroups of children (6–12 years) and adolescents (13–19 years) versus controls. **Method:** The tibia was examined with peripheral quantitative CT. Based on the assessed data, bone strength-strain index was calculated in the lateral and anterior-posterior directions. **Results:** Twenty patients with Fontan and twenty controls (mean age 13.0 ± 4.4 years; 50% females) were examined. Patients had a lower strength-strain index in the lateral direction compared to controls ($808.4 \pm 416.8\text{mm}^3$ versus $1162.5 \pm 552.1\text{mm}^3$, $p = 0.043$). Subgroup analyses showed no differences regarding strength-strain index in children (6–12 years) with Fontan circulation compared to controls. However, the adolescents (13–19 years) with Fontan circulation had lower strength-strain indexes in both the lateral and anterior-posterior directions compared to controls ($1041.4 \pm 299.8\text{mm}^3$ versus $1596.4 \pm 239.6\text{mm}^3$, $p < 0.001$, and $771.7 \pm 192.4\text{mm}^3$ versus $1084.9 \pm 215.0\text{mm}^3$, $p = 0.004$). When adjusted for height, there were differences between patients (6–19 years) and controls in strength-strain indexes in both the lateral and anterior-posterior directions. In subgroup analyses, the results remained robust. **Conclusion:** Young patients (6–19 years) with Fontan circulation have a lower strength-strain index in the tibia compared to controls. Subgroup analyses show that this deficit is mainly driven by the differences in adolescents (13–19 years), which might suggest that bone strength decreases with age.

Surgical and medical treatments of CHD have substantially improved survival.^{1,2} Despite success in survival, extra-cardiac consequences of the CHD may constitute an increasing and unexpected concern.³ Recent reports have highlighted deficits in bone mineral density in patients with complex CHD,^{4,5} especially in patients with Fontan circulation.^{3,6–11} Also, it was suggested that these deficits increase with age (7).

Most of the previous reports are based on data retrieved from dual-energy X-ray absorptiometry, which provides a two-dimensional projection of the bone, and areal bone mineral density is the most common outcome.^{4,6,7,10} Impaired bone mineral density contributes to a lower bone strength that in the long-term, if impairment progresses, increases the risk of fractures.¹²

An alternative method to assess bone health is peripheral quantitative CT. Peripheral quantitative CT provides a three-dimensional projection and enables separate analyses of trabecular and cortical bone. It also allows for calculation of the bone strength based not only on density but also on geometrical properties of the bone such as cortical thickness and diameter, *i.e.* strength-strain index.^{13–15}

A previous report using peripheral quantitative CT showed lower trabecular density, cortical thickness, and strength-strain index in the radius of children (mean age 12 years) with Fontan circulation in comparison to controls.⁸ Similar findings were reported in a small sample of adolescents and young adults with Fontan circulation ($n = 6$, age 14–24 years).¹¹ Additionally, lower bone mineral density and cortical thickness were found in the tibia of patients with Fontan circulation (age 5–33 years) when compared to reference data.⁹ Furthermore, a reduced strength-strain index was shown in the tibia and the radius in adults with Fontan circulation, despite findings of a normal bone mineral density.⁵ Taken together, recent reports indicate bone deficits in patients with Fontan circulation. However, the consequences these deficits have on the strength-strain index are unclear. It is also unclear when in the lifespan these deficits occur.

The primary aim of the present study was to determine differences in the tibia strength-strain index between young patients (6–19 years) with Fontan circulation and age- and sex-matched

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

controls. Other aims were to determine (a) differences in trabecular and cortical bone measures between patients and controls, (b) the above-mentioned bone variables in subgroups of children (6–12 years) and adolescents (13–19 years), and (c) the associations between clinical data, self-reported physical activity, and the tibia strength-strain index.

Materials and method

Study population

There were twenty-two eligible participants with Fontan circulation (6–19 years), and of these twenty agreed to participate. They were recruited from the Pediatric Cardiology Units in Västerbotten, Västernorrland, and Norrbotten. Most of the study population ($n = 18$) participated in a previous cross-sectional research project regarding muscle function,^{16,17} body composition (assessed with dual-energy X-ray absorptiometry), and vitamin D intake.¹⁸ Inclusion criteria were Fontan circulation and an age of 6–19 years. In addition, twenty age- and sex-matched controls were recruited by convenience sampling, *i.e.* from the population near the study centre. Exclusion criteria for both patients and controls were presence of diagnosis known to affect the bone health, *e.g.* juvenile rheumatoid arthritis, cognitive and neurological disabilities, or limited knowledge of the Swedish language. Data on diagnosis, age at the total cavo-pulmonary connection surgery, oxygen saturation, systemic ventricle, weight, height, presence of growth pain in the legs, and blood test results (*i.e.* pro-BNP, 25-hydroxy-vitamin D, gamma-glutamyltransferase) were retrieved at the clinical visit and via the medical record. Prior to participation, a consent form was signed by both parents if the participant was under 18 years of age. Participants that were ≥ 18 years of age provided a written informed consent themselves. The study conformed with the principles outlined by the declaration of Helsinki¹⁶ and was approved by the Regional Ethics Review Board, Umeå (Dnr: 2016-445-31 M, 2018-07-32M).

Peripheral quantitative computed tomography

The bone and soft tissue composition of the tibia were assessed with peripheral quantitative CT (pQCT, Stratec XCT2000, Medizintechnik GmbH, Pforzheim, Germany) at Livsmedicin in Umeå. Peripheral quantitative CT enables a non-invasive and three-dimensional evaluation of the microarchitecture and strength of the peripheral bones, most commonly of the radius or the tibia. The equipment was calibrated prior to the examination and the voxel size was $0.5 \times 0.5 \times 2.0$ mm (with slice thickness of 2 mm). The tibia length was measured manually with a tape measure from the medial malleolus to the medial site of the knee joint. The examinations were made at 4 and 66% of the total length of the bone in the proximal direction. An illustrative description of projection sites was previously published.⁵ Variables regarding the trabecular bone and total bone mineral content were extracted from the distal tibia at the 4% site where the cortex is thin and most of the bone cross-sectional area consists of trabecular bone. Variables regarding the cortical bone were extracted from the proximal tibia at the 66% site since the cortical bone at this site is thick and nearly circular in shape. Data on muscle and fat tissue were also extracted from the 66% site. In addition, the strength-strain index was calculated based on bone geometry and bone mineral density data extracted from the 66% site. The strength-strain index represents the resistance to bending in the lateral and anterior-posterior direction with a higher value indicating that the

bone is more resistant to bending before fracturing. This variable has been evaluated in a bending test, showing that the strength-strain index predicted approximately 80% of the force required to fracture the bone at the diaphysis.¹⁹

Physical activity

Data on habitual physical activity and self-reported participation in free-time sports were retrieved at the time of peripheral quantitative CT analysis. Hours spent physically active during the week, *i.e.* physical activity in school, free-time sports, and other habitual physical activities, were added to the total number of hours.

Biochemical analyses

Venous blood was sampled in the fasting state to analyse 25-hydroxy-vitamin D (nmol/l), gamma-glutamyltransferase (μ -kat/L) and pro-BNP (ng/L). Prior to blood sampling, local anaesthetic cream (EMLA, Astra Zeneca) was applied. The samples were analysed at the Department of Clinical Chemistry, Umeå University Hospital. More detailed information is presented elsewhere.¹⁸

Statistical analysis

Statistical analyses were performed using IBM SPSS statistics 27.0 (IBM, Armonk, NY, USA). Data were assessed for normality and presented as means with standard deviations. Frequencies were presented as numbers with percentages. For comparison, Chi² test was used for categorical variables, and student's t-test or Mann-Whitney *U*-test was used for analyses of continuous variables. Cohen's *D* was used to calculate the effect size of the peripheral quantitative CT variables. A one-way analysis of covariance was conducted to determine if the difference in bone strength could be attributed to differences in height between patients and controls. The dependent variable was strength-strain index, the independent variable was the two groups of patients and controls, and the covariate was height. Univariate linear regression analyses were performed to determine factors associated with strength-strain index in the lateral direction in patients. The independent variables were chosen due to their suggested influence on the bone quality and strength, *i.e.* age at total cavo-pulmonary connection surgery, oxygen saturation, systemic ventricle, weight, height, and blood test results (*i.e.* pro-BNP, 25-hydroxy-vitamin D, gamma-glutamyltransferase). A multivariable linear regression model was built by entering variables significant in univariate analyses ($p < 0.15$). The variables in the multivariable model were checked for multicollinearity. To ensure that the associations remained robust, a sensitivity analysis of the model was performed by stepwise excluding one independent variable at the time. A p -value < 0.05 was considered significant.

Results

Study population

For the study group of twenty patients with Fontan circulation and twenty sex- and age-matched controls, the mean age was 13.0 ± 4.4 years and 50% were females. There were no differences between patients and controls in height or weight. Both groups participated in school sports to the same extent. However, the patients had a lower self-reported participation in free-time sports compared to the controls. Also, the total hours of self-reported physical activity

during the week were lower among the patients. There was no difference between patients and controls regarding the presence of growth pain in legs (Supplementary Table S1).

The most common cardiac diagnoses were hypoplastic left heart syndrome and pulmonary atresia. Sixty-five per cent of the patients had a left systemic ventricle. Most of the patients had sufficient 25-hydroxy-vitamin D levels. The mean gamma-glutamyltransferase was slightly increased and pro-BNP was within the normal range (Supplementary Table S2).

Bone variables assessed with peripheral quantitative CT

The patients with Fontan circulation had a lower strength-strain index in the lateral direction and a lower total bone mineral content of the tibia compared to controls (Table 1). No other differences were found regarding bone or soft tissue measures (Supplementary Table S3). However, in subgroup analyses, there were no differences regarding strength-strain index or total bone mineral content in the tibia of children (6–12 years, $n = 9$) with Fontan circulation compared to controls ($n = 9$). Nevertheless, children with Fontan circulation had a lower trabecular density and a lower cortical thickness than controls. In addition, adolescents (13–19 years, $n = 11$) with Fontan circulation had a lower total bone mineral content and a lower strength-strain index in both the lateral and anterior-posterior direction compared to the controls ($n = 11$). Also, the adolescents had smaller cortical and trabecular cross-sectional areas than controls (Table 1).

Height as a covariate

After adjusting for height, the difference in tibia strength-strain index in the lateral direction remained between patients and controls [$F(1, 37) = 12.3, p = 0.001, n^2 = 0.25$]. This difference also remained in the subgroup of adolescents (13–19 years) [$F(1, 19) = 14.7, p = 0.001, n^2 = 0.44$]. In the corresponding analysis in the subgroup of children (6–12 years), there was no difference in strength-strain index in the lateral direction when compared to controls (data not shown).

When using tibia strength-strain index in the anterior-posterior direction as the dependent variable, a difference was found between patients and controls whilst adjusting for height [$F(1, 37) = 4.8, p = 0.036, n^2 = 0.114$]. This difference remained for the subgroup of adolescents (13–19 years) [$F(1, 19) = 6.5, p = 0.02, n^2 = 0.25$], yet no difference was found in the subgroup of children (6–12 years) when compared to controls (data not shown).

Factors associated with bone strength

In univariate linear regression analysis, height, weight, and age at total cavo-pulmonary connection was associated with tibia strength-strain index in the lateral direction. No other variables were associated with the strength-strain index in the lateral direction. In the multivariable mode, height was associated with strength-strain index in the lateral direction whilst adjusting for age at total cavo-pulmonary connection (Table 2).

Discussion

To the best of our knowledge, this is the first study reporting a lower tibia strength-strain index assessed with peripheral quantitative CT in young patients (6–19 years) with Fontan circulation when compared to controls. In addition, a lower tibia bone mineral content was found in the patients.

Bone strength and bone mineral content

At present, there are only a few reports assessing strength-strain index and total bone mineral content with peripheral quantitative CT in patients with Fontan circulation. A previous study found no difference in tibia strength-strain index in young patients with Fontan circulation compared to controls.⁸ However, in another study, a lower tibia strength-strain index and a lower total bone mineral content were shown in adults with Fontan circulation when compared to controls,⁵ which is in line with our report on children and adolescents.

In the present report, subgroup analyses show that the children (6–12 years) with Fontan circulation had a lower cortical thickness and trabecular bone mineral density compared to the controls. This conforms with a previous study assessing bone quality in children with Fontan circulation with peripheral quantitative CT.⁹ However, these differences in cortical thickness and trabecular bone mineral density were not present in the subgroup of adolescents, which is difficult to explain. Instead, lower trabecular and cortical cross-sectional areas and bone mineral content were found in this group. Although speculative, a lower cross-sectional area in combination with a lower total bone mineral content could lead to a proportionally smaller bone, thereby maintaining the density within the normal range. In addition, similar findings of a lower strength-strain index among both adolescents and adults with Fontan circulation⁵ suggest the development of bone deficits over time. Similar findings of a decrease in bone health with age were reported in children and adolescents with cystic fibrosis.²⁰

Since the peak bone mass is usually reached before the age of 18,²¹ the presence of a lower bone strength at a young age may pose a higher risk of future fractures compared to the general population. However, according to recent reports, there are no indications of an increased risk of fractures in young patients (<18 years) with complex CHD.^{22,23} Despite lower bone strength, it is important to keep in mind that this is still a young population with no indication of osteoporosis *per se*. However, there is a slightly increased risk for fractures among older patients with CHD compared to controls.²² Therefore, it is important to monitor the bone development over time to detect further deterioration of bone quality, and if possible, even minimise future fractures.

Height as a covariate

There are previous reports on a shorter stature in patients with Fontan circulation,²⁴ yet no significant differences regarding height were found in the present cohort. However, to ensure that the differences in strength-strain index were not due to differences in height, post hoc analyses adjusting for height were conducted. When adjusted for height, an additional difference was found in the anterior-posterior direction of the strength-strain index between patients and controls. The rest of the results remained robust.

Factors associated with bone quality

A number of factors are known to affect bone quality and bone development, e.g. vitamin D levels,²⁵ presence of hypoxia,²⁶ physical activity,²¹ and growth trajectory.²⁷ Previous reports have shown that vitamin D deficits are common among young patients with Fontan circulation. However, these reports failed to show an association between vitamin D levels and bone mineral density.^{7,18} Also, in the present report, vitamin D levels were not associated with strength-strain index or bone mineral content. Furthermore, hypoxia has been suggested as a factor that could affect bone

Table 1. Tibia bone measures assessed with peripheral quantitative computed tomography

Variable	Unit	Patients (n = 20)	Controls (n = 20)	p	Effect size
Age 6–19					
Dimensions					
Trabecular CSA ¹	mm ²	399.5 ± 101.1	467.1 ± 106.6	0.06	–0.65
Cortical CSA ¹	mm ²	197.8 ± 79.2	246.11 ± 87.2	0.1	–0.58
Cortical thickness	mm	3.2 ± 0.9	3.6 ± 0.9	0.1	–0.47
Mass and density					
Total BMC ²	g	2.6 ± 0.7	3.17 ± 0.6	0.009	–0.89
Trabecular BMD ³	mg/cm ³	242.9 ± 42.6	277.1 ± 59.9	0.06	–0.66
Cortical BMD ³	mg/cm ³	1069.1 ± 62.7	1064.5 ± 58.6	0.8	0.09
Strength					
xSSI ⁴	mm ³	808.4 ± 416.8	1162.5 ± 552.1	0.043	–0.72
ySSI ⁵	mm ³	610.0 ± 289.1	811.0 ± 365.3	0.1	–0.61
Age 6–12					
		Patients (n = 9)	Controls (n = 9)		
Dimensions					
Trabecular CSA ¹	mm ²	335.9 ± 89.6	385.3 ± 91.3	0.4	–0.55
Cortical CSA ¹	mm ²	137.3 ± 62.9	160.0 ± 36.6	0.3	–0.44
Cortical thickness	mm	2.5 ± 0.7	2.9 ± 0.3	0.024	–0.58
Mass and density					
Total BMC ²	g	2.2 ± 0.7	2.8 ± 0.6	0.1	–0.89
Trabecular BMD ³	mg/cm ³	259.3 ± 55.0	317.96 ± 62.3	0.019	–0.99
Cortical BMD ³	mg/cm ³	1021.2 ± 56.6	1021.35 ± 30.6	0.7	–0.003
Strength					
xSSI ⁴	mm ³	523.6 ± 366.4	632.2 ± 277.6	0.4	–0.33
ySSI ⁵	mm ³	413.5 ± 269.9	476.2 ± 172.7	0.5	–0.28
Age 13–19					
		Patients (n = 11)	Controls (n = 11)		
Dimensions					
Trabecular CSA ¹	mm ²	451.5 ± 79.9	534.0 ± 63.2	0.02	–1.15
Cortical CSA ¹	mm ²	247.3 ± 52.7	316.6 ± 35.3	0.007	–1.55
Cortical thickness	mm	3.8 ± 0.6	4.3 ± 0.6	0.9	–0.88
Mass and density					
Total BMC ²	g	2.9 ± 0.6	3.5 ± 0.4	0.034	–1.17
Trabecular BMD ³	mg/cm ³	229.5 ± 24.1	243.6 ± 31.3	0.2	–0.51
Cortical BMD ³	mg/cm ³	1109.8 ± 32.1	1099.8 ± 52.3	1.0	0.23
Strength					
xSSI ⁴	mm ³	1041.4 ± 299.8	1596.4 ± 239.6	<0.001	–2.05
ySSI ⁵	mm ³	771.7 ± 192.4	1084.9 ± 215.0	0.004	–1.54

Data are presented as the means with standard deviations. Effect size, standardised mean difference Cohen's D. Bold number denotes p-values < 0.05.

¹CSA, cross-sectional area.

²BMC, bone mineral content.

³BMD, bone mineral density.

⁴xSSI, strength-strain index in the lateral direction.

⁵ySSI, strength-strain index in the anterior-posterior direction.

development.²⁶ Most of the patients in our cohort were not hypoxic (<90%) at the time of the peripheral quantitative CT examination, and no associations were found between oxygen saturation and strength-strain index. However, age at total cavo-pulmonary

connection surgery, which reflects time spent in hypoxia, was associated with a lower strength-strain index. Hypoxia during a sensitive period for bone development could possibly explain the association. Nevertheless, this association did not remain in the

Table 2. Univariate and multivariable linear regression analyses of factors associated with strength-strain index

Variable	Unit	Univariate				Multivariable		
		n	B	95% CI	p-value	B	95% CI	p-value
Height	cm	20	22.1	18.5–25.7	<0.001	18.2	12.3–24.0	<0.001
Age at TCPC ¹	months	19	–26.9	–8.9– –55.8	0.006	1.3	–11.8–14.5	0.83
Weight	kg	20	20.9	17.2–24.5	<0.001			
Oxygen saturation	%	20	28.1	–55.4–111.6	0.5			
Serum 25 (OH) D ²	nmol/L	20	0.69	–15.1–16.4	0.9			
GGT ³	μkat/L	20	87.0	–321.1–495.2	0.7			
pro-BNP ⁴	ng/L	20	–0.04	–0.7–0.7	0.9			
Growth pain legs	yes	19	–74.4	–409.4–260.7	0.7			

In the multivariable mode $r = 0.9$. Number in model $n = 19$. Bold numbers denote p -values < 0.05 .

¹TCPC, total cavo-pulmonary connection.

²25(OH) D, 25-hydroxy-vitamin D.

³GGT, gamma-glutamyltransferase.

⁴pro-BNP, B-type natriuretic protein.

multivariable regression model. However, the analysis reflects a small sample, which did not allow for extensive multiple regression models. Therefore, it is suggested that future prospective study protocols include hypoxia as a variable, preferably in a larger population. It would also be of interest to investigate if this association is present in other cyanotic heart diseases.

It is known that physical activity has an important effect on bone development.²¹ Therefore, the findings of a lower strength-strain index and bone mineral content in the tibia are of particular interest considering that the tibia is a weight-bearing bone, and hence constantly exposed to stress during daily activities.²¹ The lower self-reported physical activity among the patients compared to the controls could potentially explain the bone deficits. The lower physical activity level is mostly explained by the lower participation in free-time sports. However, these data must be interpreted with caution since they are self-reported activities. The findings of a lower self-reported participation in free-time sports are in line with a previous study by Hedlund et al.²⁸ Nevertheless, when objectively measuring the time spent at moderate physical activity in the same cohort, these previous authors found no difference between the patients with Fontan circulation and controls. The effect of physical activity on bone development in patients with Fontan circulation needs further investigation.

An interesting observation is that the adolescents with Fontan circulation had a bone mineral content level comparable to the children (6–12 years) of the control group (2.9 g versus 2.8 g). It is known that the growth in infancy is associated with bone mineral content levels later in life.²⁷ Therefore, a possible explanation for the lower bone mineral content found in the adolescents with Fontan circulation could be a decrease in growth trajectory during the first years of life, especially prior to total cavo-pulmonary connection surgery.²⁹ Unfortunately, data on growth trajectory were not available in the present population.

Contrary to our findings of a lower tibia strength-strain index and bone mineral content, Sarafoglou et al. found no differences in the tibia of children with Fontan circulation.⁸ A possible explanation for the conflicting findings could be that the lower strength-strain index and bone mineral content in our study were mostly driven by the adolescent patients. Even though the mean age between our study and the study by Sarafoglou et al. appears similar (13.0 versus 12.1 years), there are indications of a younger population in their

study since the majority of the patients (7 out of 10) had a Tanner stage ≤ 2 . This further stresses the need for future prospective studies investigating whether these differences in bone quality arise during adolescence.

Unfortunately, the cross-sectional design of present study does not allow analyses of causal relationships. Yet potential causes could be the low cardiac output and increased central venous pressure or mild residual hypoxia that the Fontan circulation entails.³⁰ In addition, many other organs are affected by a single ventricle physiology, resulting in secondary effects on e.g., bone mineral content.³¹ However, previous cross-sectional studies have shown presence of reduced bone in other CHDs as well,⁵ thus a Fontan circulation is probably not the only explanation. To address this topic, larger prospective studies including multiple CHDs are desirable.

Limitations

Since the cross-sectional study design only provides a momentary description of the population, it is not possible to analyse the individual development of bone health over time. Although the results of the present study might suggest that strength-strain index and bone mineral content, on a group level, decreases with age, prospective studies are needed to confirm these findings. Another limitation is the rather small study population, which does not allow for extensive analyses using multiple regression models or linear regression analyses in the subgroups. Nevertheless, the number of participants was adequate to enable subgroup analyses. In addition, age- and sex-matched controls were included, which can be seen as a strength of the present study. This is especially important since large national reference values for peripheral quantitative CT are not available for children. A known limitation with existing reference data for children is that site of measurements, age ranges, and groups of subjects may differ.²⁰ An additional limitation is the lack of information regarding Tanner stage since this could have enabled the comparison of maturity between patients and controls.

Conclusion

Young patients (6–19 years) with Fontan circulation have a lower strength-strain index compared to controls. Subgroup analyses

show that this deficit is mainly driven by the differences in adolescents (13–18), which might suggest that impaired bone strength develops with age. Our findings highlight the need to monitor bone development in patients with Fontan circulation in order to detect further deterioration of bone quality, and if possible, to prevent future fractures.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S1047951124000404>.

Acknowledgements. We would like to thank all the patients who generously committed time and effort to the current study. We would also like to thank all the personnel at Livsmedicin, Umeå, for the substantial cooperation regarding examinations with peripheral quantitative CT.

Author contribution. Annika Rydberg and Camilla Sandberg contributed equally to last authorship.

Financial support. This work was supported by the Swedish Heart-Lung Foundation (20160496) and the Swedish Odd Fellow order.

Competing interests. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the Helsinki Declaration of 1975, as revised in 2008, and have been approved by the Regional Ethics Review Board, Umeå (Dnr: 2016-445-31M, 2018-07-32M).

References

- Mandalenakis Z, Giang KW, Eriksson P, et al. Survival in children with congenital heart disease: have we reached a peak at 97%? *J Am Heart Assoc* 2020; 9: e017704.
- Moons P, Bovijn L, Budts W, Belmans A, Gewillig M. Temporal trends in survival to adulthood among patients born with congenital heart disease from 1970 to 1992 in Belgium. *Circulation* 2010; 122: 2264–2272.
- Ritmeester E, Veger VA, van der Ven JPG, et al. Fontan circulation associated organ Abnormalities beyond the heart, lungs, liver, and gut: a systematic review. *Front Cardiovasc Med* 2022; 9: 826096.
- Sandberg C, Johansson K, Christersson C, Hlebowicz J, Thilén U, Johansson B. Low bone mineral density in adults with complex congenital heart disease. *Int J Cardiol* 2020; 319: 62–66.
- Johansson K, Johansson B, Sandberg C. Reduced bone strength in adults with moderate or complex congenital heart disease. *Int J Cardiol Congenit Heart Dis* 2021; 6.
- D'Ambrosio P, Tran D, Verrall CE, et al. Prevalence and risk factors for low bone density in adults with a Fontan circulation. *Congenit Heart Dis* 2019; 14: 987–995.
- Diab SG, Godang K, Müller LO, et al. Progressive loss of bone mass in children with Fontan circulation. *Congenit Heart Dis* 2019; 14: 996–1004.
- Sarafoglou K, Petryk A, Mishra PE, et al. Early characteristics of bone deficits in children with Fontan palliation. *Cardiol Young* 2020; 30: 468–475.
- Avitabile CM, Goldberg DJ, Zemel BS, et al. Deficits in bone density and structure in children and young adults following Fontan palliation. *Bone* 2015; 77: 12–16.
- Vaikunth SS, Leonard MB, Whitehead KK, et al. Deficits in the functional muscle-bone unit in youths with Fontan physiology. *J Pediatr* 2021; 238: 202–207.
- Witzel C, Sreeram N, Coburger S, Schickendantz S, Brockmeier K, Schoenau E. Outcome of muscle and bone development in congenital heart disease. *Eur J Pediatr* 2006; 165: 168–174.
- Kanis JA, Cooper C, Rizzoli R, Reginster JY. European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int* 2019; 30: 3–44.
- Fonseca A, Gordon CL, Barr RD. Peripheral quantitative computed tomography (pQCT) to assess bone health in children, adolescents, and young adults: a review of normative data. *J Pediatr Hematol Oncol* 2013; 35: 581–589.
- Jiang H, Robinson DL, McDonald M, et al. Predicting experimentally-derived failure load at the distal radius using finite element modelling based on peripheral quantitative computed tomography cross-sections (pQCT-FE): a validation study. *Bone* 2019; 129: 115051.
- Samelson EJ, Broe KE, Xu H, et al. Cortical and trabecular bone microarchitecture as an independent predictor of incident fracture risk in older women and men in the bone microarchitecture international consortium (BoMIC): a prospective study. *Lancet Diabetes Endocrinol* 2019; 7: 34–43.
- Sandberg C, Frisk E, Hansson L, et al. Impaired knee extension muscle strength in adolescents but not in children with Fontan circulation. *Cardiol Young* 2020; 30: 1138–1143.
- Bergdahl MS, Crenshaw AG, Hedlund ER, Sjöberg G, Rydberg A, Sandberg C. Calf muscle oxygenation is impaired and may decline with age in young patients with total cavopulmonary connection. *Pediatr Cardiol* 2022; 43: 449–456.
- Hansson L, Sandberg C, Öhlund I, et al. Vitamin D, liver-related biomarkers, and distribution of fat and lean mass in young patients with Fontan circulation. *Cardiol Young* 2021; 32: 1–8.
- Kontulainen SA, Johnston JD, Liu D, Leung C, Oxland TR, McKay HA. Strength indices from pQCT imaging predict up to 85% of variance in bone failure properties at tibial epiphysis and diaphysis. *J Musculoskelet Neuronal Interact* 2008; 8: 401–409.
- Stagi S, Cavalli L, Cavalli T, de Martino M, Brandi ML. Peripheral quantitative computed tomography (pQCT) for the assessment of bone strength in most of bone affecting conditions in developmental age: a review. *Ital J Pediatr* 2016; 42: 88.
- Soyka LA, Fairfield WP, Klibanski A. Clinical review 117: hormonal determinants and disorders of peak bone mass in children. *J Clin Endocrinol Metab* 2000; 85: 3951–3963.
- Kröönström LA, Dellborg M, Giang KW, Eriksson P, Mandalenakis Z. Fractures in patients with and without congenital heart disease – a nationwide register-based cohort study. *Int J Cardiol Congenit Heart Dis* 2022; 10: 100413.
- Kröönström LA, Dellborg M, Cider Å, Eriksson P, Rosengren A, Mandalenakis Z. Fractures in children and young adults with and without congenital heart disease. *Int J Cardiol Congenit Heart Dis* 2021; 2: 100072.
- Avitabile CM, Leonard MB, Zemel BS, et al. Lean mass deficits, vitamin D status and exercise capacity in children and young adults after Fontan palliation. *Heart* 2014; 100: 1702–1707.
- Khazai N, Judd SE, Tangpricha V. Calcium and vitamin D: skeletal and extraskelatal health. *Curr Rheumatol Rep* 2008; 10: 110–117.
- Arnett TR, Gibbons DC, Utting JC, et al. Hypoxia is a major stimulator of osteoclast formation and bone resorption. *J Cell Physiol* 2003; 196: 2–8.
- Cooper C, Fall C, Egger P, Hobbs R, Eastell R, Barker D. Growth in infancy and bone mass in later life. *Ann Rheum Dis* 1997; 56: 17–21.
- Hedlund ER, Lundell B, Villard L, Sjöberg G. Reduced physical exercise and health-related quality of life after Fontan palliation. *Acta Paediatr* 2016; 105: 1322–1328.
- Daymont C, Neal A, Prosnitz A, Cohen MS. Growth in children with congenital heart disease. *Pediatrics* 2013; 131: e236–e242.
- Gewillig M, Brown SC. The fontan circulation after 45 years: update in physiology. *Heart* 2016; 102: 1081–1086.
- Rychik J. The relentless effects of the Fontan paradox. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu* 2016; 19: 37–43.