## **NOTES FOR AUTHORS**

The Journal of Helminthology publishes papers and reviews on all aspects of pure and applied helminthology, particularly those helminth parasites of environmental health, medical or veterinary importance. For those wishing to contribute research papers, work on helminths in wildlife hosts including plant and insect parasites will also be welcome. Taxonomic contributions will be acceptable if they contribute to the systematics of a group and particularly if they employ biochemical or molecular biological techniques.

**Page Format.** The *Journal* is printed in a two-column format (column width of 80 mm) with a text area of 170×225 mm.

**Text.** Papers should be typed, on one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used.

When the paper has been accepted word-processed text stored on floppy disc is encouraged, proving the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with fewer type-setting errors.

**Abstract.** Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words. A short title should be provided as a running head.

**Tables.** Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or landscape (if absolutely necessary) format. Table captions should be typed on a separate sheet.

Illustrations. Copies only of artwork should be submitted. The original illustrations should accompany the paper after acceptance and revision. Text figures, line drawings, computer-generated figures and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Halftone photographs are acceptable where they are a real contribution to the text. They should be glossy prints of the same size as they are to appear in the Journal. All figures and letters on photographs must be inserted by the author. Figure and captions should be typed on a separate sheet.

**Voucher specimens.** The deposition of voucher specimen should be considered where appropriate.

**References.** References must be based on the name and year system, give full journal titles and conform to the following styles:

Grønvold, J., Wolstrup, J., Larsen, M., Henriksen, S.A. & Nansen, P. (1993) Biological control of Ostertagia ostertagi by feeding selected nematode-trapping fungi to calves. Journal of Helminthology 67, 31–36.

**Grove**, **D.I.** (1990) A history of human helminthology. 850 pp. Wallingford, CAB International.

Southgate, V.R. & Rollinson, D. (1987) Natural history of transmission and schistosome interactions. pp. 347–378 in Rollinson, D. & Simpson, A.J.G. (Eds) The biology of schistosomes: from genes to latrines. London, Academic Press.

Citation of authors in the text should appear in the form: Polaszek (1990) or (Polaszek, 1990). More than one author should be cited in chronological order as: (Holloway et al., 1987; Walker & Huddleston, 1988).

**Offprints.** 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

**Manuscript.** Three copies of the manuscript, which must be in English or French (with an English summary) should be accompanied by a letter signed by all the authors and together with artwork submitted to:

The Editor
Journal of Helminthology
International Institute of Parasitology
395A Hatfield Road
St Albans, Herts
AL4 0XU, UK.

## Journal of Helminthology

Editorial	
Research Papers	
Akufongwe, P.F., Dakul, D.A., Michael, P.D., Dajagat, P.D. & Arabs, W.L. Urinary schistosomia	sis
in rural communities of some Local Government Areas in Plateau State, Nigeria:	
a preliminary parasitological and malacological survey	
Bæk, J., Simonsen, P.E., Friis, H. & Christensen, N.Ø. Zinc deficiency and host response to	
helminth infection: Echinostoma caproni infections in CBA mice	
Ferrer, J.R., Gracenea, M., Trullols, M., & Gonzalez-Moreno, O. Ultrastructural observations	
the tegument of Postorchigenes gymnesicus (Digenea: Lecithodendriidae)	
Jones, A., Bailey, T.A., Nothelfer, H.B., Gibbons, L.M., Samour, J.H., Al Bowardi, M. & Osborr	
Parasites of wild houbara bustards in the United Arab Emirates	
Kennedy, C.R. Colonization and establishment of <i>Pomphorhynchus laevis</i> (Acanthocephala) in a isolated English river	
Niwa, A. & Miyazato, T. Enhancement of intestinal eosinophilia during Hymenolepis nana infection in mice	
Nollen, P.M. The mating behaviour of Echinostoma paraensei grown in mice	
Omar, M.S. & Raoof, A.M.S. Onchocerca fasciata: histochemical demonstration of succinate and NADH dehydrogenase	
Patel, M.N. & Wright, D.J. The influence of neuroactive pesticides on the behaviour of entomopathogenic nematodes	
Richards, D.T., Hoole, D., Lewis, J.W., Ewens, E. & Arme, C. Adherence of carp leucocytes to adults and cercariae of the blood fluke Sanguinicola inermis	
Theodore, J.G. & Kaliraj, P. Wuchereria bancrofti recombinant antigen-derived poly-	
and monoclonal antibodies for the detection of circulating antigen(s) in the sera of lymphatic filarial patients	
Udonsi, J.K., Behnke, J.M. & Gilbert, F.S. Analysis of the prevalence of infection and associate between human gastrointestinal nematodes among different age classes living in the urban and suburban communities of Port Harcourt, Nigeria	ions
Veltkamp, C.J., Richards, G.R. & Chubb, J.C. Freeze fixation-dehydration as a method of	
preparation of Gyrodactylus (Monogenea) for SEM	
Research Notes	
Pillay, D. Electrophoretic differentiation of soluble egg antigens from Schistosoma mansoni isola	
Pillay, D. Electrophoretic differentiation of soluble egg antigens from Schistosoma mansoni isola using SDS-PAGE	
mehendazole and arprinocid on Echinostoma caproni in ICR mice	

## © CAB INTERNATIONAL, 1996

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.