





# First Red List of Ecosystems assessment of a tropical glacier ecosystem to diagnose the pathways towards imminent collapse

JOSÉ R. FERRER-PARIS<sup>\*1,2,3</sup> , LUIS D. LLAMBÍ<sup>4,5</sup>   
ALEJANDRA MELFO<sup>6</sup>  and DAVID A. KEITH<sup>1,3</sup> 

**Abstract** Tropical glaciers are rapidly disappearing, particularly in isolated mountain peaks below 5,000 m elevation. These glaciers are fundamental substrates for unique cryogenic ecosystems in high tropical environments where the ice, melting water and rocky substrate sustain microbiological communities and other meso- and macro-biota. This study uses the Red List of Ecosystems guidelines to diagnose the collapse of the tropical glacier ecosystem of the Cordillera de Mérida, Venezuela. We undertook the assessment with existing estimates of glacier ice extent, indirect historical estimates of ice mass balance and global mechanistic models of future ice mass balance. We complemented these with additional statistical analysis of trends and bioclimatic suitability modelling to calculate and predict rates of decline and relative severity of degradation in selected ecosystem indicators. The evidence suggests an extreme risk of collapse (Critically Endangered) because of a prolonged and acute reduction in ice extent and changes in climatic conditions that are leading to the complete loss of ice mass. The ice substrate has declined 90% in the last 20 years, and observed acceleration of the rate of decline suggests it will probably disappear within the next 5 years. Loss of ice substrate will trigger an immediate loss of supraglacial, englacial and subglacial biotic compartments and initiate a decades-long succession of forefield vegetation. However, ongoing inventories of native biota and monitoring of ecosystem transitions can provide valuable insights and lessons for other ecosystems facing similar risks. The Red List of Ecosystems assessment protocol provides a useful framework for comparative analysis of cryogenic ecosystems.

**Keywords** Climate change, cryogenic biome, forefield biota, glacial biota, ice mass balance, Red List of Ecosystems, risk assessment, tropical Andes

\*Corresponding author, [j.ferrer@unsw.edu.au](mailto:j.ferrer@unsw.edu.au)

<sup>1</sup>Centre for Ecosystem Science, University of New South Wales, Sydney, Australia

<sup>2</sup>UNSW Data Science Hub, University of New South Wales, Sydney, Australia

<sup>3</sup>IUCN Commission on Ecosystem Management, Gland, Switzerland

<sup>4</sup>Instituto de Ciencias Ambientales y Ecológicas, Universidad de Los Andes, Mérida, Venezuela

<sup>5</sup>Consorcio para el Desarrollo Sostenible de la Ecoregión Andina (CONDESAN), Quito, Ecuador

<sup>6</sup>Centro de Física Fundamental, Departamento de Física, Universidad de Los Andes, Mérida, Venezuela

Received 8 June 2023. Revision requested 18 September 2023.

Accepted 13 November 2023. First published online 10 September 2024.

The supplementary material for this article is available at [doi.org/10.32942/X2VK54](https://doi.org/10.32942/X2VK54)

## Introduction

Cryogenic ecosystems by definition depend on the dynamics of snow and ice. They are undergoing widespread and intense transformations because of climate change, especially in the tropics (Huss et al., 2017; Masiokas et al., 2020). Tropical glacier ecosystems are restricted to the highest elevations of mountain ranges (typically above 4,700 m) where snow and ice have accumulated over years, and the ice, melting water and rocky substrates sustain microbiological communities and other meso- and macro-biota (Hotaling et al., 2017). They are naturally isolated and increasingly exposed to rising temperatures and changing precipitation patterns that create an imbalance in ice mass, with more ice lost because of melting and ablation than the amount gained by snowfall and rainfall (Sagredo & Lowell, 2012; Rabatel et al., 2013; Veettil & Kamp, 2019). Historical losses of glacial ice mass have been documented in several countries, with > 60% having been lost in Colombia and > 50% in Peru over the last 60 years (Masiokas et al., 2020). The tropical Andes are considered a global hotspot for cryogenic change (Vuille et al., 2018), having lost over 1 gigatonne of ice during 2000–2018 (Dussallant et al., 2019), and the whole tropics region is expected to lose at least 60% of its glacier ice mass by 2100 if current global climate change trends are maintained (Rounce et al., 2023).

The functions of glacier ice as a water reservoir, a regulator of local temperature and a substrate for biotic processes are being disrupted by continuing losses of mass (Buytaert et al., 2017; Stibal et al., 2020). The changes in biota and ecosystem processes associated with glacial retreat will have cascading effects as novel ecosystems emerge in recently exposed areas (Llambí et al., 2021; Rosero et al., 2021; Anthelme et al., 2022) and changes in downstream aquatic habitats, biota and water quality affect human livelihoods in the surrounding landscape (Cauvy-Fraunié et al., 2016; Huss et al., 2017; Llambí et al., 2020). Long-term monitoring in the region indicates that plant diversity in the tropical Andes is particularly vulnerable given the high proportion of endemic species that are dependent on low temperatures

This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Downloaded from <https://doi.org/10.1017/S0030605323001771> on 24 May 2024 at 02:09:52, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>. Published by Cambridge University Press on behalf of The Author(s), 2024.

<https://doi.org/10.1017/S0030605323001771>

and have narrow thermal niches (Cuesta et al., 2020, 2023). Andean landscapes are being transformed, including via the contraction of puna and paramo and degradation/loss of high Andean wetlands (Cuesta et al., 2019).

Monitoring the status and trends of ecosystems is fundamental to informing biodiversity conservation and sustaining the benefits from nature (Bland et al., 2019). The IUCN Red List of Ecosystems provides a standard protocol for monitoring through a rigorous assessment of the risk of ecosystem collapse. Ecosystem collapse is defined as a large transformational change involving loss of biodiversity and/or major degradation and loss of ecosystem function. The risk status of an ecosystem is represented by the assigned Red List of Ecosystems category of risk that summarizes the probability of collapse based on the diagnosis of qualitative and quantitative indicators (Keith et al., 2013).

Previous Red List of Ecosystems assessments in the tropical Andes have focused on vegetated ecosystem types including tropical alpine vegetation, but have not considered glacial ecosystems (Oliveira-Miranda et al., 2010; Etter et al., 2018). Globally there are few examples of such assessments of cryogenic ecosystems, and those that exist are primarily focused on seasonal snow (Williams et al., 2015; Kontula & Raunio, 2019), freeze-thaw glacial lakes or permanent snowfields (Murray et al., 2020).

Here we contribute to filling this gap by presenting a precursory example of a global assessment of all tropical glacier ecosystems. We focus on the glaciers of the Cordillera de Mérida in Venezuela, which have a documented history of decline in number and extent (Ramírez et al., 2020). We use information from local, regional and global studies to apply the Red List of Ecosystems criteria, and we provide a comprehensive ecosystem description supported by knowledge from related ecosystem types and information from local studies, a diagnosis of ecosystem functioning, threats and state of collapse and the ecosystem risk status based on spatial and functional symptoms of collapse.

## Study area

The tropical glacier ecosystem of the Cordillera de Mérida is a distinct ecosystem type that formerly occupied several peaks above 4,600 m in Sierra Nevada National Park in Venezuela (Fig. 1; Braun & Bezada, 2013; Ramírez et al., 2020).

Following the IUCN Global Ecosystem Typology, all tropical glacier ecosystems belong to the Ecosystem Functional Group T6.1 Ice sheets, glaciers and perennial snowfields in the Cryogenic functional biome (T6; Keith et al., 2022). Tropical glaciers share key features with other glaciers, such as the icy substrate, atmospheric deposition of




FIG. 1 The study area in the Cordillera de Mérida, Venezuela. This assessment focuses on three glaciated peaks: Bolívar, La Concha and Humboldt. These peaks and other historically glaciated areas are encompassed in a single 10 × 10 km cell (highlighted square, area of occupancy = 1). The isolated Mucuñuque Peak at 4,609 m lost its ice before 1930.

nutrients, a biota dominated by cold-adapted microorganisms that inhabit several habitat compartments on (supraglacial), in (englacial) or below (subglacial) the icy substrate, truncated trophic networks and low productivity and diversity (Anesio et al., 2017; Hotaling et al., 2017; Keith et al., 2022). Tropical glaciers differ from others in their exposure to daily variations in temperature that far exceed the differences in monthly mean temperatures across the year. Precipitation can occur throughout the year as rainfall or snowfall, although it tends to concentrate during the rainy or wet season, which typically extends for > 8 months (Sagredo & Lowell, 2012). Annual precipitation at the highest elevations in the Cordillera de Mérida is estimated to be 1,000–1,200 mm, with 10–17% of the total falling during December–March, and the mean annual temperature is  $-0.4$  °C at the highest station where records are available in the country (4,766 m), with a ratio of diurnal variation to annual variation of 2.15–2.36 (Pulwarty et al., 1998; Andressen, 2007).

The long-term mass balance of the icy substrate of tropical glaciers is therefore probably dominated by interannual fluctuations in precipitation and exposure to solar radiation, but this process can be accelerated by scale and edge effects (Ceballos et al., 2006; Andressen, 2007; Vuille et al., 2018). The Cordillera de Mérida has lost several glaciated areas in the last 150 years (Braun & Bezada, 2013), and the glacier retreat rates increased after 1998, exceeding 7% area loss per year, which is higher than the rates reported for the last 30 years in other glaciers of the tropical Andes (Ramírez et al., 2020).

Atmospheric or aeolian deposition (windfall) provide key nutrients such as carbon, nitrate and ammonium to the micro- and mesobiota of the supraglacial zone (Edwards, 1987; Hotaling et al., 2017), but this is also a source of light-absorbing particles that increase melting processes (Gilardoni et al., 2022). Nutrients and meltwater can be transported through interglacial cracks and crevasses to

reach the subglacial zone (contact zone between rock and ice substrate), where they combine with small particles produced by rock comminution (Hotaling et al., 2017). Greater concentrations of black carbon (sub-micron particle volume increasing from 0.19 to  $1.4 \mu\text{m}^3/\text{cm}^3$ ) at high elevations of the Cordillera de Mérida have been linked to biomass burning in Venezuelan savannah, with greater fire activity and higher concentrations observed following El Niño years (Hamburger et al., 2013).

The ice substrate, proglacial waters and glacier forefield share part of their microbiota because of dispersal through melting water (Hotaling et al., 2017) and are here considered compartments of the same ecosystem. Glacier retreat associated with climate change and subsequent soil development triggers multitrophic changes from simple food webs dominated by microbes and heterotrophic organisms to more complex networks as deglaciated areas are colonized by lichen, mosses and vascular plants, marking the transition to a novel ecosystem (Ficetola et al., 2021; Khedim et al., 2021; Llambí et al., 2021; Rosero et al., 2021; Anthelme et al., 2022).

Prospective samples from englacial and subglacial microbiota were collected from Bolívar and Humboldt peaks (Fig. 1, Plates 1 & 2; Ball et al., 2014; Balcazar et al., 2015; Rondón et al., 2016). These include samples taken prior to the complete disappearance of ice from Bolívar peak, but they do not allow for a full assessment of the role of the microbiota in the ecosystem. However, a more detailed study of the microbiota in the glacier forefield at Humboldt peak is underway (B. Huber, pers. comm., 2023). In the forefield area, a diverse array of pioneer lichen and bryophyte species (including many new records for Venezuela and several endemic species) appears to facilitate the establishment of pioneer wind-dispersed and wind-pollinated vascular plants during the first 6 decades of primary succession after glacier ice retreat (Llambí et al., 2021).



PLATE 1 View of the glacier at Humboldt Peak in the Cordillera de Mérida, Venezuela, in August 2022. The photograph was taken from the peak, looking west towards the direction of (1) Bolívar Peak and (2) La Concha Peak. Photo: J.A. González.




PLATE 2 Mosaic of satellite images (Sentinel-2 MultiSpectral Instrument, Level 2A; December 2021–March 2022; clouds and shadows removed) over the three peaks in the Cordillera de Mérida, Venezuela: (1) Bolívar, (2) La Concha and (3) Humboldt. The square represents the same 10 × 10 km grid cell shown in Fig. 1.

## Methods

We applied the IUCN Red List of Ecosystems protocol (Bland et al., 2017) that requires assessment of five criteria: A, declining distribution; B, restricted distribution and exposure to threats; C, degradation of abiotic environment; D, disruption of biotic processes and E, quantitative risk analysis. Three of the five criteria (A, C and D) include sub-criteria applied to three different timeframes to capture the effects of historical, current and future threats. Criterion B evaluates the current exposure to threats using various spatial metrics, and criterion E is evaluated over 50 and 100 years into the future. The complete assessment is provided in the Supplementary Material.

We captured the main ecosystem processes and the threats of climate change and air pollution in a conceptual ecosystem model (Fig. 2) and identified spatiotemporal indicators for the assessment of each criterion and sub-criterion. We defined a threshold of collapse for each indicator and interpolated or extrapolated time series data to infer rates of change over the relevant timeframes.

We defined the collapsed state of the ecosystem as the complete loss of the icy substrate that can sustain a cold-resistant microbiota (zero ice extent for spatial criteria A and B or zero ice mass for criterion E). We used three indicators to assess climatic suitability for glacial function under criterion C: the equilibrium-line altitude (the altitude at which rates of ice accumulation and ablation are equal); the atmospheric freezing level height (the altitude of the 0 °C isotherm); and a suitability threshold based on a correlative bioclimatic suitability model. An increase in equilibrium-line altitude or freezing level height reduces the available area for long-term glacier persistence, with

thresholds of collapse for both indicators set to the altitude of the mountain summit (Polissar et al., 2006; Braun & Bezada, 2013). We based the threshold of bioclimatic suitability on the optimal classification of the current occurrence of glacier outlines. There is currently no suitable indicator of collapse for criterion D because the microbiota of the supraglacial zone and its relationships with other ecological zones of the glacier are still poorly understood (Ball et al., 2014; Rondón et al., 2016). Data collected at Humboldt Peak in 2019 and 2021 (including supraglacial, englacial and subglacial samples as well as soil from the glacial forefront) may shed light on this issue. Preliminary results indicate marked changes in microbiota composition and function and a sizeable role of the microbiota on ecosystem processes in four chronosequence sites deglaciated during 1910–2009 (B. Huber, pers. comm., 2023); similar findings have been reported for a tropical glacier in Ecuador (Díaz et al., 2023).

We used available measures of glacier ice extent to calculate best estimates of past decline and to extrapolate rates of future declines in ice extent (sub-criteria A1, A2b and A3). We obtained measurements and their standard errors from published, ground-validated reconstructions based on cartographic data from 1910, aerial photographs from 1952 and 1998, satellite images from 2009, 2015 and 2016 and field measurements using GPS and drone imagery from 2019 (Ramírez et al., 2020). Indirect and less accurate evidence of the extent of glacier ice prior to 1910 is available from a few other sources and is provided for context (Polissar et al., 2006; Jomelli et al., 2009; Braun & Bezada, 2013). We calculated past declines based on pairs of measurements and considering error propagation. For future declines we calculated first proportional rates of decline (as percentage area loss per year) and then used this rate to project the expected magnitude of decline




FIG. 2 Conceptual ecosystem model for the tropical glacier ecosystem of the Cordillera de Mérida, Venezuela. White ovals and boxes: those with a dashed outline represent the main characteristic biota and biotic processes, those with a solid outline the main elements of the abiotic environment and abiotic processes, respectively. Dark grey boxes: threatening processes. Black lines: relationships between elements (arrow: increases; circle: reduces). Light grey boxes: indicator variables used in the assessment and indirectly linked to ecosystem processes or components by grey dashed lines. Some elements and processes are excluded for clarity of visualization.

over a 50-year period (1948–1998). Given the evidence suggesting accelerating rates of decline, we compared the results of using different timeframes for calculation of proportional rates of decline.

We used two spatial measures of exposure to threats (criterion B): the extent of occurrence was calculated as the area of the convex hull around the glacier outlines from the RGI 6.0 database (Randolph Glacier Inventory Consortium, 2017), and the area of occupancy was calculated as the minimum number of  $10 \times 10$  km cells that include all of these occurrences.

For criterion C<sub>1</sub> we used a time series of freezing level height in metres calculated from climate reanalysis data for 1948–2011 (Braun & Bezada, 2013) and fitted a local polynomial regression (loess with gaussian distribution, span = 0.75 and degree = 2, equivalent number of parameters = 4.35) to smooth the temporal trend. We calculated relative severity of degradation (RS) using the formula:  $RS = OD/MD$ , where OD (observed decline) is the difference between the initial and final values and MD (maximum decline) is the difference between the initial value and the collapse threshold (Keith et al., 2013). We used the initial and final values of the smoothed trend in freezing level height and two collapse thresholds of 4,920 and 4,970 m, which span the range of plausible estimates. We took estimates of historical changes in equilibrium-line altitude from Polissar et al. (2006), and these are based on reconstructed palaeo-glacier topography near Mucuñuque Peak (located in nearby Sierra de Santo Domingo; Fig. 1) and cumulative elevation profiles of modern

glaciers at Bolívar Peak. For the calculation of relative severity of degradation we used change in equilibrium-line altitude as a direct measurement of observed decline and the maximum possible change as maximum decline (sub-criterion C<sub>3</sub>).

We also used local results of correlative and mechanistic global models to project trends in climatic suitability (sub-criterion C<sub>2a</sub>) and the probability of persistence of tropical glaciers into the future (criterion E). These models incorporated different global circulation models and scenarios of projected greenhouse gas emissions based on alternative socio-economic developments and climate policies (shared socio-economic pathways; Karger et al., 2017).

The correlative model of environmental suitability focused exclusively on tropical regions and compares the bioclimatic conditions of high-elevation areas with and without glaciers using the Gradient Boosting Machine algorithm (Ferrer-Paris & Keith, 2024). We selected occurrence records using stratified random sampling from glacier outlines in tropical areas (from the tropical Andes, Mexico, Africa and Asia) and non-glaciated areas with a minimum of 3,500 m elevation and a maximum distance of 25 km from the nearest glacier outlines. We withheld occurrence records from the Cordillera de Mérida for final evaluation of prediction performance of the model; we divided all other records into calibration (80%) and test partitions (20%) and used them for tuning of the model parameters (number of trees, interaction depth, shrinkage and minimum number of observations per node) and final model fitting using

cross-validation. We compared the predicted probability or suitability index with the known occurrences in the test partition and in the Cordillera de Mérida to calculate cut-off values of equal sensitivity and specificity (balancing errors of omission and commission) and maximum accuracy (minimizing misclassification errors). We then used the model to predict future suitability under 15 combinations of global circulation models and socio-economic pathways, assuming the alternative cut-off values represented thresholds of ecosystem collapse (sub-criterion C2a).

The quantitative analysis of collapse for criterion E was based on a global hybrid model (Rounce et al., 2023) that combined a mass balance module and a glacier dynamics module to model all glaciers in the world independently using globally available datasets of glacier outlines (Randolph Glacier Inventory Consortium, 2017), glacier-wide geodetic mass balance data and regional ice volume estimates for calibration (Farinotti et al., 2019; Hugonnet et al., 2021). In the case of small regions with no direct measurements (such as the Cordillera de Mérida) the model used initial estimates of ice volume based on digital elevation models and most likely overestimated initial mass, making the projections conservative (Rounce et al., 2023). The model was run with monthly and annual time steps from 2000 to 2100 for various ensembles of global circulation models and socio-economic pathways. We used the ice mass projections of the model for the glacier outlines of the Cordillera de Mérida (Rounce et al., 2022) to estimate the year of collapse (first year when ice mass reaches zero) for 48 combinations of global circulation models and socio-economic pathways. We used the empirical cumulative distribution function of the year of collapse to calculate the proportion of models indicating collapse for each year and estimate probability of collapse.

We performed all statistical analysis using *R 4.3.1* (R Core Team, 2023). For the suitability model we used functions in the *R* package *caret* (Kuhn, 2008).

## Results

There are no direct field measurements of decline in extent of glacier ice for the last 50 years, but cartographic estimates of glacier extent suggest a decline of at least 89% over

21 years (proportional rate of decline =  $5.173 \pm \text{SE } 0.130\%$ /year) and nearly 99% if we consider the last 67 years (proportional rate of decline =  $10.220 \pm \text{SE } 0.473\%$ /year). These rates of decline would translate to future declines of at least 94% by the year 2048 (Table 1).

Given the rapid decline and disappearance of the glacier at Bolívar Peak during 1998–2017 and that the size of the Humboldt glacier at the time of the last measurement ( $0.045 \text{ km}^2$  in 2019) was comparable to that of the Bolívar glacier in 1998, it is reasonable to expect that the Humboldt glacier will disappear completely over the next 20 years. The area of glacier ice in Humboldt Peak was last measured at  $0.046 \pm 0.004 \text{ km}^2$  in July 2019 (Ramírez et al., 2020) and was estimated to be c.  $0.01 \text{ km}^2$  in August 2022 (Plate 1; N. Ramírez, pers. comm., 2023). This last field observation would confirm the recent acceleration of the rate of decline of the Humboldt glacier and increases the likelihood that it will disappear sooner than predicted, probably within the next 5 years.

Reconstruction of the historical evolution of glaciers in South America suggests that the most recent maximum glacial extent in Venezuela occurred c. 1730 (during the Little Ice Age), and glaciers retreated continuously in the following centuries, with only minor readvances c. 1760, 1820 and 1880 (Polissar et al., 2006; Jomelli et al., 2009). Thus, we can assume that the observed decline between 1910 and the present represents a lower bound of the total decline between 1750 and the present (Table 1).

The extent of occurrence for the Humboldt glacier and surrounding areas is  $0.892 \text{ km}^2$ . If the proglacial waters and glacier forefields of recently collapsed glaciers are included, the extent of occurrence is at most  $5.957 \text{ km}^2$ . The main occurrences of known glaciers (extant and recently collapsed) occupy a single  $10 \times 10 \text{ km}$  cell (Fig. 1, Plate 2). This cell probably contains all occurrences of associated habitats with connected microbiota (proglacial waters and glacier forefields).

The relationship between recent temporal changes in climatic conditions and the disappearance of glacier ice is best illustrated with the case of La Concha Peak. The estimated maximum elevation of ice was 4,840 m in the year 1952, and it disappeared before 1998. The mean freezing level height was almost 10 m below the maximum elevation in 1948

TABLE 1 Estimated magnitude of past and future tropical glacier declines in the Cordillera de Mérida, Venezuela (Fig. 1), based on previous measurements (Ramírez et al., 2020) and estimated proportional rates of decline.

Start date	End date	Timeframe	Method	Decline $\pm$ SE (%)
1910	2019	109 years	Reconstructed map + field observations	$99.10 \pm 0.080$
1952	2019	67 years	Aerial photographs + field observations	$98.06 \pm 0.180$
1998	2019	21 years	Aerial photographs + field observations	$89.61 \pm 1.149$
1998	2048	50 years	Projected using proportional rate of decline = $5.71 \pm 0.130\%$	$94.72 \pm 0.611$
1998	2048	50 years	Projected using proportional rate of decline = $10.20 \pm 0.473\%$	$99.54 \pm 0.127$

and rose to more than 83 m above the maximum elevation in 2010, and the smoothed freezing level height mean surpassed the 4,840 m threshold during 1972–1981 (Fig. 3). Using the thresholds of maximum elevation of glacier ice in the two other peaks, we calculated the relative severity of change for this indicator to be  $67.2 \pm \text{SE } 18.5\%$  (Bolívar Peak) and  $104.4 \pm 28.5\%$  (Humboldt Peak). The large standard error is because of the wide variation in the time series.

The best-performing bioclimatic suitability model had high predictive performance during model training (area under the curve =  $0.967 \pm \text{SE } 0.008$ , sensitivity =  $0.621 \pm 0.058$  and specificity =  $0.981 \pm 0.004$ , with Gradient Boosting Machine model parameters: 200 trees, interaction depth = 5, shrinkage = 0.1 and a minimum of 12 observations per node) and when evaluating model predictions under current climatic conditions in the Cordillera de Mérida (area under the curve = 0.990; sensitivity = 0.667 and specificity = 0.996; full model details in Supplementary Material). The best estimate of mean relative severity for this indicator across all combinations of five global circulation models, three pathways, two timeframes and two collapse thresholds is 97%, with a 90% confidence interval of 63–100%.

There are three plausible values of difference in equilibrium-line altitude: the historical reconstruction of equilibrium-line altitude between the maximum glacier extent (pre-1820) and 1972 suggests a value of  $-300$  m, whereas comparison with the historically and recently collapsed Mucuñuque and Bolívar Peaks suggests values between  $-400$  and  $-500$  m (Polissar et al., 2006). Using a value of  $-550$  m as the maximum decline, we calculated




FIG. 3 Local polynomial regression of freeze level height in the Cordillera de Mérida, Venezuela, for the period 1948–2011, with 95% confidence intervals (based on data from Braun & Bezada, 2013). Horizontal lines represent the last recorded elevation of glacier ice at the different peaks, with years given in parentheses.

the relative severity of degradation for this indicator and timeframe to be 54–90%, with a median value of 72%.

The hybrid model predicted a high probability of collapse in the future. Focusing on the 50-year period 2020–2070, we found that 79% of the models end in collapse (Fig. 4). Uncertainty in mass estimates (mean absolute deviation) does not have a significant effect on the estimated year of collapse. Considering each scenario separately, the proportion of models that predict collapse by 2070 is higher than 50% in all cases, except for the sustainable development scenario (socio-economic pathway 1-2.6).

These quantitative results support the assessment of the risk of collapse using the IUCN Red List of Ecosystems categories and criteria (Table 2).

## Discussion

Based on the analyses presented here, the current Red List status of Cordillera de Mérida glacier is Critically Endangered, with plausible evidence that it may already be Collapsed: CR (CR-CO). Almost all of the criteria and sub-criteria assessed indicate the CR status, implying similar magnitudes of historical, present and future threats (Table 2). For the Cordillera de Mérida the evidence from recent field visits suggests that its glacier ice will probably disappear before 2048 (sub-criterion A2b). Other lines of evidence suggest similar conclusions, although with more




FIG. 4 Empirical cumulative distribution function of year of collapse from a global hybrid model of mass balance and glacier dynamics based on future climate predictions from 48 combinations of global circulation models and shared socio-economic pathways (SSPs; thick solid line) and disaggregated by the four different pathway (labelled lines): SSP1-2.6 is a sustainable development scenario, SSP2-4.5 is intermediate, SSP3-7.0 prioritizes national development and SSP5-8.5 is fossil-fuelled development.

TABLE 2 Summary of the IUCN Red List of Ecosystems assessment of the tropical glacier ecosystem of the Cordillera de Mérida, Venezuela. The first column indicates the sub-criteria assessed under each criterion. A more detailed report is available in the Supplementary Material. The categories of ecosystem risk are: Collapsed (CO), Critically Endangered (CR), Endangered (EN), Vulnerable (VU), Near Threatened (NT), Least Concern (LC) and Data Deficient (DD). A threat defined location is ‘a geographically or ecologically distinct area in which a single threatening event can rapidly affect all occurrences of an ecosystem type’ (Bland et al., 2017).

Indicator data & analysis applied	Estimates & uncertainty	Rationale	Category of risk
<b>Criterion A</b>			
A1 Cartographic estimates of glacier extent for three peaks (La Concha, Bolívar & Humboldt) in the Cordillera de Mérida (Ramírez et al., 2020)	Plausible bounds of decline in extent: 89–98%	There is no direct estimate of decline in extent for the last 50 years, but calculations of rates of decline between 1952–2019 & 1998–2019 used as minimum & maximum bounds	CR
A2b Data from Ramírez et al. (2020) used to estimate proportional rates of decline for the last remaining patch (Humboldt) & comparison with recently collapsed patch (Bolívar)	Best estimates of decline based on alternative proportional rate of decline estimates are > 94%. Assuming a similar trajectory between both patches, it is expected to reach 100% decline within the next 20 years	Recent acceleration of the rate of decline suggests that the higher proportional rate of decline is more likely; this is consistent with the observed timeframe of collapse in Bolívar Peak (Ramírez et al., 2020)	CR (CR-CO)
A3 Cartographic estimates of glacier extent for three peaks (La Concha, Bolívar & Humboldt) in the Cordillera de Mérida (Ramírez et al., 2020)	Best estimate of decline in extent: 99 ± SE 0.1%	Reconstruction of the historical evolution of glaciers suggests that the maximum glacial extent during the Little Ice Age occurred c. 1730 (Jomelli et al., 2009; Rabatel et al., 2013). The observed decline during 1910–2019 represents a lower bound of the total decline	CR (CR-CO)
<b>Criterion B</b>			
B1 Data from Ramírez et al. (2020), Braun & Bezada (2013) & RGI 6.0 database (Randolph Glacier Inventory Consortium, 2017)	Extent of occurrence < 20,000 km <sup>2</sup>	Restricted to one threat defined location, with evidence of continuing decline & inferred threatening processes	CR
B2 Same as B1	All occurrences (extant & collapsed) occupy one 10 × 10 km cell	Restricted to one threat defined location, with evidence of continuing decline & inferred threatening processes	CR
<b>Criterion C</b>			
C1 Freezing level height (in metres) from climate reanalysis data for the period 1948–2011 (Braun & Bezada, 2013), with a local polynomial regression	Averaged relative severity of degradation considering two collapse thresholds: 84% (plausible bounds 48–100%)	We used the values of the smoothed freezing level height to calculate initial & final values & assumed that the collapse value is 4,920–4,970 m	CR (VU-CO)
C2a Estimated change in suitability of bioclimatic conditions using a Gradient Boosting Machine model fitted to current climate (1980–2010) & projected to future timeframes (2010–2040 & 2040–2070) considering uncertainty	Best estimate of mean relative severity: 97% (95% CI 63–100%)	Predicted suitability represents the initial & final values; different cut-off values represent potential collapse thresholds. We considered uncertainty in climate models, scenarios & cut-off values	CR (EN-CO)
C3 Equilibrium-line altitude estimated from reconstructed palaeo-glacier topography & elevation profiles of existing glaciers (Polissar et al., 2006)	Best estimate of relative severity of degradation considering three probable values: 72% (plausible bounds 54–90%)	Change in equilibrium-line altitude was calculated for a historical timeframe (1820–1972) & an extended timeframe (1820–2006). This last value is considered to be near to the collapse threshold	EN (VU-CR)
<b>Criterion D</b>			
We considered prospective samples of the supraglacial microbiota & postglacial chronosequence of the glacier forefield (Ball et al., 2014; Rondón et al., 2016; Llambí et al., 2021)	No measure of relative severity could be calculated from the available data	The biota of this assessment unit is poorly known; there is no direct information on temporal changes in microbial communities in the different habitats after the loss of the ice substrate, but studies investigating this are underway	DD



TABLE 2 (Cont.)

Indicator data & analysis applied	Estimates & uncertainty	Rationale	Category of risk
<b>Criterion E</b>			
Ice mass balance projections from a global glacier evolution model (Rounce et al., 2022). We calculated the cumulative distribution of predicted year of collapse (first year when ice mass reaches zero) for 48 combinations of models & scenarios	Probability of collapse during 2020–2070: 79%	Uncertainty in mass estimates from replicates does not have a significant effect on the estimated year of collapse. The proportion of models that predict collapse by 2070 is higher than 50% in three out of four scenarios	CR (EN-CR)

uncertainty because of the use of indirect indicators and global instead of local data. During the last 100 years, the freezing level height and equilibrium-line altitude have increased close to the point of collapse, and permanent ice has completely disappeared from all previously glaciated peaks except Humboldt (sub-criteria A1, A3 and C1). The current distribution of glaciated areas is much reduced and exposed (sub-criteria B1 and B2), is undergoing rapid decline at a rate similar to the recently disappeared Bolívar glacier and is unlikely to recover mass before its complete collapse (sub-criterion C2a and criterion E).

This is the first Red List of Ecosystems assessment of a tropical glacier ecosystem. Cryogenic ecosystems are often under-represented in national assessments that focus on vegetation units, but despite differences in processes and functions they can be assessed using the same criteria and assigned to comparable categories of risk (Keith et al., 2013). Risk assessments and long-term monitoring of all ecosystems in tropical mountains can provide valuable information for comprehensive conservation planning and ecosystem management in these complex landscapes (Cuesta et al., 2019; Llambí et al., 2020).

This quantitative assessment of several symptoms of collapse illustrates how the Red List of Ecosystems protocol can be applied to other tropical glaciers (and cryogenic ecosystems outside tropical areas) by combining all available evidence within a common framework. We anticipate that similarly isolated and exposed tropical glaciers in Indonesia, East Africa, Mexico and some Andean regions (e.g. Santa Isabel in Colombia, Carihuairazo in Ecuador) will also be at extreme risk of collapse, and the larger regional units in the tropical Andes may experience widespread decline and degradation (Ceballos et al., 2006; Rabatel et al., 2018; Veettil & Kamp, 2019; Ferrer-Paris & Keith, 2024). The Red List of Ecosystems protocol provides a valuable framework for consistent and comparable assessments across all continents that synthesizes local and regional data from geology, climatology, palaeontology/palynology, microbiology, entomology and soil and vegetation ecology to describe ecosystems and diagnose the threats that they face.

The Red List of Ecosystems assessment explores different pathways towards collapse, and the associated uncertainty of methods and projections is consolidated within the categories of risk and their plausible bounds. Here we fully exploited this versatility, assessing spatial and functional symptoms, undertaking quantitative risk analysis and considering alternative interpretations of collapse thresholds. Our study shows that the multiple lines of evidence offer a broad picture of threatening processes and pathways towards collapse and that the assessment protocol handles several sources of uncertainty (Keith et al., 2013; Bland et al., 2017). Sufficient data are available for the comprehensive risk assessment of tropical glacier ecosystems in different regions of the world, and global datasets and analyses can be used as informative indicators of the climate and cryosphere dynamics that have not been studied locally, with the caveat of the limited ground validation and potential bias of these sources (Sagredo & Lowell, 2012; Rounce et al., 2023).

Because of the rapid decline of the icy substrate in recent decades, the small remnants of ice in Venezuela and on other tropical mountains are often considered static or extinct glaciers because dynamic processes of ice accumulation no longer operate and ice loss is accelerated by scale and edge effects (Ceballos et al., 2006; Braun & Bezada, 2013; Rabatel et al., 2018; Ramírez et al., 2020). Despite rapid contraction, the little remaining ice substrate could sustain much of its original microbiota (Ball et al., 2014; Balcazar et al., 2015; Rondón et al., 2016). Studies of the glacier forefield at Humboldt Peak provide insights into the postglacial chronosequence: pioneer lichens, mosses and vascular vegetation are already present 10 years after the retreat of the glacier. However, vascular plants only increase slowly in cover, species and functional diversity during the first 100 years of primary succession, and soil properties (e.g. soil organic matter, total nitrogen and exchangeable bases) change significantly after 21 and 60 years (Llambí et al., 2021).

Although collapse seems imminent and is probably unavoidable, we can still learn much from inventories of native biota and monitoring of ecosystem transitions. Multiple

changes in climate, soil and ecological communities operate at different temporal scales in high Andean summits and the recently deglaciated lands (Ficetola et al., 2021; Khedim et al., 2021; Llambí et al., 2021; Anthelme et al., 2022; Cuesta et al., 2023). A key research question is whether some biotic components of glacial ecosystems will be lost or whether they will persist in cold, ice-free refuges after glacial collapse (Stibal et al., 2020). However, we have limited time to study the remainder of this glacier microbiota in the novel ecosystems that form after the complete retreat of ice.

**Author contributions** Study design: JRF-P, DAK; data review: JRF-P, LDL, AM; data analysis: JRF-P; writing: all authors.

**Acknowledgements** We thank Nerio Ramírez for his invaluable work documenting glacier retreat and Barbara Huber for kindly sharing preliminary results on glacier microbiota in the Venezuelan Andes. The National Geographic Society provided financial support for LDL and AM (Grant No. NGS-55170R-19). We also thank the Mucumbarila cable car system and INPARQUES for their support in accessing the study sites near Humboldt Peak.

**Conflicts of interest** None.

**Ethical standards** This research abided by the *Oryx* guidelines on ethical standards.

**Data availability** The data and code that support the findings of this study are available at [osf.io/y3279](https://osf.io/y3279).

## References

- ANDRESEN, R. (2007) Circulación atmosférica y tipos de climas. In *GeoVenezuela 2. Medio Físico y Recursos Naturales* (ed. P.C. Grau), pp. 238–329. Fundación Polar, Caracas, Venezuela.
- ANESIO, A.M., LUTZ, S., CHRISMAS, N.A.M. & BENNING, L.G. (2017) The microbiome of glaciers and ice sheets. *npj Biofilms and Microbiomes*, 3, 10.
- ANTHELME, F., CARRASQUER, I., CEBALLOS, J.L. & PEYRE, G. (2022) Novel plant communities after glacial retreat in Colombia: (many) losses and (few) gains. *Alpine Botany*, 132, 211–222.
- BALCAZAR, W., RONDÓN, J., RENGIFO, M., BALL, M.M., MELFO, A., GÓMEZ, W. & YARZÁBAL, L.A. (2015) Bioprospecting glacial ice for plant growth promoting bacteria. *Microbiological Research*, 177, 1–7.
- BALL, M.M., GÓMEZ, W., MAGALLANES, X., ROSALES, R., MELFO, A. & YARZÁBAL, L.A. (2014) Bacteria recovered from a high-altitude, tropical glacier in Venezuelan Andes. *World Journal of Microbiology and Biotechnology*, 30, 931–941.
- BLAND, L., KEITH, D., MILLER, R., MURRAY, N. & RODRÍGUEZ, J.P. (2017) *Guidelines for the Application of IUCN Red List of Ecosystems Categories and Criteria. Version 1.1*. IUCN, Gland, Switzerland. [doi.org/10.2305/IUCN.CH.2016.RLE.3.en](https://doi.org/10.2305/IUCN.CH.2016.RLE.3.en).
- BLAND, L.M., NICHOLSON, E., MILLER, R.M., ANDRADE, A., CARRÉ, A., ETTER, A. et al. (2019) Impacts of the IUCN Red List of Ecosystems on conservation policy and practice. *Conservation Letters*, 12, e12666.
- BRAUN, C. & BEZADA, M. (2013) The history and disappearance of glaciers in Venezuela. *Journal of Latin American Geography*, 12, 85–124.
- BUYTAERT, W., MOULDS, S., ACOSTA, L., DE BIÈVRE, B., OLMOS, C., VILLACIS, M. et al. (2017) Glacial melt content of water use in the tropical Andes. *Environmental Research Letters*, 12, 114014.
- CAUVY-FRAUNIE, S., ANDINO, P., ESPINOSA, R., CALVEZ, R., JACOBSEN, D. & DANGLES, O. (2016) Ecological responses to experimental glacier-runoff reduction in alpine rivers. *Nature Communications*, 7, 12025.
- CEBALLOS, J.L., EUSCÁTEGUI, C., RAMÍREZ, J., CAÑON, M., HUGGEL, C., HAEBERLI, W. & MACHGUTH, H. (2006) Fast shrinkage of tropical glaciers in Colombia. *Annals of Glaciology*, 43, 194–201.
- CUESTA, F., CARILLA, J., LLAMBÍ, L.D., MURIEL, P., LENCINAS, M.V., MENESES, R.I. et al. (2023) Compositional shifts of alpine plant communities across the high Andes. *Global Ecology and Biogeography*, 32, 1591–1606.
- CUESTA, F., LLAMBÍ, L.D., HUGGEL, C., DRENKHAN, F., GOSLING, W.D., MURIEL, P. et al. (2019) New land in the Neotropics: a review of biotic community, ecosystem, and landscape transformations in the face of climate and glacier change. *Regional Environmental Change*, 19, 1623–1642.
- CUESTA, F., TOVAR, C., LLAMBÍ, L.D., GOSLING, W.D., HALLOY, S., CARILLA, J. et al. (2020) Thermal niche traits of high alpine plant species and communities across the tropical Andes and their vulnerability to global warming. *Journal of Biogeography*, 47, 408–420.
- DÍAZ, M., MONFORT-LANZAS, P., QUIROZ-MORENO, C., RIVADENEIRA, E., CASTILLEJO, P., ARNAU, V. et al. (2023) The microbiome of the ice-capped Cayambe Volcanic Complex in Ecuador. *Frontiers in Microbiology*, 14, 1154815.
- DUSSAILLANT, I., BERTHIER, E., BRUN, F., MASIOKAS, M., HUGONNET, R., FAVIER, V. et al. (2019) Two decades of glacier mass loss along the Andes. *Nature Geoscience*, 12, 802–808.
- EDWARDS, J.S. (1987) Arthropods of alpine aeolian ecosystems. *Annual Review of Entomology*, 32, 163–179.
- ETTER, A., ANDRADE, A., SAAVEDRA, K. & CORTÉS, J. (2018) Actualización de la Lista Roja de los Ecosistemas Terrestres de Colombia: conocimiento del riesgo de ecosistemas como herramienta para la gestión. In *Biodiversidad 2017. Estado y Tendencias de la Biodiversidad Continental de Colombia* (eds L.A. Moreno, C. Rueda & G.I. Andrade), pp. 1–9. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia.
- FARINOTTI, D., HUSS, M., FÜRST, J.J., LANDMANN, J., MACHGUTH, H., MAUSSION, F. & PANDIT, A. (2019) A consensus estimate for the ice thickness distribution of all glaciers on Earth. *Nature Geoscience*, 12, 168–173.
- FERRER-PARIS, J.R. & KEITH, D.A. (2024) Trade-offs in the use of direct and indirect indicators of ecosystem degradation for risk assessment. *Ecological Indicators*, 160, 111790.
- FICETOLA, G.F., MARTA, S., GUERRIERI, A., GOBBI, M., AMBROSINI, R., FONTANETO, D. et al. (2021) Dynamics of ecological communities following current retreat of glaciers. *Annual Review of Ecology, Evolution, and Systematics*, 52, 405–426.
- GILARDONI, S., DI MAURO, B. & BONASONI, P. (2022) Black carbon, organic carbon, and mineral dust in South American tropical glaciers: a review. *Global and Planetary Change*, 213, 103837.
- HAMBURGER, T., MATISÁNS, M., TUNVED, P., STRÖM, J., CALDERÓN, S., HOFFMANN, P. et al. (2013) Long-term in situ observations of biomass burning aerosol at a high altitude station in Venezuela – sources, impacts and interannual variability. *Atmospheric Chemistry and Physics*, 13, 9837–9853.
- HOTALING, S., HOOD, E. & HAMILTON, T.L. (2017) Microbial ecology of mountain glacier ecosystems: biodiversity, ecological connections and implications of a warming climate. *Environmental Microbiology*, 19, 2935–2948.
- HUGONNET, R., MCNABB, R., BERTHIER, E., MENOUNOS, B., NUTH, C., GIROD, L. et al. (2021) Accelerated global glacier mass loss in the early twenty-first century. *Nature*, 592, 726–731.

- HUSS, M., BOOKHAGEN, B., HUGGEL, C., JACOBSEN, D., BRADLEY, R.S., CLAGUE, J.J. et al. (2017) Toward mountains without permanent snow and ice. *Earth's Future*, 5, 418–435.
- JOMELLI, V., FAVIER, V., RABATEL, A., BRUNSTEIN, D., HOFFMANN, G. & FRANCOU, B. (2009) Fluctuations of glaciers in the tropical Andes over the last millennium and palaeoclimatic implications: a review. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 281, 269–282.
- KARGER, D.N., CONRAD, O., BÖHNER, J., KAWOHL, T., KREFT, H., SORIA-AUZA, R.W. et al. (2017) Climatologies at high resolution for the Earth's land surface areas. *Scientific Data*, 4, 170122.
- KEITH, D.A., FERRER-PARIS, J.R., NICHOLSON, E., BISHOP, M.J., POLIDORO, B.A., RAMIREZ-LLODRA, E. et al. (2022) A function-based typology for Earth's ecosystems. *Nature*, 610, 513–518.
- KEITH, D.A., RODRÍGUEZ, J.P., RODRÍGUEZ-CLARK, K.M., NICHOLSON, E., AAPALA, K., ALONSO, A. et al. (2013) Scientific foundations for an IUCN Red List of Ecosystems. *PLOS One*, 8, e62111.
- KHEDIM, N., CÉCILLON, L., POULENARD, J., BARRÉ, P., BAUDIN, F., MARTA, S. et al. (2021) Topsoil organic matter build-up in glacier forelands around the world. *Global Change Biology*, 27, 1662–1677.
- KONTULA, T. & RAUNIO, A. (eds) (2019) *Threatened Habitat Types in Finland 2018 – Red List of Habitats Results and Basis for Assessment*. The Finnish Environment 2/19, Finnish Environment Institute and Ministry of the Environment, Helsinki, Finland. [urn.fi/URN:ISBN:978-952-11-5110-1](https://urn.fi/URN:ISBN:978-952-11-5110-1) [accessed April 2024].
- KUHN, M. (2008) Building predictive models in R using the *caret* package. *Journal of Statistical Software*, 28, 1–26.
- LLAMBÍ, L.D., BECERRA, M.T., PERALVO, M., AVELLA, A., BARUFFOL, M. & DÍAZ, L.J. (2020) Monitoring biodiversity and ecosystem services in Colombia's high Andean ecosystems: toward an integrated strategy. *Mountain Research and Development*, 39, A8–A20.
- LLAMBÍ, L.D., MELFO, A., GAMEZ, L.E., PELAYO, R.C., CÁRDENAS, M., ROJAS, C. et al. (2021) Vegetation assembly, adaptive strategies and positive interactions during primary succession in the forefield of the last Venezuelan glacier. *Frontiers in Ecology and Evolution*, 9, 657755.
- MASIOKAS, M.H., RABATEL, A., RIVERA, A., RUIZ, L., PITTE, P., CEBALLOS, J.L. et al. (2020) A review of the current state and recent changes of the Andean cryosphere. *Frontiers in Earth Science*, 8, 99.
- MURRAY, N.J., KEITH, D.A., DUNCAN, A., TIZARD, R., FERRER-PARIS, J.R., WORTHINGTON, T.A. et al. (2020) Myanmar's terrestrial ecosystems: status, threats and conservation opportunities. *Biological Conservation*, 252, 108834.
- OLIVEIRA-MIRANDA, M.A., HUBER, O., RODRÍGUEZ, J.P., ROJAS-SUAREZ, F., DE OLIVEIRA-MIRANDA, R., ZAMBRANO-MARTINEZ, S. & HERNANDEZ, D.G. (2010) Riesgo de eliminación de los ecosistemas terrestres de Venezuela. In *Libro Rojo de los Ecosistemas Terrestres de Venezuela* (eds J.P. Rodríguez, F. Rojas-Suárez & D.G. Hernández), pp. 107–236. Proviata, Shell Venezuela and Lenovo (Venezuela), Caracas, Venezuela.
- POLISSAR, P.J., ABBOTT, M.B., WOLFE, A.P., BEZADA, M., RULL, V. & BRADLEY, R.S. (2006) Solar modulation of little ice age climate in the tropical Andes. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 8937–8942.
- PULWARTY, R.S., BARRY, R.G., HURST, C.M., SELLINGER, K. & MOGOLLON, L.F. (1998) Precipitation in the Venezuelan Andes in the context of regional climate. *Meteorology and Atmospheric Physics*, 67, 217–237.
- R CORE TEAM (2023) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. [r-project.org](https://r-project.org) [accessed April 2024].
- RABATEL, A., CEBALLOS, J.L., MICHELETTI, N., JORDAN, E., BRAITMEIER, M., GONZÁLEZ, J. et al. (2018) Toward an imminent extinction of Colombian glaciers? *Geografiska Annaler: Series A, Physical Geography*, 100, 75–95.
- RABATEL, A., FRANCOU, B., SORUCO, A., GOMEZ, J., CÁCERES, B., CEBALLOS, J.L. et al. (2013) Current state of glaciers in the tropical Andes: a multi-century perspective on glacier evolution and climate change. *The Cryosphere*, 7, 81–102.
- RAMÍREZ, N., MELFO, A., RESLER, L.M. & LLAMBÍ, L.D. (2020) The end of the eternal snows: integrative mapping of 100 years of glacier retreat in the Venezuelan Andes. *Arctic, Antarctic, and Alpine Research*, 52, 563–581.
- RANDOLPH GLACIER INVENTORY CONSORTIUM (2017) *Randolph Glacier Inventory 6.0*. [glims.org/RGI/randolph60.html](https://glims.org/RGI/randolph60.html) [accessed February 2024].
- RONDÓN, J., GÓMEZ, W., BALL, M.M., MELFO, A., RENGIFO, M., BALCAZAR, W. et al. (2016) Diversity of culturable bacteria recovered from Pico Bolívar's glacial and subglacial environments, at 4950 m, in Venezuelan tropical Andes. *Canadian Journal of Microbiology*, 62, 904–917.
- ROSETO, P., CRESPO-PÉREZ, V., ESPINOSA, R., ANDINO, P., BARRAGÁN, Á., MORET, P. et al. (2021) Multi-taxa colonisation along the foreland of a vanishing equatorial glacier. *Ecography*, 44, 1010–1021.
- ROUNCE, D.R., HOCK, R. & MAUSSION, F. (2022) *Global PyGEM-OGGM Glacier Projections with RCP and SSP Scenarios, Version 1*. NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, USA. [doi.org/10.5067/P8BN9VO9N5C7](https://doi.org/10.5067/P8BN9VO9N5C7).
- ROUNCE, D.R., HOCK, R., MAUSSION, F., HUGONNET, R., KOCHTITZKY, W., HUSS, M. et al. (2023) Global glacier change in the 21st century: every increase in temperature matters. *Science*, 379, 78–83.
- SAGREDO, E.A. & LOWELL, T.V. (2012) Climatology of Andean glaciers: a framework to understand glacier response to climate change. *Global and Planetary Change*, 86–87, 101–109.
- STIBAL, M., BRADLEY, J.A., EDWARDS, A., HOTALING, S., ZAWIERUCHA, K., ROSVOLD, J. et al. (2020) Glacial ecosystems are essential to understanding biodiversity responses to glacier retreat. *Nature Ecology & Evolution*, 4, 686–687.
- VEETTL, B.K. & KAMP, U. (2019) Global disappearance of tropical mountain glaciers: observations, causes, and challenges. *Geosciences*, 9, 196.
- VUILLE, M., YARLEQUE, C., ELISON TIMM, O., CAREY, M., HUGGEL, C., SALZMANN, N. et al. (2018) Rapid decline of snow and ice in the tropical Andes – impacts, uncertainties and challenges ahead. *Earth-Science Reviews*, 176, 195–213.
- WILLIAMS, R.J., STOTT, K.A.J., CAMAC, J.S., WHITE, M., BURNS, E., HARRIS, S. et al. (2015) An International Union for the Conservation of Nature Red List ecosystems risk assessment for alpine snow patch herbfields, South-Eastern Australia. *Austral Ecology*, 40, 433–443.