

This is a “preproof” accepted article for Invasive Plant Science and Management.

This version may be subject to change in the production process, *and does not include access to supplementary material*. DOI: 10.1017/inp.2025.14

Invasion alert: Lindenleaf sage (*Salvia tiliifolia*) in the Tigray highlands

Jan Nyssen^{1,*}, Rishan Haile², Getachew Gebremedhin³, Zbelo Tesfamariam⁴, Jesús Guadalupe González-Gallegos⁵ and Tesfaalem Ghebreyohannes⁴

¹Professor emeritus, Department of Geography, Ghent University, Ghent, Belgium, and Guest professor, Department of Geography and Environmental Studies, Mekelle University, Mekelle, Tigray, Ethiopia

²Lecturer, Department of Geography and Environmental Studies, Mekelle University, Mekelle, Tigray, Ethiopia

³Research technician, Hagere Selam, Tigray, Ethiopia

⁴Associate professor, Department of Geography and Environmental Studies, Mekelle University, Mekelle, Tigray, Ethiopia

⁵Senior scientist, Instituto Politécnico Nacional, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional – Unidad Durango, Durango, Mexico

***Author for correspondence:** Jan Nyssen, Professor emeritus, Department of Geography, Ghent University, Krijgslaan 281 (S8), B-9000 Gent, Belgium. (E-mail: jan.nyssen@ugent.be)

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Abstract

Lindenleaf sage (*Salvia tiliifolia* Vahl) is an annual plant native to Mesoamerica that thrives in tropical, seasonally dry environments. It was introduced to Ethiopia in the late 20th century and made its way to the Tembien highlands in Tigray in 2018. With its many branching stems and profusion of seeds, this upright pioneer plant becomes dominant in the herbaceous layer of intensively used rangeland and roadsides before spreading to areas with other land uses, including agricultural land. Visual observations show that in areas with less grazing pressure, *S. tiliifolia* is not prevalent. Farmers uproot *S. tiliifolia* from their crops, which is rather simple because this species is easily recognized, and its roots do not withstand much force. On rangeland, weeding is labor-intensive. As an alternative, we recommend converting rangelands into managed exclosures to maximize grass production, with initial *S. tiliifolia* removal. Natural vegetation regrowth on infested places would keep the most significant *S. tiliifolia* seed supply under control and reduce the species' invasion of farmlands.

Keywords: Invasive exotic plants, phytogeography, weed control, seed bank, exclosure.

Introduction

Native to Mesoamerica, lindenleaf sage or *Salvia tiliifolia* Vahl (Lamiaceae) (Figure 1) is mostly found in tropical, seasonally dry habitat (Kew 2024; Standley and Williams 1973); it has expanded to other parts of the world, including Ethiopia. There, the growth of grasses in rangeland is strongly suppressed by *S. tiliifolia* (Supplementary Figure S1).

In its native Mexico, *S. tiliifolia* grows as weed in crops, but also in ruderal environments or in the borders and clearings of native vegetation (Pichardo et al. 2009; González-Gallegos et al. 2016). In recent centuries, the plant has been transported to other continents, most probably accidentally together with commercial products. Deliberate transplantation of *S. tiliifolia* seems unlikely, as the species does not exhibit a high ornamental value with its small and short-lived flowers and its weedy behavior (Froissart 2008); for instance, Clebsch does not include it in her compilation of ornamental salvias (Clebsch 1997). The expansion of *S. tiliifolia* has been quite rapid after 1980 (Supplementary Figure S2).

In South Africa, *S. tiliifolia* has been in the Pretoria area since the 1940s (pers. comm. Mark van Dalsen, iNaturalist, 23 September 2024) but it became a menace in many parts of the country after 2010 (Anon. 2015). There, it grows in dense stands in both sunny and partially shaded sites, overruns rocky hillsides, roadsides, waste sites, urban open spaces, and prefers moist places (Anon. 2024, 2015). It competes with native species and may eventually supplant them (Anon. 2024). In southern China, *S. tiliifolia* first appeared in the 1990s and is regarded there as having a high risk of becoming an invasive plant (Hu et al. 2013).

Salvia tiliifolia spreads by seeds that drop when the plant dries after the rainy season (Supplementary Figure S3); the seeds may be transported to adjacent places by mammals and birds (Anon. 2024). In its region of origin, Mesoamerica, it is not considered as a major problem, though it occurs as weed in coffee orchards (Standley and Williams 1973). Further, it grows in secondary vegetation, along roads and at the edges of field crops. It usually does not grow in big numbers. Whereas *S. tiliifolia* is weeded out from croplands, jointly with other species (Molina-Freaner et al. 2008), there are no targeted weed control programs for it. The situation is however different in other continents. In southern China, it was suggested that land managers in the area monitor the species and take action to stop its spread (Hu et al. 2013). In South Africa, *S. tiliifolia* is regarded as a difficult plant to control since it just keeps growing back and cannot be eradicated until every plant in an area is removed (Kyle

Campbell, pers. comm., 25 September 2024); hence, there it is recommended to uproot the plants before they flower and set seed (Anon. 2015).

In Ethiopia, *Salvia tiliifolia* is a recent invasive exotic species, first formally collected and described near Kombolcha in 1996 (Demissew 1996). It is now quite widespread over the Ethiopian highlands. Whereas naturalists (Supplementary Figure S4) and field inventories (Dagne and Birhanu 2023; Atsbeha 2012; Bersisa et al. 2021) reported observations on the occurrence of *S. tiliifolia*, there are few publications about the behavior and environmental preferences of this new invasive species in Ethiopia. Near Harar in eastern Ethiopia, *S. tiliifolia* was observed to be the dominant herbaceous species in coffee plantations, with highest mean field density and highest relative abundance (32%) (Bersisa et al. 2021). In 2012, the species was reported around Aksum in Tigray (Atsbeha 2012). Most studies in Ethiopia (Bersisa et al. 2021; Atsbeha 2012; Demissew 1996) explicitly mention that the species has no local name (except ‘new weed’), which stresses its relatively recent introduction. In the Tigray region, where we studied it, farmers are very worried by this invasive species, especially as they are already coping with the social, agronomic and environmental impacts of the recent Tigray war (Nyssen et al. 2023; Weldegiargis et al. 2023).

Here we discuss the spread of *S. tiliifolia* in the Tigray highlands (northern Ethiopia), and efforts to control it. As management options on communal lands, the systematic removal approach is contrasted to the nature-based approach, i.e. the restriction of ranging livestock to allow native vegetation to outcompete *S. tiliifolia*.

Methods

Plant traits

Salvia tiliifolia is an erect, short hairy, annual herb with many branched stems, between 20 cm and 1.6 m high (Figure 1). The soft, bright green leaves are arranged in pairs that alternate at a 90° angle from one another. The leaves are simple, ovate with truncate base, serrate margin, usually with wide teeth. *S. tiliifolia* has tiny blue flowers, upper calyx lip with three veins, corolla 4 - 10 mm long, clustered in spikes up to 30 cm long. The calyx elongates to 10 mm in fruit, enclosing 3-angled ovoid nutlets (mericarps), brown and irregularly marbled with a darker tone (Anon. 2024, 2015; Pichardo et al. 2009; Standley and Williams 1973).

The seeds (mericarps) are known as “chía” or “chía cimarrona” and constitute a staple food in Mexico and Central America, next to the cultivated *Salvia hispanica*, the species commercially known as chía (Ayerza and Coates 2005; Martínez Hernández 2017; Pichardo et al. 2009).

Field investigation in Tembien

The research region, the Tembien area is situated between 30 and 120 km west of Mekelle, the regional capital of Tigray. The geology of the area is made up of Precambrian metamorphics, Mesozoic sedimentary rocks, Tertiary volcanics and Quaternary lava flows (Gebreyohannes et al. 2010). A network of deeply incised rivers defines the landscape, with elevations between 1500 and 2900 meters. A stepped geomorphology was produced by selective erosion as a result of the alternation of different lithologies (Nyssen et al. 2006; Coltorti et al. 2007).

The region is classified as hot semi-arid (BSH) by the Köppen climate classification (Peel et al. 2007). The depth of annual rainfall ranges from 500 mm to 900 mm (Jacob et al. 2013). Most rain falls in the main rainy season, which usually lasts from June to September. Except for the rainy season, when there are fewer sunshine hours and more rainfall, monthly potential evapotranspiration exceeds monthly rainfall. In one of the major towns, Hagere Selam, yearly rainfall from 2016 to 2022 ranged from 770 to 850 mm, 7 to 18% over the long-term normal. There is a tendency toward increased precipitation and warmer temperatures over time. While the rainy season in 2024 was unusually wet, the drought in 2023 was accompanied with anomalies in rain seasonality (Nyssen et al. 2024b).

In this mountainous area, the farming communities typically use steeper slopes as communal rangeland, and then they use the more level areas for crops. Free grazing is still the dominant way of feeding livestock, which has even expanded due to the Tigray War (Nyssen et al. 2024b). Oxen are used for farm operations such as plowing and threshing in the small-scale family farms of Tigray, which follow a permanent farming system based on cereals (Westphal 1975). Goats and sheep are also raised, primarily as a safety net in case of emergencies (Nyssen et al. 2008). Since crop cultivation has been practiced in Tigray for at least three millennia, the agricultural system has been gradually optimized (D’Andrea 2008; Blond et al. 2018). After harvest, stubble grazing occurs on cropland as well. Over the past 40 years, many sloping lands—both rangelands and marginal croplands—have been turned into

exclosures, fended from human and livestock interference (Aerts et al. 2009; Nyssen et al. 2008). Linear landscape features, including cliffs and gullies, act as demarcations for these exclosures, with regulations outlining their use. Unlike open grazing land, where those with larger herds tend to monopolize the available biomass, exclosures facilitate communal collection and sharing of grasses, fostering equity within the community. Therefore, in Tigray, the transformation of an area of rangeland into an exclosure is a decision made with considerable deliberation. The natural vegetation is mainly open *Olea-Acacia* woodland, remnants of the primary dry Afromontane evergreen forests (Asmelash and Rannestad 2024). The woodlands have a low, single-story, discontinuous canopy and comparatively few tree species. Under this open canopy, grasses and herbaceous vegetation appear during the rainy season (Aerts 2019).

To investigate *S. tiliifolia*'s rapid spread in Ethiopia's semi-arid Tigray region, and how to control it both on privately managed cropland and on communally owned rangeland and exclosures, we carried out field observations on its distribution and density in various habitats of Tembien between November 2023 and October 2024 (Nyssen 2024), both during the growing and the dry season. Forty-three different sites were visited, and at fourteen locations open-ended interviews (Albudaiwi 2017) were conducted with community members who were present on-site.

Results and discussion

Salvia tiliifolia invasion in Tembien

Salvia tiliifolia arrived suddenly but relatively late in Tembien. Dr. Raf Aerts (K.U. Leuven University), who made extensive inventories of plant species in Tembien during the period 2001-2006, confirmed that this species was absent at the time (personal communication). Farmers commonly mention approx. 2018 as its time of arrival around Hagere Selam, where it has invaded rangelands surrounding the town, particularly those with heavy livestock browsing (Supplementary Figures S1 and S5). There are several native *Salvia* species in the study region (*S. schimperi*, *S. merjamie*, *S. nilotica*), with their own local names and which do not have such an invasive behavior (Endeshaw et al. 2000; November et al. 2002; Seegeler 1983; Edwards 1991).

Besides rangeland near towns, we observed *S. tiliifolia* as a dominant species in nearly all grazed areas: open *Eucalyptus* or *Acacia* woodlands, formal waste dumps, rocky gorges, coarse alluvial deposits, sides of footpaths and dirt roads. The plants showed however stunted growth under the shadow of trees, and in such places seeds do not fully mature. *S. tiliifolia* was conspicuous, but not dominant, in the herbaceous layer in places with less grazing pressure such as grass strips between cropped parcels, gullies and banks of ephemeral waterways, as well as on permanently moist places such as banks of grassed waterways and at the edges of a permanent springs. We have not seen *S. tiliifolia* in rainfed and irrigated croplands, because the species is weeded out at early growth stage. There were no observations either in exclosures with dense vegetation. In some pocket areas with various land cover, *S. tiliifolia* was also absent. The dominance of *S. tiliifolia* in the herbaceous layer of grazed areas seems an extreme manifestation of the typical mechanism by which invasive species proliferate in rangelands. Browsing livestock continually removes the other herbs while avoiding the unpalatable *S. tiliifolia*, except for occasional browsing of the infructescence. The low grass and other herbs are further suppressed by *S. tiliifolia*. Visual observations show that *S. tiliifolia* is not prevalent in places with reduced grazing pressure - such as exclosures, i.e. communal lands where open access for grazing or wood harvesting has been abandoned (Aerts et al. 2009). Quantitative ecological research on this aspect would be appreciated.

Voucher Specimen. Ethiopia, Tigray, Dogu'a Tembien: Hagere Selam, northern part of the town, in secondary vegetation on side of gravel road. 13°39'05" N, 39°10'21" W, 2633 m. February 5, 2025. *Getachew Gebremedhin 1* (LUX herbarium, accession number MNHNL2025/12).

Pathways of the invasion

Demissew (1996) suggests that *Salvia tiliifolia* arrived in Ethiopia through weed-seed contaminated food aid in the 1980s. Grain trade is a known pathway for the introduction of alien plants (Ikeda et al. 2022). The fact that *S. tiliifolia* was first formally observed near Kombolcha, which was a hub for international food aid since the Ethiopian Famine in 1983-1985 (Augenstein 2020) supports this interpretation. On the other hand, no extensive surveys have been done, and the plant may also have been introduced otherwise, and travelled along roads. Similarly, tree tobacco (*Nicotiana glauca*), a woody noxious invasive species in the

study area (Figure 2), is also said to have come in that period with food aid; yet specimens of the plant had already been collected in a garden in Addis Ababa in 1960, and at Haromaya University campus in 1975 (Nyssen 1997).

The pathways of the *S. tiliifolia* spread need to be studied further; the spread over the Ethiopian highlands would logically have followed roads, as dusty lorries transport thousands of seeds of alien plants (Bajwa et al. 2018; Khan et al. 2018) to suitable environmental conditions. The extent to which multiple traffic and war disturbances may have contributed to this spread is unknown. Further, seed dispersal by runoff water was evidenced in the study area by the presence of *S. tiliifolia* on sediment deposits in ephemeral streams down from major infested areas.

The wetter conditions in the study area in the last decades and particularly in recent years (Nyssen et al. 2024b), jointly with increasing temperatures, may have created a niche for the species, in line with observations of natural vegetation and crop belts that shift up the mountains in northern Ethiopia (Nyssen et al. 2015; Jacob et al. 2017). The fact that in South Africa, the *S. tiliifolia* spread was observed to be strongly subdued in dry years (Kyle Campbell, iNaturalist, pers. comm., 25 September 2024) supports this hypothesis.

The large extent and dominance of the plant, with its specific phenology (annual pattern of *S. tiliifolia*'s lifecycle) would be ideal for studies using multispectral satellite imagery with high spatiotemporal resolution. The plant with its dark-blueish green color appears in July, occupies large patches, and grows up to the end of September, when it rapidly decays to homogenous brown-yellowish patches. These colors are unique among the background vegetation and therefore can be picked up through multispectral imagery analysis, what allows detailed studies of its current extent, as well as pathways of diffusion over the last decades.

***Salvia tiliifolia* management on cropland in Tigray**

According to local farmers, *Salvia tiliifolia* is now the main weed to be removed from croplands in Tigray; it grows rapidly and even suppresses other weed species. On well-managed farmlands one will generally not observe full-grown *S. tiliifolia* (Figure 2). The farmers in the study area have experience with manually uprooting *S. tiliifolia*. While the work in itself is quite easy, because the species is very recognizable, and the roots do not

resist much (Supplementary Figure S6), it involves additional workload, and is a burden, particularly for women and children. However, and depending on time constraints, it is often not weeded out from farm boundaries (Figure 2) and rangeland nearby farmlands, which will be a source of *S. tiliifolia* seed in the next cropping season.

***Salvia tiliifolia* management on communal lands**

The conventional strategy has long been the extensive and methodical clearance of invasive plants before they set seed (Mack and Foster 2009; Parmesan and Hanley 2015; Engel et al. 2024). Whereas it is impossible in this way to totally “get rid of” the invasive plant, it may subdue the growth in that area in the next year, and with annual repetitive removal, exhaust the seed bank (Anon. 2015, 2024). In the study area, experimental community weeding of *S. tiliifolia* on rangeland showed that 45 person-days per ha are necessary for an initial manual *S. tiliifolia* removal operation (Supplementary Figure S7) (Nyssen et al. 2024c).

Reversely, in communal lands where free grazing is forbidden (exclosures), we observed that *S. tiliifolia* is totally absent, particularly at locations away from roads (Figure 3). Also, we observed that *S. tiliifolia* has a stunted growth in shady places with seeds that do not fully mature. Locally, *S. tiliifolia* coexists with other herbs in peri-urban exclosures, but it does not dominate. In such cases, the grasses and herbs—including *S. tiliifolia*—are all harvested together (Supplementary Figure S8), to serve as fodder for livestock. In contrast to thistles for instance, farmers accept withered *S. tiliifolia* as a part of the harvested hay. Establishing exclosures hence seems the most cost-effective strategy to control *S. tiliifolia*, while sustaining fodder production. In this nature-based approach (Ngondya and Munishi 2022; Tracy et al. 2004; Ammond and Litton 2012), the recovery of herbaceous and woody vegetation in newly established exclosures would be sped up by *S. tiliifolia* removal in the first years. Reversely, opening the exclosures for free grazing to relieve the fodder crisis caused by the *S. tiliifolia* invasion would only result in the opposite effect, with the exclosures being invaded by *S. tiliifolia* in turn, due to selective grazing. All in all, the mere protection of the indigenous vegetation enables subduing *S. tiliifolia* at a lower economic cost (Lázaro-Lobo and Ervin 2021; Leuzinger and Rewald 2021; Nyssen et al. 2024a).

The ability of *S. tiliifolia* to dominate rangelands and suppress grasses, particularly in areas of intensive livestock grazing, underscores the urgent need for effective management strategies. Field observations and local knowledge suggest that a combination of weeding on

cropland and the establishment of exclosures, where only haymaking but no grazing is allowed, may offer the most sustainable approach to controlling its spread.

Supplementary material.

Supplementary Figures S1 to S8 are available as Supplementary Material at <https://www.cambridge.org/core/journals/invasive-plant-science-and-management/article/xxx>

Data availability statement

Species observation locations (with photographs) are reported in the iNaturalist database and reported as a reference to the article (Nyssen 2024).

Acknowledgments

We thank Kyle Campbell, Mark van Dalsen (iNaturalist) and Raf Aerts (K.U.Leuven) for their help in preliminary identification of the species. Abraha Teklu and the Meserete Birhan 'Idir traditional neighborhood solidarity group in Hagere Selam are also acknowledged for all assistance. Gebrekidan Mesfin and Miro Jacob (EthioTrees) provided photos and coordinates of areas with dense *S. tiliifolia* growth. Sofie Annys (Ghent University) assisted in voucher specimen collection and handling. We thank Odile Weber and the Botany team of the Musée national d'histoire naturelle du Luxembourg for mounting and housing the voucher specimen.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing interests

The authors declare none.

References

- Aerts R (2019) Forest and Woodland Vegetation in the Highlands of Dogu'a Tembien. In: Geo-trekking in Ethiopia's Tropical Mountains. Springer, pp 233-250
- Aerts R, Nyssen J, Mitiku Haile (2009) On the difference between "exclosures" and "enclosures" in ecology and the environment. *Journal of Arid Environments* 73:762-763
- Albudaiwi D (2017) Survey: Open-ended questions. In: The SAGE Encyclopedia of Communication Research Methods, vol 1. SAGE Publications, Inc., Thousand Oaks, CA, USA, pp 1715-1716
- Ammond SA, Litton CM (2012) Competition between native Hawaiian plants and the invasive grass *Megathyrsus maximus*: implications of functional diversity for ecological restoration. *Restor Ecol* 20 (5):638-646
- Anon. (2015) Lindenleaf sage: control this invader! SAPIA News - Southern African Plant Invaders Atlas 36:1 and 6
- Anon. (2024) Fact sheet: Lindenleaf sage - *Salvia tiliifolia*. Invasives South Africa. <https://invasives.org.za/fact-sheet/lindenleaf-sage/>. Accessed 24/03/2025
- Asmelash F, Rannestad M (2024) Challenges and a Strategy for Successful Restoration of Dry Evergreen Afromontane Forests of Ethiopia. *Ecological Restoration* 42:220-237. doi:10.3368/er.42.3.220
- Atsbeha G (2012) Floristic Composition of Herbaceous Flowering Plant Species with Emphasis to Ethnobotanical Importance of Wild Grasses in Laelay and Tahtay Michew Districts, Central Zone of Tigray, Ethiopia. Addis Ababa University,
- Augenstein P (2020) The End of Famine. <https://www.youtube.com/watch?v=vgP9W9amAM8>. Accessed 24/03/2025
- Ayerza R, Coates W (2005) Chia: rediscovering a forgotten crop of the Aztecs. University of Arizona Press,
- Bajwa AA, Nguyen T, Navie S, O'Donnell C, Adkins S (2018) Weed seed spread and its prevention: the role of roadside wash down. *Journal of Environmental Management* 208:8-14
- Bersisa H, Adisu Wagari, Taddese Ishetu (2021) Survey of weed flora composition in coffee (*Coffea arabica* L.) growing areas of East Ethiopia. *Results of Plant Protection Research*:278

- Blond N, Jacob-Rousseau N, Callot Y (2018) Terrasses alluviales et terrasses agricoles. Première approche des comblements sédimentaires et de leurs aménagements agricoles depuis 5000 av. n. è. à Wakarida (Éthiopie). *Géomorphologie: Relief, Processus, Environnement* 24 (3):277-300. doi:10.4000/geomorphologie.12258
- Clebsch B (1997) *A book of salvias: sages for every garden*. Timber Press, Portland, United States of America
- Coltorti M, Dramis F, Ollier C (2007) Planation surfaces in northern Ethiopia. *Geomorphology* 89 (3-4):287-296
- D'Andrea AC (2008) T'ef (*Eragrostis tef*) in ancient agricultural systems of highland Ethiopia. *Economic Botany* 62 (4):547-566. doi:doi.org/10.1007/s12231-008-9053-4
- Dagne Y, Birhanu L (2023) Floristic composition and plant community distribution along environmental gradients in Guard dry Afromontane forest of Northwestern Ethiopia. *BMC Ecology and Evolution* 23 (1):43
- Demissew S (1996) A Central American weedy *Salvia* in Ethiopia. *Lamiales Newsletter* 4:3-4
- Edwards SB (1991) Crops with wild relatives found in Ethiopia. In: Engels J, Hawkes JG, Melaku Worede (eds) *Plant genetic resources of Ethiopia*. Cambridge University Press, pp 42-74
- Endeshaw MM, Gautun O, Asfaw N, Aasen A (2000) Volatile oil constituents of the Ethiopian plant *Salvia schimperii* Benth. *Flavour and fragrance journal* 15 (1):27-30
- Engel M, Nyssen B, Desie E, den Ouden J, Raats L, Hagemann U (2024) Managing Black Cherry (*Prunus serotina* Ehrh.) in European Forests: Insights from native and non-native ranges. *Forest Ecology and Management* 562:121959
- Froissart C (2008) *La connaissance des sauges*. Édisud, Aix-en-Provence, France
- Gebreyohannes T, De Smedt F, Miruts Hagos, Solomon Gebresilassie, Kassa Amare, Kurkura Kabeto, Abdulwassie Hussein, Nyssen J, Bauer H, Moeyersons J (2010) Large-scale geological mapping of the Geba basin, northern Ethiopia, vol 9. Tigray Livelihood Papers. VLIR-Mekelle University IUC Programme, Mekelle, Ethiopia
- González-Gallegos JG, Castro-Castro A, Quintero-Fuentes V, Mendoza-López ME, de Castro Arce E (2016) Revisión taxonómica de Lamiaceae del occidente de México. *Ibutana* (7):3-545
- Hu G, Xiang C, Liu E (2013) Invasion status and risk assessment for *Salvia tiliifolia*, a recently recognised introduction to China. *Weed Research* 53 (5):355-361

- Ikeda M, Nishi T, Asai M, Muranaka T, Konuma A, Tominaga T, Shimono Y (2022) The role of weed seed contamination in grain commodities as propagule pressure. *Biological Invasions* 24 (6):1707-1723
- Jacob M, Frankl A, Hurni H, Lanckriet S, De Ridder M, Guyassa E, Beeckman H, Nyssen J (2017) Land cover dynamics in the Simien Mountains (Ethiopia), half a century after establishment of the National Park. *Regional Environmental Change* 17 (3):777-787
- Jacob M, Frankl A, Mitiku Haile, Zwertvaegher A, Nyssen J (2013) Assessing spatio-temporal rainfall variability in a tropical mountain area (Ethiopia) using NOAA's Rainfall Estimates. *International Journal of Remote Sensing* 34 (23):8305-8321. doi:10.1080/01431161.2013.837230
- Kew (2024) *Salvia tiliifolia* Vahl. Kew Royal Botanical Gardens. <https://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:457397-1>. Accessed 15/10/2024
- Khan I, Navie S, George D, O'donnell C, Adkins S (2018) Alien and native plant seed dispersal by vehicles. *Austral Ecology* 43 (1):76-88
- Lázaro-Lobo A, Ervin GN (2021) Native and exotic plant species respond differently to ecosystem characteristics at both local and landscape scales. *Biological Invasions* 23 (1):143-156
- Leuzinger S, Rewald B (2021) The who or the how? Species vs. ecosystem function priorities in conservation ecology. *Frontiers in Plant Science* 12:758413
- Mack RN, Foster SK (2009) Eradicating Plant Invaders: Combining Ecologically-Based Tactics and Broad-Sense Strategy. In: Inderjit (ed) *Management of Invasive Weeds*. Springer Netherlands, Dordrecht, pp 35-60. doi:10.1007/978-1-4020-9202-2_3
- Martínez Hernández C (2017) Propiedades funcionales de péptidos de semillas de chía comercial (*Salvia hispanica*) y silvestre (*Salvia tiliifolia*). *Jóvenes en la Ciencia* (Universidad de Guanajuato) 3 (1):139-143
- Molina-Freaner F, Espinosa-García F, Sarukhán-Kermez J (2008) Weed population dynamics in a rain-fed maize field from the Valley of Mexico. *Agrociencia* 42 (5):499-511
- Ngondya IB, Munishi LK (2022) Managing invasive plants through a nature-based approach in complex landscapes. *Trends in Ecology & Evolution* 37 (4):284-288
- November E, Aerts R, Mintesinot Behailu, Muys B (2002) Species list *Tigrinya* – Scientific. Technical note 2002/4. . Forest Rehabilitation Project, Mekelle University, Ethiopia and K.U. Leuven, Belgium.,

- Nyssen B, Den Ouden J, Bindewald A, Brancalion P, Kremer K, Lapin K, Raats L, Schatzdorfer E, Stanturf J, Verheyen K, Muys B (2024a) Established Invasive Tree Species Offer Opportunities for Forest Resilience to Climate Change. *Current Forestry Reports*:1-31. doi:10.1007/s40725-024-00232-6
- Nyssen J (1997) Vegetation and soil erosion in Dega Tembien (Tigray, Ethiopia). *Bulletin du Jardin botanique national de Belgique/Bulletin van de Nationale Plantentuin van België*:39-62
- Nyssen J (2024) *Salvia tiliifolia* observations in Dogu'a Tembien, Tanqwa Melash and Kayeh Tekli, Tigray, Ethiopia. iNaturalist. https://www.inaturalist.org/observations?subview=map&taxon_id=153953&user_id=j_an_nyssen. Accessed 24/03/2025
- Nyssen J, Emnet Negash, Hailemariam Meaza, Annys S, Mitiku Haile, Zbelo Tesfamariam, Poesen J, Deckers J, Moeyersons J, Amanuel Zenebe, Frankl A, Tesfaalem Ghebreyohannes (2024b) Land Resilience Amidst the Ravages of War: Insights from Tigray (northern Ethiopia). In preparation
- Nyssen J, Emnet Negash, Van Schaeybroeck B, Haegeman K, Annys S (2023) Crop Cultivation at Wartime – Plight and Resilience of Tigray's Agrarian Society (North Ethiopia). *Defence and Peace Economics* 34 (5):618-645. doi:10.1080/10242694.2022.2066420
- Nyssen J, Frankl A, Amanuel Zenebe, Deckers J, Poesen J (2015) Land management in the northern Ethiopian highlands: local and global perspectives; past, present and future. *Land Degradation & Development* 26 (7):759-764
- Nyssen J, Gebremedhin G, Tesfamariam Z, Haile R, González-Gallegos JG, Ghebreyohannes T (2024c) About the new salvia weed (*Salvia tiliifolia*) in Tigray. doi:<https://doi.org/10.5281/zenodo.14069091>
- Nyssen J, Naudts J, De Geyndt K, Mitiku Haile, Poesen J, Moeyersons J, Deckers J (2008) Soils and land use in the Tigray highlands (Northern Ethiopia). *Land Degradation & Development* 19 (3):257-274. doi:10.1002/ldr.840
- Nyssen J, Poesen J, Moeyersons J, Deckers J, Mitiku Haile (2006) Processes and rates of rock fragment displacement on cliffs and scree slopes in an amba landscape, Ethiopia. *Geomorphology* 81 (3-4):265-275
- Parmesan C, Hanley ME (2015) Plants and climate change: complexities and surprises. *Annals of botany* 116 (6):849-864

- Peel MC, Finlayson BL, McMahon TA (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11 (5):1633-1644. doi:10.5194/hess-11-1633-200
- Pichardo JM, Vibrans H, Lezama PT (2009) *Salvia tiliifolia* Vahl. In: Vibrans H (ed) *Malezas de México*. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México,
- Seegeler CJP (1983) *Oil Plants in Ethiopia, their Taxonomy and Agricultural Significance*. PUDOC, Wageningen
- Standley PC, Williams LO (1973) Labiatae. In: Standley PC, Steyermark JA (eds) *Flora of Guatemala*, vol 24, part 9(3). Natural History Museum, London, pp 237-317
- Tracy BF, Renne IJ, Gerrish J, Sanderson MA (2004) Effects of plant diversity on invasion of weed species in experimental pasture communities. *Basic and Applied Ecology* 5 (6):543-550
- Weldegiargis AW, Abebe HT, Abraha HE, Abrha MM, Tesfay TB, Belay RE, Araya AA, Gebregziabher MB, Godefay H, Mulugeta A (2023) Armed conflict and household food insecurity: evidence from war-torn Tigray, Ethiopia. *Conflict and Health* 17 (1):22
- Westphal E (1975) *Agricultural systems in Ethiopia*. Centre for Agricultural Publishing and Documentation, Wageningen



Figure 1. *Salvia tiliifolia* in Hagere Selam, Tigray, Ethiopia (13°39'04" N, 39°10'33" W, 2610 m), in flowering (A; 25 September 2024) and dry stages (B; 23 November 2023).



Figure 2. *Salvia tiliifolia* was weeded out of this wheat field in Hech'i (13°38'25" N, 39°12'17" W, 2258 m) but left to grow at its edge (November 2023). On the right *Nicotiana glauca*.



Figure 3. At Jira, *Salvia tiliifolia* grows densely along footpaths (A, 13°41'05" N, 38°57'27" W, 1991 m, photo Gebrekidan Mesfin), while it is totally absent from the adjacent enclosure (B, 13°40'47" N, 38°57'37" W, 1920 m, photo Miro Jacob). Both photos taken in October 2024.