INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology, CAB International Institute of Parasitology, 395A Hatfield Road, St Albans, Herts AL4 0XQ. England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size $12 \cdot 0 \times 20 \cdot 0$ cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Diagnostic M, 31 000 Schistosoma mansoni proteins: requirement of infection, but not immunization, and use of the "miniblot" technique for the production of monoclonal antibodies. A. RUPPEL, U. BRETERNITZ and R. BURGER	95–101
Onchocerciasis in British cattle: a study of <i>Onchocerca gutturosa</i> and <i>O. lienalis</i> in North Wales. A. J. TREES, P. J. McCALL and S. J. CROZIER	103-113
Inhibition phenomena mechanisms in experimental obeliscoidosis in rabbits. II. Local and systemic antibody responses. I. B. HELAL, H. WEDRYCHOWICZ, E. SINSKI and B. BEZUBIK	115–123
Cortisone-sensitive, innate resistance to <i>Hymenolepis nana</i> infection in congenitally athymic nude rats. A. ITO and T. KAMIYAMA	124–128
Changes in surface antigens of <i>Hymenolepis nana</i> during differentiation and maturation in mice. A. ITO and K. ONITAKE	129–136
The protective effect of irradiated metacercariae of Fasciola gigantica against homologous challenge in sheep. H. A/GADIR, E. M. HAROUN and A. A. GAMEEL	137–142
Polarographic determination of mitochondrial respiration in Gastrothylax crumenifer. (Research Note). J. SIDDIQUI and A. H. SIDDIQI	143–145
Purification and properties of haemoglobin from Gastrothylax crumenifer (Trematoda: Paramphistomatidae). J. SIDDIQUI and A. H. SIDDIQI	146–150
Immunodepression in hamsters experimentally infected with <i>Opisthorchis viverrini</i> . S. WONGRATANACHEEWIN, W. RATTANASIRIWILAI, R. PRIWAN and S. SIRISINHA	151–156
Eosinophilia in chimeric mice infected with <i>Toxocara canis</i> . K. SUGANE and T. MATSUURA	157–162
Development, growth and migration of <i>Philophthalmus gralli</i> (Trematoda) from Jordan in the eyes of chicks. N. S. ISMAIL and I. M. ISSA	163-168
The neural commissural rings of the adult male Schistosoma mansoni. B. WITHYACHUMNARNKUL, A. PONGSA-ASAWAPAIBOON, P. SOBHON and E. S. UPATHAM	169–172
Studies of the effect of contralateral versus ipselateral challenge of rats primed on one side with <i>Strongyloides ratti</i> . P. A. G. WILSON	173–178
The pathogenesis of benzimidazole-resistant and benzimidazole-susceptible strains of <i>Trichostrongylus colubriformis</i> in the Mongolian gerbil (<i>Meriones unguiculatus</i>). J. M. MACLEAN, D. LEWIS and P. H. HOLMES	179–189
Book review	114
Announcements	102, 190